

Energy-dependent dipole form factor in a QCD-inspired model

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 J. Phys.: Conf. Ser. 706 052006

(<http://iopscience.iop.org/1742-6596/706/5/052006>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 200.130.19.173

This content was downloaded on 24/05/2017 at 18:48

Please note that [terms and conditions apply](#).

You may also be interested in:

[Resonance measurements in pp collisions with the ALICE detector and prospects for Pb-Pb](#)

A Pulvirenti and the ALICE Collaboration

[Electromagnetic form factors of nucleons](#)

P N Pandit, M P Khanna and M Gupta

[Hadronic resonances at ALICE](#)

A G Knospe and the Alice Collaboration

[The deuteron wavefunction in the quark compound bag model](#)

I L Grach, Yu S Kalashnikova and I M Narodetskii

[A new molecular dynamics calculation for the light and strange baryons](#)

T Watanabe, M Oosawa, K Saito et al.

[System size dependence of hadron pT spectra in p + p and Au+Au collisions at \$s_{\sqrt{s_{NN}}}\$ GeV](#)

P K Khandai, P Sett, P Shukla et al.

[Bottomonium states versus recent experimental observations in the QCD-inspired potential model](#)

Tian Wei-Zhao, Cao Lu, Yang You-Chang et al.

[Inelastic cross sections, overlap functions and Cq moments from ISR to LHC energies in proton interactions](#)

P C Boggio

Energy-dependent dipole form factor in a QCD-inspired model

C.A.S. Bahia, M. Broilo, E.G.S. Luna

Instituto de Física, Universidade Federal do Rio Grande do Sul, Caixa Postal 15051,
91501-970, Porto Alegre, RS, Brazil

E-mail: andre.bahia@ufrgs.br, mateus.broilo@ufrgs.br, luna@if.ufrgs.br

Abstract. We consider the effect of an energy-dependent dipole form factor in the high-energy behavior of the forward amplitude. The connection between the semihard parton-level dynamics and the hadron-hadron scattering is established by an eikonal QCD-based model. Our results for the proton-proton (pp) and antiproton-proton ($\bar{p}p$) total cross sections, $\sigma_{tot}^{pp,\bar{p}p}(s)$, obtained using the CTEQ6L1 parton distribution function, are consistent with the recent data from the TOTEM experiment.

1. Introduction

The total cross section $\sigma_{tot}(s)$ is a fundamental quantity in collisions of strongly interacting particles. At present one of the main theoretical approaches for the description of σ_{tot} is the QCD-inspired formalism [1, 2, 3, 4]. In this approach the energy dependence of the total cross section is obtained from QCD via an eikonal formulation. The high-energy dependence of σ_{tot} is driven by the rapid increase in gluon density at small- x . In this work we explore the non-perturbative dynamics of QCD in order to describe, in both pp and $\bar{p}p$ channels, the total cross sections $\sigma_{tot}^{pp,\bar{p}p}(s)$ and the ratios of the real to imaginary part of the forward scattering amplitude, $\rho^{pp,\bar{p}p}(s)$. In our calculations we introduce an energy-dependent dipole form factor which represents the overlap density for the partons at impact parameter b . The behavior of these forward quantities is derived from the QCD parton model using standard parton-parton elementary processes and an updated set of gluon distribution, namely the CTEQ6L1 set. In our analysis we have included recent measurements of pp total cross sections at the LHC by the TOTEM Collaboration [5]. This work is organized as follows: in the next section we introduce our QCD-based model where the onset of the dominance of semihard partons is managed by the dynamical gluon mass. We investigate a form factor which the spatial distribution of gluons changes with the energy and introduce integral dispersion relations to connect the real and imaginary parts of eikonals. In Section III we present our results and in Section IV we draw our conclusions.

2. The QCD-inspired eikonal model

In our model the increase of $\sigma_{tot}^{pp,\bar{p}p}(s)$ is associated with elementary semihard processes in the hadrons. In the eikonal representation $\sigma_{tot}(s)$ and $\rho(s)$ are given by

$$\sigma_{tot}(s) = 4\pi \int_0^\infty b db [1 - e^{-\chi_R(s,b)} \cos \chi_I(s,b)] \quad (1)$$



and

$$\rho(s) = \frac{-\int_0^\infty b db e^{-\chi_R(s,b)} \sin \chi_I(s,b)}{\int_0^\infty b db [1 - e^{-\chi_R(s,b)} \cos \chi_I(s,b)]}, \quad (2)$$

respectively, where s is the square of the total center-of-mass energy, b is the impact parameter, and $\chi(s,b) = \text{Re} \chi(s,b) + i \text{Im} \chi(s,b) \equiv \chi_R(s,b) + i \chi_I(s,b)$ is the (complex) eikonal function. The eikonal functions for pp and $\bar{p}p$ scatterings are the sum of the soft and semihard (SH) parton interactions in the hadron-hadron collision,

$$\chi(s,b) = \chi_{soft}(s,b) + \chi_{SH}(s,b). \quad (3)$$

It follows from the QCD parton model that the real part of the eikonal, $\text{Re} \chi_{SH}(s,b)$, can be factored as

$$\text{Re} \chi_{SH}(s,b) = \frac{1}{2} W_{SH}(b) \sigma_{QCD}(s), \quad (4)$$

where $W_{SH}(b)$ is an overlap density for the partons at impact parameter space b ,

$$W_{SH}(b) = \int d^2b' \rho_A(|\mathbf{b} - \mathbf{b}'|) \rho_B(b'), \quad (5)$$

and $\sigma_{QCD}(s)$ is the usual QCD cross section

$$\begin{aligned} \sigma_{QCD}(s) &= \sum_{ij} \frac{1}{1 + \delta_{ij}} \int_0^1 dx_1 \int_0^1 dx_2 \int_{Q_{min}^2}^\infty d|\hat{t}| \frac{d\hat{\sigma}_{ij}}{d|\hat{t}|}(\hat{s}, \hat{t}) \\ &\times f_{i/A}(x_1, |\hat{t}|) f_{j/B}(x_2, |\hat{t}|) \Theta\left(\frac{\hat{s}}{2} - |\hat{t}|\right), \end{aligned} \quad (6)$$

with $|\hat{t}| \equiv Q^2$, $\hat{s} = x_1 x_2 s$ and $i, j = q, \bar{q}, g$.

The eikonal functions for pp and $\bar{p}p$ scatterings are written as $\chi_{pp}^{\bar{p}p}(s,b) = \chi^+(s,b) \pm \chi^-(s,b)$, with $\chi^+(s,b) = \chi_{soft}^+(s,b) + \chi_{SH}^+(s,b)$ and $\chi^-(s,b) = \chi_{soft}^-(s,b) + \chi_{SH}^-(s,b)$. Since in the parton model $\chi_{SH}^-(s,b)$ decreases rapidly with increasing s , the difference between pp and $\bar{p}p$ cross sections is due only to the different weighting of the quark-antiquark (valence) annihilation cross sections in the two channels. Thus the crossing-odd eikonal $\chi^-(s,b)$ receives no contribution from semihard processes at high energies, and it is sufficient to take $\chi_{SH}(s,b) = \chi_{SH}^+(s,b)$ and, consequently, $\chi^-(s,b) = \chi_{soft}^-(s,b)$. The connection between the real and imaginary parts of the crossing-even eikonal $\chi^+(s,b)$ is obtained by means of dispersion relation

$$\text{Im} \chi^+(s,b) = -\frac{2s}{\pi} \mathcal{P} \int_0^\infty ds' \frac{\text{Re} \chi^+(s',b)}{s'^2 - s^2}, \quad (7)$$

valid at $s \gg 1 \text{ GeV}^2$. The soft term is given by

$$\chi_{soft}^+(s,b) = \frac{1}{2} W_{soft}^+(b; \mu_{soft}^+) \left[A' + \frac{B'}{(s/s_0)^\gamma} e^{i\pi\gamma/2} - iC' \left[\ln\left(\frac{s}{s_0}\right) - i\frac{\pi}{2} \right] \right], \quad (8)$$

where $\sqrt{s_0} \equiv 5 \text{ GeV}$ and A', B', C', γ and μ_{soft}^+ are fitting parameters. The odd eikonal $\chi^-(s,b)$ is given by

$$\chi^-(s,b) = \frac{1}{2} W_{soft}^-(b; \mu_{soft}^-) D' \frac{e^{-i\pi/4}}{\sqrt{s}}, \quad (9)$$

where $\mu_{soft}^- \equiv 0.5$ GeV and D' is also a fitting parameter.

In the expression (6) the term $d\hat{\sigma}_{ij}/d\hat{t}$ is the differential cross section for ij scattering ($i, j = q, \bar{q}, g$). We select parton-parton processes containing at least one gluon in the initial state, i.e., we consider the processes

$$\frac{d\hat{\sigma}}{d\hat{t}}(gg \rightarrow gg) = \frac{9\pi\bar{\alpha}_s^2}{2\hat{s}^2} \left(3 - \frac{\hat{t}\hat{u}}{\hat{s}^2} - \frac{\hat{s}\hat{u}}{\hat{t}^2} - \frac{\hat{t}\hat{s}}{\hat{u}^2} \right), \quad (10)$$

$$\frac{d\hat{\sigma}}{d\hat{t}}(qg \rightarrow qg) = \frac{\pi\bar{\alpha}_s^2}{\hat{s}^2} (\hat{s}^2 + \hat{u}^2) \left(\frac{1}{\hat{t}^2} - \frac{4}{9\hat{s}\hat{u}} \right), \quad (11)$$

$$\frac{d\hat{\sigma}}{d\hat{t}}(gg \rightarrow \bar{q}q) = \frac{3\pi\bar{\alpha}_s^2}{8\hat{s}^2} (\hat{t}^2 + \hat{u}^2) \left(\frac{4}{9\hat{t}\hat{u}} - \frac{1}{\hat{s}^2} \right), \quad (12)$$

where

$$\bar{\alpha}_s(Q^2) = \frac{4\pi}{\beta_0 \ln \left[(Q^2 + 4M_g^2(Q^2))/\Lambda^2 \right]}, \quad (13)$$

$$M_g^2(Q^2) = m_g^2 \left[\frac{\ln \left(\frac{Q^2 + 4m_g^2}{\Lambda^2} \right)}{\ln \left(\frac{4m_g^2}{\Lambda^2} \right)} \right]^{-12/11}; \quad (14)$$

here $\bar{\alpha}_s(Q^2)$ is a non-perturbative QCD effective charge, obtained by Cornwall through the use of the pinch technique in order to derive a gauge invariant Schwinger-Dyson equation for the gluon propagator [6].

In the case of semihard gluons we consider the possibility of a ‘‘broadening’’ of the spatial distribution. Our assumption suggests an increase of the average gluon radius when \sqrt{s} increases, and can be properly implemented using the energy-dependent dipole form factor

$$G_{SH}^{(d)}(s, k_{\perp}; \nu_{SH}) = \left(\frac{\nu_{SH}^2}{k_{\perp}^2 + \nu_{SH}^2} \right)^2, \quad (15)$$

where $\nu_{SH} = \nu_1 - \nu_2 \ln(\frac{s}{s_0})$, with $\sqrt{s_0} \equiv 5$ GeV. Here ν_1 and ν_2 are constants to be fitted. From equation (5) we have

$$\begin{aligned} W_{SH}(s, b; \nu_{SH}) &= \frac{1}{2\pi} \int_0^{\infty} dk_{\perp} k_{\perp} J_0(k_{\perp} b) [G_{SH}^{(d)}(s, k_{\perp}; \nu_{SH})]^2 \\ &= \frac{\nu_{SH}^2}{96\pi} (\nu_{SH} b)^3 K_3(\nu_{SH} b). \end{aligned} \quad (16)$$

Since semihard interactions dominate at high energies, we consider an energy-dependence only in the case of $W_{SH}(s, b; \nu_{SH})$. In this way the soft overlap densities $W_{soft}^+(b; \mu_{soft}^+)$ and $W_{soft}^-(b; \mu_{soft}^-)$ are static, i.e., μ_{soft}^+ and μ_{soft}^- are not energy dependent.

3. Results

First, in order to determine the model parameters, we set the value of the gluon mass scale to $m_g = 400$ MeV. This choice for the mass scale is not only consistent to our LO procedure, but is also the one usually obtained in other calculations of strongly interacting processes [2, 3, 7, 8]. Next, we carry out a global fit to high-energy forward pp and $\bar{p}p$ scattering data above $\sqrt{s} = 10$ GeV, namely the total cross section $\sigma_{tot}^{pp,\bar{p}p}$ and the ratio of the real to imaginary part of the forward scattering amplitude $\rho^{pp,\bar{p}p}$. We use data sets compiled and analyzed by the Particle Data Group [9] as well as the recent data at LHC from the TOTEM Collaboration, with the statistic and systematic errors added in quadrature. We include in the dataset the first estimate for the ρ parameter made by the TOTEM Collaboration in their ρ -independent measurement at $\sqrt{s} = 7$ TeV, namely $\rho^{pp} = 0.145 \pm 0.091$ [5].

The values of the fitted parameters are given in the legend of Figure 1. The result $\chi^2/DOF = 1.062$ for the fit was obtained for 154 degrees of freedom. The results of the fit to σ_{tot} and ρ for both pp and $\bar{p}p$ channels are displayed in Figure 1, together with the experimental data. The curves depicted in Figure 1 were all calculated using the cutoff $Q_{min} = 1.3$ GeV, the value of the CTEQ6 fixed initial scale Q_0 .




Figure 1. The total cross section (upper) and the ratio of the real to imaginary part of the forward scattering amplitude (lower), for pp (\bullet) and $\bar{p}p$ (\circ). The values of the fitted parameters are: $\nu_1 = 2.770 \pm 0.865$ GeV, $\nu_2 = (7.860 \pm 5.444) \times 10^{-2}$ GeV, $A' = 108.9 \pm 8.6$ GeV $^{-1}$, $B' = 30.19 \pm 15.78$ GeV $^{-1}$, $C' = 1.260 \pm 0.437$ GeV $^{-1}$, $\gamma = 0.719 \pm 0.200$, $\mu_{soft}^+ = 0.457 \pm 0.209$ GeV and $D' = 21.73 \pm 2.26$ GeV $^{-1}$.

4. Conclusions

In this work we have investigated infrared effects on semihard parton-parton interactions. We have studied pp and $\bar{p}p$ scattering in the LHC energy region with the assumption that the increase of their total cross sections arises exclusively from semihard interactions. In the calculation of $\sigma_{tot}^{pp,\bar{p}p}$ and $\rho^{pp,\bar{p}p}$ we have considered the phenomenological implication of an energy-dependent form factor for semihard partons. We introduce integral dispersion relations to connect the real and imaginary parts of eikonals with energy-dependent form factors. In our analysis we have included the recent data at LHC from the TOTEM Collaboration. Our results show that very good descriptions of $\sigma_{tot}^{pp,\bar{p}p}$ and $\rho^{pp,\bar{p}p}$ data are obtained by constraining the value of the cutoff at $Q_{min} = 1.3$ GeV. The χ^2/DOF for the global fit was 1.062 for 154 degrees of freedom. This good statistical result shows that our eikonal model, where non-perturbative effects are naturally included via a QCD effective charge, is well suited for detailed predictions of the forward quantities to be measured at higher energies. In the semihard sector we have considered a form factor in which the average gluon radius increases with \sqrt{s} . With this assumption we have obtained another form in which the eikonal can be factored into the QCD parton model, namely $\text{Re } \chi_{SH}(s, b) = \frac{1}{2} W_{SH}(s, b) \text{Re } \sigma_{QCD}(s)$. The imaginary part of this *semi-factorizable* eikonal was obtained by means of appropriate integral dispersion relations which take into account eikonals with energy-dependent form factors.

At the moment we are analyzing the effects of different updated sets of parton distributions on the forward quantities, namely CTEQ6L and MSTW, investigating the uncertainty on these forward observables coming from the uncertainties associated with the dynamical mass scale and the parton distribution functions, and exploring a new ansatz for the energy-dependent form factor.

References

- [1] L. Durand and H. Pi, Phys. Rev. Lett. **58**, 303 (1987); Phys. Rev. D **38**, 78 (1988); **40**, 1436 (1989).
- [2] E.G.S. Luna, A.F. Martini, M.J. Menon, A. Mihara, and A.A. Natale, Phys. Rev. D **72**, 034019 (2005); E.G.S. Luna, Phys. Lett. B **641**, 171 (2006); E.G.S. Luna and A.A. Natale, Phys. Rev. D **73**, 074019 (2006); E.G.S. Luna, Braz. J. Phys. **37**, 84 (2007); D.A. Fagundes, E.G.S. Luna, M.J. Menon, and A.A. Natale, Nucl. Phys. A **886**, 48 (2012); E.G.S. Luna and P.C. Beggio, Nucl. Phys. A **929**, 230 (2014).
- [3] E.G.S. Luna, A.L. dos Santos, and A.A. Natale, Phys. Lett. B **698**, 52 (2011).
- [4] A. Corsetti, A. Grau, G. Pancheri, and Y.N. Srivastava, Phys. Lett. B **382**, 282 (1996); A. Grau, G. Pancheri, and Y.N. Srivastava, Phys. Rev. D **60**, 114020 (1999); R.M. Godbole, A. Grau, G. Pancheri, and Y.N. Srivastava, Phys. Rev. D **72**, 076001 (2005); A. Achilli *et al.*, Phys. Lett. B **659**, 137(2008); A. Grau, R.M. Godbole, G. Pancheri, and Y.N. Srivastava, Phys. Lett. B **682**, 55 (2009); G. Pancheri, D.A. Fagundes, A. Grau, S. Pacetti, and Y.N. Srivastava, arXiv:1301.2925 [hep-ph].
- [5] G. Antchev *et al.*, EPL **96**, 21002 (2011); G. Antchev *et al.*, EPL **101**, 21002 (2013); G. Antchev *et al.*, EPL **101**, 21004 (2013); G. Antchev *et al.*, Phys. Rev. Lett. **111**, 012001 (2013).
- [6] J. M. Cornwall, Phys. Rev. D **22**, 1452 (1980); D **26**, 1453 (1982).
- [7] A. Doff, E.G.S. Luna, and A.A. Natale, Phys. Rev. D **88**, 055008 (2013).
- [8] E.G.S. Luna and A.A. Natale, J. Phys. G **42**, 015003 (2015).
- [9] K.A. Olive *et al.*, Chin. Phys. C **38**, 090001 (2014).