

UNIVERSIDADE FEDERAL DO RIO GRANDE DO SUL
INSTITUTO DE MATEMÁTICA
CADERNOS DE MATEMÁTICA E ESTATÍSTICA
SÉRIE A: TRABALHO DE PESQUISA

SPECTRAL ANALYSIS OF EXPANDING
ONE-DIMENSIONAL CHAOTIC TRANSFORMATIONS

ARTUR OSCAR LOPES
RAFAEL RIGÃO SOUZA
SILVIA REGINA COSTA LOPES

SÉRIE A, N° 46
PORTO ALEGRE, ABRIL DE 1996

SPECTRAL ANALYSIS OF EXPANDING ONE-DIMENSIONAL CHAOTIC TRANSFORMATIONS

Artur Lopes, Sílvia Lopes and Rafael Souza

Instituto de Matemática
Universidade Federal do Rio Grande do Sul
Av. Bento Gonçalves, 9500
Porto Alegre – RS – 91540-000 – Brasil

SUMMARY

The purpose of this paper is to show explicitly the spectral distribution function of some stationary stochastic processes as

$$X_t = F(X_{t-1}), \quad \text{for } t \in \mathbb{Z},$$

where F is a deterministic two-dimensional invertible map. The invertible map F that will be considered in this paper is the natural extension of a map T on a class \mathcal{F}_2 (see Section 1 for definition) of one-dimensional piecewise linear expanding monotonic transformations.

Any non-linear expanding piecewise monotonic transformation $g \in \mathcal{F}_1$ (see Section 1 for definition) can be approximated by a map $T \in \mathcal{F}_2$. From the structural stability of the maps we consider here, it will follow that the spectral density function of the natural extension of any non-linear expanding piecewise monotonic transformation $g \in \mathcal{F}_1$ can be approximated by explicit expressions obtained for the spectral density function of the natural extension of maps T in \mathcal{F}_2 .

Results for the one-dimensional map T can be obtained from results for the two-dimensional map F .

Keywords: CHAOTIC TIME SERIES; SPECTRAL ANALYSIS; EXPANDING TRANSFORMATION.

1. INTRODUCTION

We shall consider a special class of *non-linear piecewise monotonic expanding C^2 transformations* g in which the image of any interval of monotonicity is all the interval $(0, 1)$ (see Lasota and Mackey (1994) or Section 3 for definition). For instance, Figure 1 shows an example of a map g of such class while Figure 2 shows the graph of a map that is not of the above defined class. The number of intervals of monotonicity of g will be assumed to be finite.

We will denote the set of such class of maps g by \mathcal{F}_1 .

Consider the class of piecewise linear monotonic continuous expanding transformations T of the following form.

Let A_i , $1 \leq i \leq n$, be an open interval and a_i be the length of A_i , $1 \leq i \leq n$, the intervals of monotonicity (we assume that $T(A_i) = (0, 1)$, for $1 \leq i \leq n$), and suppose that $a_i = \sum_{j=1}^m b_{ij}$. Denote by B_{ij} , for $1 \leq i \leq n$ and $1 \leq j \leq m$, intervals such that the length of B_{ij} is b_{ij} and $\cup_{j=1}^m B_{ij} = A_i$, for all $1 \leq i \leq n$. Suppose there exists C_j , for $1 \leq j \leq m$, such that $g(B_{ij}) = C_j$, independent of i . Finally, assume that each C_j is contained in a unique A_i and each C_j is a union of sets of the form B_{ul} . Denote by c_j the length of C_j , for $1 \leq j \leq m$.

The analytic expression of $T(x)$ is given by

$$T(x) = \sum_{k=0}^{j-1} c_k + \left(x - \sum_{\alpha=1}^{i-1} a_\alpha - \sum_{\beta=1}^{j-1} b_{i\beta} \right) \frac{c_j}{b_{ij}}.$$

We will denote the set of this second class of maps T by \mathcal{F}_2 .

In Góra and Boyarsky (1989) and Parry and Pollicott (1990) the explicit expression of the invariant density

$$\sum_{j=1}^m I_{C_j}(x)p_j, \quad p_j \in (0, 1), \quad \text{with} \quad \sum_{j=1}^m p_j = 1$$

of $T \in \mathcal{F}_2$ is obtained by finding the eigenvector (p_1, p_2, \dots, p_m) of a large matrix. In this way the number p_j , for $1 \leq j \leq m$, can be explicitly obtained.

It is easy to see (we refer the reader to Góra and Boyarsky (1989)) that each $g \in \mathcal{F}_1$ can be C^1 approximated by $T \in \mathcal{F}_2$ (up to a finite number of points where T is not differentiable). In Figure 3, we show the graph of a map $T \in \mathcal{F}_2$.

We will show an explicit formula (see Section 4) for the spectral density function of the natural extension of the piecewise linear map $T \in \mathcal{F}_2$ described above.

Góra and Boyarsky (1989), Li (1976), Ding and Li (1991) and also Parry and Pollicott (1990) show that the invariant density $\eta_n(x)$ of a sequence of maps $T_n \in \mathcal{F}_2$ converging to $g \in \mathcal{F}_1$ satisfies the weak convergence

$$\eta_n(x) \rightarrow \eta(x),$$

where $\eta(x)$ is the invariant density for the map g .

Therefore, one can obtain an approximation of the spectral density function of $g \in \mathcal{F}_1$ by an explicit formula for the spectral density function of $T_n \in \mathcal{F}_2$.

The spectral density function of such map $g \in \mathcal{F}_1$ will be a meromorphic function (see Ruelle (1987)).

The spectral density function of maps g of the class \mathcal{F}_1 are important for the spectral analysis of chaotic time series and also because the zeta function associated with the potential $-\log g'(x)$ has poles on the same values of the poles of the spectral density function (see Ruelle (1978, 1987) and Rugh (1992)).

2. STATIONARY STOCHASTIC PROCESSES

The general setting of chaotic time series we shall analyze is the following. Consider K a compact subset of \mathbf{R}^n with a given Borel σ -algebra \mathcal{F} , a bijective continuous transformation $F : K \rightarrow K$, an invariant probability \mathcal{P} on K (that is, $\mathcal{P}(F^{-1}(A)) = \mathcal{P}(A)$, for any set $A \in \mathcal{F}$) and $\phi : K \rightarrow \mathbf{R}$ a continuous function. We will analyze the stationary stochastic process $\{Z_t\}_{t \in \mathbf{Z}}$ given by

$$Z_t = X_t + \xi_t = (\phi \circ F)(X_{t-1}) + \xi_t, \quad \text{for } t \in \mathbf{Z}. \quad (2.1)$$

The natural measure on $K^{\mathbf{Z}}$ is the product measure $\mathcal{P}^{\mathbf{Z}}$ on $K^{\mathbf{Z}}$ and it is invariant for the stationary process $\{X_t\}_{t \in \mathbf{Z}}$ or $\{Z_t\}_{t \in \mathbf{Z}}$. The process $\{\xi_t\}_{t \in \mathbf{Z}}$ is considered to be a Gaussian white noise process (see Brockwell and Davis (1987)) independent of $\{(\phi \circ F)(X_t)\}_{t \in \mathbf{Z}}$, with zero mean and variance σ_ξ^2 . One observes that in the model (2.1) the random variables X_t (or Z_t) and X_{t+1} (or Z_{t+1}) are generally not independent.

We shall denote the above system by

$$(K, F, \mathcal{P}, \phi, \mathcal{F}, \sigma_\xi^2). \quad (2.2)$$

Following the terminology in Tong (1990) we may call the system (2.1), when $\sigma_\xi^2 = 0$, the *skeleton* of the system.

Given a certain measurable function $\phi : K \rightarrow \mathbf{R}$ the *autocovariance function at lag* $h \in \mathbf{Z}$ (see Brockwell and Davis (1987)) of the process $\{X_t\}_{t \in \mathbf{Z}}$ as in (2.1) is given by

$$R_{XX}(h) = E(X_t X_{t+h}) - [E(X_t)]^2 = \int \phi(x)\phi(F^h(x))d\mathcal{P}(x) - \left[\int \phi(x)d\mathcal{P}(x) \right]^2. \quad (2.3)$$

The autocovariance function $R_{XX}(h)$ in (2.3) measures the covariance between two values of the process $\{X_t\}_{t \in \mathbf{Z}}$ separated by lag h . The *autocorrelation function at lag* h of the process $\{X_t\}_{t \in \mathbf{Z}}$ (see Brockwell and Davis (1987)) is given by

$$\rho_X(h) = \frac{R_{XX}(h)}{R_{XX}(0)}, \quad \text{for } h \in \mathbf{Z}, \quad (2.4)$$

where $R_{XX}(0) = E[(X_t - E(X_t))^2] = \text{Var}(X_t)$ is the variance of the process.

From the Herglotz's theorem (see Brockwell and Davis (1987)) a function $\rho_X(h)$ is non-negative definite if and only if

$$\rho_X(h) = \int_{-\pi}^{\pi} e^{i\lambda h} dF_X(\lambda), \quad \text{for any } h \in \mathbf{Z}, \quad (2.5)$$

where $F_X(\cdot)$ is a right-continuous, non-decreasing, bounded function on $[-\pi, \pi]$ with $F_X(-\pi) = 0$. The function $F_X(\cdot)$ is called the *spectral distribution function* of $\{X_t\}_{t \in \mathbf{Z}}$ and if

$$F_X(\lambda) = \int_{-\pi}^{\lambda} f_X(\omega) d\omega, \quad \text{for } -\pi < \lambda \leq \pi, \quad (2.6)$$

then $f_X(\cdot)$ is called *the spectral density function* of the process $\{X_t\}_{t \in \mathbf{Z}}$. When

$$\sum_{h=-\infty}^{\infty} |\rho_X(h)| < \infty,$$

then $\rho_X(h) = \int_{-\pi}^{\pi} e^{ih\lambda} f_X(\lambda) d\lambda$, for $h \in \mathbf{Z}$, where $f_X(\cdot)$ is given by

$$f_X(\lambda) = \frac{1}{2\pi} \sum_{h=-\infty}^{\infty} e^{-ih\lambda} \rho_X(h). \quad (2.7)$$

This function has real values if $\rho_X(h) = \rho_X(-h)$, for all $h \in \mathbf{N}$.

The reason to consider F a bijective map and not just a non-invertible map is for defining $R_{XX}(h)$ also for negative values of $h \in \mathbf{Z}$ and, from this, (2.7) will be well defined.

Each particular invertible transformation F will require different technique in order to obtain explicitly the spectral distribution function (see Lopes and Lopes (1995)).

Example: When the compact subset K is equal to $[-\pi, \pi]$, the transformation F is given by $F(x) = \omega_0 + x \pmod{2\pi}$, with $\omega_0 \in (0, \pi)$, and $\phi(x) = \cos(x)$. This is the classical harmonic model $Z_t = \cos(\omega_0 t + x) + \xi_t$. The spectral measure of the process $\{X_t\}_{t \in \mathbf{Z}} = \{(\phi \circ F)(X_{t-1})\}_{t \in \mathbf{Z}}$ as in (2.1) is not a function but a *distribution function* given by

$$dF_X(\lambda) = \frac{1}{2}(\delta_{\omega_0} + \delta_{-\omega_0}), \quad (2.8)$$

where δ_{ω_0} is the Dirac delta function concentrated at ω_0 .

Remark: Expanding maps (see Section 3 for the definition) always have an exponential decay of autocorrelations, for any ϕ Holder continuous function (see Parry and Pollicott (1990)). Therefore, in this case the spectral density function always exists and it is a meromorphic function (see Ruelle (1978, 1987)).

3. THE NATURAL EXTENSION F OF T

It is well known that in general larger the dimension of the set K , more difficult is to analyze the dynamics of the map F .

When K is one-dimensional, that is, when K is a segment, the diffeomorphism $F : K \rightarrow K$ has simple dynamics.

In general, the dynamics of an one-dimensional diffeomorphism is very simple.

The simplest example in dimension 2, that is, when K is a square $[0, 1] \times [0, 1]$, is obtained when F is the natural extension of an one-dimensional map T . The map T is not an one-to-one map, but F is.

When the transformation T is an *expanding map*, that is, there exists $\lambda > 1$ such that $|T'(x)| > \lambda$, for all $x \in [0, 1]$, then there exists (see Lasota and Yorke (1973) and also Parry and Pollicott (1990)) a density $\eta(x)$ such that $d\mu(x) = \eta(x) dx$ is invariant for T (that is,

$\mu(T^{-1}(A)) = \mu(A)$, for any Borel set A). The probability μ is *ergodic* (see Cornfeld, et al. (1982) for the definition) for such map T . There exists a natural way to obtain from such T a bijective map F , called *the natural extension of T* . Denote by (x, y) a vector in the domain K and by $(x', y') = F(x, y)$ its image by the map F . Then, (see Bogomolny and Carioli (1995))

$$T(x) = x' \quad \text{and} \quad T(y') = y$$

defines F .

If T is an expanding map the corresponding F is *Axiom A* (see Robinson (1995) for definition).

The invariant probability μ for T on $[0, 1]$ has a natural extension to a probability $\nu = \mathcal{P}$ (according to the notation of Section 2) on $K = [0, 1] \times [0, 1]$ invariant for F .

Consider now the random variable $\phi : K \rightarrow \mathbf{R}$ of the form $\phi(x, y) = \phi(x)$. Then, the time series

$$X_t = \phi(F^t(x, y)) = \phi(T^t(x)), \quad \text{for } 1 \leq t \leq N,$$

and the probability ν define the simplest example of a chaotic time series on dimension 2.

The dynamics comes basically from an one-dimensional map even if the setting is for a two-dimensional bijective map. As we mentioned before the reason to consider bijective maps is to obtain $R_{XX}(h)$, for $h \in \mathbf{Z}$.

The analysis of the dynamics of F is more general and results for T can be derived from the former transformation.

When $\phi(x, y) = x$, for a certain class of maps T in \mathcal{F}_2 (see Section 4) we shall be able to show explicitly the spectral density function. This is obtained by solving some linear systems as we shall explain later. We call a stochastic process obtained from the system (F, ϕ) as above *a standard stochastic process* obtained from (T, ϕ) .

Any expanding map g in \mathcal{F}_1 can be approximated by maps T in \mathcal{F}_2 and the corresponding absolutely continuous invariant measure of T will converge to the corresponding one for g (see Góra and Boyarsky (1989)). Therefore, we will be able to approximate the spectral density function of expanding maps by known expressions. This is the main purpose of this paper.

In the sequel, we shall omit the noise process $\{\xi_t\}_{t \in \mathbf{Z}}$ of the system due to the fact that it does not interfere in the dynamics of T and that the spectral density function of the whole system with noise can be easily obtained from the one without noise (see Lopes et al. (1995, 1996)).

4. THE SPECTRAL DENSITY OF PIECEWISE LINEAR EXPANDING TRANSFORMATIONS

In this section we will show the explicit expression of the spectral density function of the system $X_t = T^t(X_0)$, where $T \in \mathcal{F}_2$.

We will denote by μ the T -invariant measure absolutely continuous with respect to the Lebesgue measure. Denote by $p(x)$ the density of such measure μ . It is well known

(see Parry and Pollicott (1990)) that $p(x)$ is of the form

$$p(x) = \sum_{j=1}^m I_{C_j}(x) p_j.$$

Therefore, $\mu(C_j) = \int_{C_j} p(x) dx = p_j$, for $1 \leq j \leq m$. The number p_j , for $1 \leq j \leq m$, can be obtained by finding an eigenvector of a large matrix (see Góra and Boyarsky (1989) and Li (1976)). Denote also by p_{ij} the measure μ of the interval B_{ij} , $1 \leq i \leq n$ and $1 \leq j \leq m$.

It is enough to show the explicit expression of

$$\gamma(z) = \sum_{k=0}^{\infty} \left(\int_0^1 x T^k(x) p(x) dx \right) z^k$$

and the corresponding explicit expression for the spectral density function (2.7) will easily follow (see Lopes et al. (1996)).

In the sequel, we shall consider the following notation:

$$A(k, i, j) = \int_{B_{ij}} x T^k(x) p(x) dx, \quad (4.1)$$

$$V(k, i, j) = \int_{B_{ij}} T^k(x) dx, \quad (4.2)$$

$$B(k, i, j) = \int_{B_{ij}} x T^k(x) dx \quad (4.3)$$

and

$$A(k) = \int_0^1 x T^k(x) p(x) dx. \quad (4.4)$$

First of all we shall compute a recursive formula for $V(k, i, j)$. One observes, from the expression (4.2), that

$$\begin{aligned} V(k+1, r, s) &= \int_{B_{rs}} T^{k+1}(x) dx = \int_{B_{rs}} T^k(T(x)) dx = \\ &= \int_{C_s} T^k(y) dy \frac{b_{rs}}{c_s} = \sum_{B_{uv} \subset C_s} \left(\int_{B_{uv}} T^k(y) dy \right) \frac{b_{rs}}{c_s} = \frac{b_{rs}}{c_s} \sum_{B_{uv} \subset C_s} V(k, u, v). \end{aligned} \quad (4.5)$$

Now we shall obtain the recursive formula for $B(k, i, j)$. One observes, from the expression (4.3), that

$$B(k+1, r, s) = \int_{B_{rs}} x T^{k+1}(x) dx = \int_{B_{rs}} x T^k(T(x)) dx =$$

$$\begin{aligned}
&= \int_{B_{r,s}} x T^k \left[\sum_{k=0}^{s-1} c_k + \left(x - \sum_{\alpha=1}^{r-1} a_\alpha - \sum_{\beta=1}^{s-1} b_{r\beta} \right) \frac{c_s}{b_{rs}} \right] dx = \\
&= \int_{C_s} \left[\left(y - \sum_{k=0}^{s-1} c_k \right) \frac{b_{rs}}{c_s} + \sum_{\alpha=1}^{r-1} a_\alpha + \sum_{\beta=1}^{s-1} b_{r\beta} \right] T^k(y) \frac{b_{rs}}{c_s} dy = \\
&= \sum_{B_{l,s} \subset C_s} \frac{b_{rs}}{c_s} \left[\int_{B_{l,s}} y T^k(y) dy \frac{b_{rs}}{c_s} + \right. \\
&\quad \left. + \left(- \sum_{k=0}^{s-1} c_k \frac{b_{rs}}{c_s} + \sum_{\alpha=1}^{r-1} a_\alpha + \sum_{\beta=1}^{s-1} b_{r\beta} \right) \int_{B_{l,s}} T^k(y) dy \right] = \\
&= \sum_{B_{l,s} \subset C_s} \frac{b_{rs}}{c_s} \left[B(k, l, s) \frac{b_{rs}}{c_s} + \left(- \sum_{k=0}^{s-1} c_k \frac{b_{rs}}{c_s} + \sum_{\alpha=1}^{r-1} a_\alpha + \sum_{\beta=1}^{s-1} b_{r\beta} \right) V(k, l, s) \right]. \quad (4.6)
\end{aligned}$$

We denote $\psi_{ij}(z)$ by

$$\psi_{ij}(z) = \sum_{k \geq 0} V(k, i, j) z^k = \sum_{k \geq 0} \left(\int_{B_{ij}} T^k(x) dx \right) z^k \quad (4.7)$$

and $\varphi_{ij}(z)$ by

$$\varphi_{ij}(z) = \sum_{k \geq 0} B(k, i, j) z^k = \sum_{k \geq 0} \left(\int_{B_{ij}} x T^{k+1}(x) dx \right) z^k. \quad (4.8)$$

Our purpose is to estimate

$$\gamma(z) = \sum_{k \geq 0} A_k z^k = \sum_{k \geq 0} \left(\int x T^k(x) p(x) dx \right) z^k, \quad (4.9)$$

but first we need to estimate $\psi_{ij}(z)$.

From (4.5), the power series of $V(k, i, j)$ satisfies the following equation

$$\begin{aligned}
\psi_{ij}(z) &= \sum_{k \geq 0} V(k, i, j) z^k = V(0, i, j) + \sum_{k \geq 0} V(k+1, i, j) z^{k+1} = \\
&= V(0, i, j) + z \sum_{k \geq 0} V(k+1, i, j) z^k = \\
&= V(0, i, j) + z \sum_{k \geq 0} \left(\frac{b_{ij}}{c_j} \sum_{B_{uv} \subset C_j} V(k, u, v) \right) z^k =
\end{aligned}$$

$$\begin{aligned}
&= V(0, i, j) + z \frac{b_{ij}}{c_j} \sum_{B_{uv} \subset C_j} \left(\sum_{k \geq 0} V(k, u, v) z^k \right) = \\
&= V(0, i, j) + z \frac{b_{ij}}{c_j} \sum_{B_{uv} \subset C_j} \psi_{u,v}(z).
\end{aligned}$$

Therefore, one can estimate $\psi_{i,j}(z)$ by solving the linear system

$$\psi_{ij}(z) = V(0, i, j) + z \frac{b_{ij}}{c_j} \sum_{B_{uv} \subset C_j} \psi_{u,v}(z). \quad (4.10)$$

Finally, the power series of $B(k, i, j)$ satisfies the equation

$$\begin{aligned}
\varphi_{i,j}(z) &= \sum_{k \geq 0} B(k, i, j) z^k = B(0, i, j) + z \sum_{k \geq 0} B(k+1, i, j) z^k = \\
&= B(0, i, j) + \\
&+ z \sum_{k \geq 0} z^k \left(\sum_{B_{lj} \subset C_j} \frac{b_{ij}}{c_j} \left[B(k, l, j) \frac{b_{ij}}{c_j} + \left(- \sum_{k=0}^{j-1} c_k \frac{b_{ij}}{c_j} + \sum_{\alpha=1}^{i-1} a_\alpha + \sum_{\beta=1}^{j-1} b_{i\beta} \right) V(k, l, j) \right] \right) = \\
&= B(0, i, j) + z \left[\sum_{B_{lj} \subset C_j} \left(\frac{b_{ij}}{c_j} \right)^2 \varphi_{lj}(z) + \right. \\
&\left. + \sum_{B_{lj} \subset C_j} \left(\frac{b_{ij}}{c_j} \right)^2 \left(- \sum_{k=0}^{j-1} c_k \frac{b_{ij}}{c_j} + \sum_{\alpha=1}^{i-1} a_\alpha + \sum_{\beta=1}^{j-1} b_{i\beta} \right) \psi_{l,j}(z) \right]. \quad (4.11)
\end{aligned}$$

As we know the values $\psi_{ij}(z)$ from (4.10), one can obtain $\varphi_{i,j}(z)$ from the linear equation (4.11).

Finally, we obtain $\gamma(z)$ explicitly by

$$\begin{aligned}
\gamma(z) &= \sum_{k \geq 0} A_k z^k = \sum_{k \geq 0} \left(\int_0^1 x T^k(x) p(x) dx \right) z^k \\
&= \sum_{i,j} \sum_{k \geq 0} \left(\int_{B_{ij}} x T^k(x) p(x) dx \right) z^k = \\
&= \sum_{i,j} p_{ij} \sum_{k \geq 0} \left(\int_{B_{ij}} x T^k(x) dx \right) z^k = \sum_{i,j} p_{ij} \varphi_{i,j}(z). \quad (4.12)
\end{aligned}$$

The spectral density function of X_t is given by

$$f_X(\lambda) = \frac{1}{2\pi \text{Var}(X_t)} [\gamma(e^{i\lambda}) + \gamma(e^{-i\lambda}) - E(X_t^2)], \quad \text{for any } \lambda \in (-\pi, \pi), \quad (4.13)$$

where $Var(X_t) = E(X_t^2) - [E(X_t)]^2$ and $\gamma(z)$ is given by the expression (4.12).

Remark: The power series $\gamma(z)$ is an analytic function on the unit disc $\{z \in \mathbf{C} \mid \|z\| < 1\}$ and the expression (4.13) has the meaning of the radial limit

$$\lim_{r \rightarrow 1} r e^{i\lambda} = e^{i\lambda} = z.$$

In this sense, the series

$$\sum_{n \in \mathbf{Z}} e^{in\lambda} = 2 \operatorname{Re} \left(\frac{1}{1 - e^{i\lambda}} \right) - 1 = 0, \quad \text{for } \lambda \neq 0,$$

even though the series $\sum_{n \in \mathbf{Z}} e^{in\lambda}$ does not converge. We are using this fact in the expression (4.13) above.

REFERENCES

1. Bogomolny, E. and Carioli, M. (1995) Quantum Maps of Geodesic Flows on Surfaces of Constant Negative Curvature. To appear in *Physica D*.
2. Brockwell, P. J. and Davis, R. A. (1991) *Time Series: Theory and Methods*, 2nd. edn. New York: Springer-Verlag.
3. Cornfeld, I. P., Fomin, S. V. and Sinai, Ya. G. (1982) *Ergodic Theory*. New York: Springer-Verlag.
4. Ding, J. and Li, T-Y. (1991) Markov Finite Approximation of Frobenius-Perron Operator. *Nonlinear Analysis, Theory, Methods & Applications*, **17**, No. 8, 759-772.
5. Góra, P. and Boyarsky, A. (1989) Compactness of Invariant Densities for Families of Expanding, Piecewise Monotonic Transformations. *Canadian Journal Mathematics*, **41**, No. 5, 855-869.
6. Lasota, A. and Yorke, J. A. (1973) On the existence of invariant measures for piecewise monotonic transformations. *Transactions of the American Mathematical Society*, **186**, 481-488.
7. Lasota, A. and Mackey, M. C. (1994) *Chaos, Fractals and Noise*. New York: Springer-Verlag.
8. Li, T-Y. (1976) Finite Approximation for the Frobenius-Perron Operator. A Solution to Ulam's Conjecture. *Journal of Approximation Theory*, **17**, No. 2, 177-186.

9. Lopes, A. and Lopes, S. (1995) Parametric Estimation and Spectral Analysis of Chaotic Time Series. Submitted.
10. Lopes, A., Lopes, S. and Souza, R. (1995) On the Spectral Density of a Class of Chaotic Time Series. To appear in *Journal of Time Series Analysis*.
11. Lopes, A., Lopes, S. and Souza, R. (1996) Spectral Analysis of Chaotic Transformations. To appear in *Brasilian Journal of Probability and Statistics*, 10 , No. 2.
12. Parry, W. and Pollicott, M. (1990) Zeta Functions and the Periodic Orbit Structure of Hyperbolic Dynamics. *Asterisque*, 187-188.
13. Robinson, C. (1995) *Dynamical Systems: Stability, Symbolic Dynamics and Chaos*. Boca Raton: CRC Press.
14. Ruelle, D. (1987) One-Dimensional Gibbs States and Axiom A Diffeomorphisms. *Journal of Differential Geometry*, 25 , 117-137.
15. Ruelle, D. (1978) *Thermodynamic Formalism*. Massachusetts: Addison-Wesley.
16. Rugh, H. H. (1992) The correlation spectrum for hyperbolic analytic maps. *Nonlinearity*, 5 , 1237-1263.
17. Tong, H. (1990) *Non-linear Time Series: A Dynamical System Approach*. Oxford: Clarendon.




Figure 1




Figure 2




Figure 3

Publicações do Instituto de Matemática da UFRGS
Cadernos de Matemática e Estatística

Série A: Trabalho de Pesquisa

1. Marcos Sebastiani Artecona - Transformation des Singularités - MAR/89
2. Jaime Bruck Ripoll - On a Theorem of R. Langevin about Curvature and Complex Singularities - MAR/89
3. Eduardo Cisneros, Miguel Ferrero e Maria Inés Gonzales - Prime Ideals of Skew Polynomial Rings and Skew Laurent Polynomial Rings - ABR/89
4. Oclide José Dotto - ε -Dilations - JUN/89
5. Jaime Bruck Ripoll - A Characterization of Helicoids - JUN/89
6. Mark Thompson e V. B. Moscatelli - Asymptotic Distribution of Liusternik-Schnirelman Eigenvalues for Elliptic Nonlinear Operators - JUL/89
7. Mark Thompson - The Formula of Weyl for Regions with a Self-Similar Fractal Boundary - JUL/89
8. Jaime Bruck Ripoll - A Note on Compact Surfaces with Non Zero Constant Mean Curvature - OUT/89
9. Jaime Bruck Ripoll - Compact ε -Convex Hypersurfaces - NOV/89
10. Jandyra Maria G. Fachel - Coeficientes de Correlação Tipo-Contigência - JAN/90
11. Jandyra Maria G. Fachel - The Probability of Occurrence of Heywood Cases - JAN/90

12. Jandyra Maria G. Fachel - Heywood Cases in Unrestricted Factor Analysis - JAN/90
13. Julio Cesar R. Claeysen e Tereza Tsukazan de Ruiz - Dynamical Solutions of Linear Matrix Differential Equations - JAN/90
14. Maria T. Albanese - Behaviour of the Likelihood in Latent Analysis of Binary Data - ABR/91
15. Maria T. Albanese - Measurements of the Latent Trait Analysis of Binary Data - ABR/91
16. Maria T. Albanese - Adequacy of the Asymptotic Variance-Covariance Matrix Using Bootstrap Jackknife Techniques in Latent Trait Analysis of Binary Data - ABR/91
17. Maria T. Albanese - Latent Variable Models for Binary Response - ABR/91
18. Mark Thompson - Kinematic Dynamo in Random Flows - DEZ/90
19. Jaime Bruck Ripoll e Marcos Sebastiani Artecona - The Generalized Map and Applications - AGO/91
20. Jaime Bruck Ripoll, Suzana Fornari e Katia Frensel - Hypersurfaces with Constant Mean Curvature in the Complex Hyperbolic Space - AGO/91
21. Suzana Fornari e Jaime Bruck Ripoll - Stability of Compact Hypersurfaces with Constant Mean Curvature - JAN/92
22. Marcos Sebastiani Artecona - Une Généralisation de L'Invariant de Malgrange - FEV/92
23. Cornelis Kraaikamp e Artur Lopes - The Theta Group and the Continued Fraction with Even Partial Quotients - MAR/92

24. Silvia Lopes - Amplitude Estimation in Multiple Frequency Spectrum - MAR/92
25. Silvia Lopes e Benjamin Kedem - Sinusoidal Frequency Modulated Spectrum Analysis - MAR/92
26. Silvia Lopes e Benjamin Kedem - Iteration of Mappings and Fixed Spectrum Analysis - MAR/92
27. Miguel Ferrero, Eduardo Cisneros e Maria Ines Gonzales - Ore Extensions and Jacobson Rings - MAI/92
28. Sara C. Carmona - An Asymptotic Problem for a Reaction-Diffusion Component - JUL/92
29. Luiz Fernando Carvalho da Rocha - Unique Ergodicity of Interval Exchange Maps - JUL/92
30. Sara C. Carmona - Wave Front Propagation for a Cauchy Problem With a Fast Component - OUT/92
31. Marcos Sebastiani Artecona e Iván Pan Pérez - Intersections Transverses dans l'Espace Projectif - OUT/92
32. Miguel Ferrero - Closed Bimodules over Prime Rings: Closed Submodules and Applications to Rings Extensions - DEZ/92
33. Dinara W. X. Fernandez - Método da Máxima Verossimilhança Restrita para Estimação de Componentes de Variância - SET/93
34. Martin Knott e M. Teresa Albanese - Polymiss: A Computer Program for Fitting a One- or Two-Factor Logit-Probit Latent Variable Model to Polynomous Data when Observations may be Missing - OUT/93
35. Peter Struss e Waldir L. Roque - Foundations and Applications of Qualitative Reasoning and Model-Based-Diagnosis - OUT/93

36. Edmund R. Puczylowski - On Koethe's Problem - OUT/93
37. Luis G. Mendes e Marcos A. A. Sebastiani Artecona - Sur la Densité des Systèmes de Pfaff sans Solution Algébrique - JAN/94
38. Artur O. Lopes e Luiz F. C. da Rocha - Invariant Measures for Gauss Maps Associated with Interval Exchange Maps - MAR/94
39. Artur O. Lopes e Roberto Markarian - Open Billiards: Cantor Sets, Invariant and Conditionally Invariant Probabilities - AGO/94
40. Pierre Collet, Antonio Galves e Artur O. Lopes - Maximum Likelihood and Minimum Entropy Identification of Grammars - AGO/94
41. Ondina F. Leal e Jandyra M. G. Fachel - Antropologia do Corpo e Pesquisa sobre Sexualidade: Dados Qualitativos e Tratamento Estatístico, uma Proposta Metodológica - DEZ/94
42. Artur O. Lopes e Sílvia R. C. Lopes - Parametric Estimation and Spectral Analysis of Chaotic Time Series - NOV/95
43. Artur O. Lopes, Rafael R. Souza e Sílvia R. C. Lopes - On the Spectral Density of a Class of Chaotic Time Series - DEZ/95
44. Marcelo Bergmann e Sílvia R. C. Lopes - Instantaneous Frequency Detection by the Contraction Mapping Method - FEV/96
45. Artur O. Lopes, Rafael R. Souza e Sílvia R. C. Lopes - Spectral Analysis of Chaotic Transformations - MAR/96
46. Artur O. Lopes, Rafael R. Souza e Sílvia R. C. Lopes - Spectral Analysis of Expanding One-dimensional Chaotic Transformation - ABR/96

UNIVERSIDADE FEDERAL DO RIO GRANDE DO SUL
INSTITUTO DE MATEMÁTICA
NÚCLEO DE ATIVIDADES EXTRACURRICULARES

Os Cadernos de Matemática e Estatística publicam as seguintes séries:

Série A: Trabalho de Pesquisa

Série B: Trabalho de Apoio Didático

Série C: Colóquio de Matemática SBM/UFRGS

Série D: Trabalho de Graduação

Série F: Trabalho de Divulgação

Série G: Textos para Discussão

Toda correspondência com solicitação de números publicados e demais informações deverá ser enviada para:

NAEC - NÚCLEO DE ATIVIDADES EXTRACURRICULARES
INSTITUTO DE MATEMÁTICA - UFRGS
AV. BENTO GONÇALVES, 9500 - PRÉDIO 43111
CEP 91509 - 900 AGRONOMIA - POA/RS
FONE: 316 6197
FAX: 336 15 12

