




Revista Brasileira de Psiquiatria

RBPPsychiatry

Official Journal of the Brazilian Psychiatric Association
Volume 34 • Number 3 • October/2012



ORIGINAL ARTICLE

Psychiatric disorders among individuals who drive after the recent use of alcohol and drugs

Sibele Faller,¹ J. Matthew Webster,² Carl G. Leukefeld,² Daniela Benzano Bumaguin,¹ Paulina do Carmo Arruda Vieira Duarte,³ Raquel De Boni,¹ Flavio Pechansky¹

¹ Center for Drug and Alcohol Research, Universidade Federal do Rio Grande do Sul, Brazil

² Center on Drug and Alcohol Research, University of Kentucky (UK), Lexington, KY, USA

³ Secretaria Nacional de Políticas sobre Drogas (Brazilian Secretariat for Drug Policies), Brazil

Received on October 7, 2011; accepted on February 29, 2012

DESCRIPTORS

Drivers;
Alcohol;
Drugs;
Psychiatric Disorders;
Telephone.

Abstract

Introduction: This cross-sectional study assessed the prevalence of psychiatric disorders among drivers, as well as the association between recent alcohol and drug use and psychiatric diagnoses using telephone interviews. **Methods:** Drivers (n = 1,134) included in a roadside survey from 25 Brazilian state capitals were given a breathalyzer test, and their saliva was tested for psychoactive drugs. A telephone interview was conducted to perform psychiatric disorder evaluations using the MINI. This association was analyzed with a Poisson regression model. **Results:** The prevalence of any psychiatric disorder was 40.5% among drivers with recent alcohol or drug use, compared with 12.9% among the other drivers. Alcohol/drug-positive drivers reported a higher prevalence of depression (19.4%), mania (6.5%), hypomania (5.4%), post-traumatic stress disorder (8.6%), antisocial personality (7.8%), and substance/alcohol abuse or dependence (48.1%) compared with other drivers (3.5, 2.5, 2.1, 0.5, 1.3 and 18.3% [p < 0.001], respectively). Drivers with recent alcohol or drug use were 2.5 times more likely to have a psychiatric diagnosis (CI: 1.8-3.6, p < 0.001). **Discussion:** This is the first study in a low-/middle-income country to evaluate psychiatric disorders in drivers with recent alcohol or drug using telephone interviews. Psychiatric disorders were found to be associated with drug and alcohol use. This type of epidemiological information for curtailing related driving problems, as these psychiatric conditions are diagnosable. **Final Remarks:** The results of this study can aid in the design of interventions, treatment programs and focused psychiatric evaluations, both in Brazil and abroad.

DESCRITORES:

Motoristas;
Álcool;
Drogas;
Transtornos
psiquiátricos;
Telefone.

Transtornos psiquiátricos entre indivíduos que dirigem sob uso recente de álcool e drogas**Resumo**

Introdução: Este estudo transversal avaliou a prevalência de transtornos psiquiátricos e a associação entre dirigir alcoolizado e transtornos psiquiátricos entre motoristas por meio de entrevistas telefônicas. **Método:** Motoristas (n = 1.134) selecionados a partir de um estudo realizado nas rodovias de 25 capitais brasileiras foram etilometrizados e tiveram sua saliva testada para substâncias psicoativas. Uma entrevista por telefone foi realizada para obter dados sobre transtornos psiquiátricos através do MINI. A associação entre transtornos psiquiátricos e dirigir com uso recente de substâncias foi analisada com um modelo de regressão de Poisson. **Resultados:** A prevalência de transtornos psiquiátricos foi de 40,5% entre os motoristas que beberam/usaram drogas, em comparação com 12,9% nos demais motoristas. Motoristas que dirigiram com uso recente de substâncias relataram uma maior prevalência de depressão (19,4%), mania (6,5%), hipomania (5,4%), transtorno de estresse pós-traumático (8,6%), personalidade antissocial (7,8%), e abuso ou dependência de álcool/drogas (48,1%) quando comparados com os outros motoristas (3,5%, 2,5%, 2,1%, 0,5%, 1,3% e 18,3%, p < 0,001), respectivamente). Motoristas com uso recente de substâncias apresentaram 2,5 mais chance de ter qualquer diagnóstico psiquiátrico (IC: 1,8-3,6, p < 0,001). **Discussão:** Este é o primeiro estudo em países de baixa e média renda cujo objetivo é avaliar transtornos psiquiátricos em motoristas com uso recente de substâncias por meio de entrevistas telefônicas. Transtornos psiquiátricos foram associados com o comportamento de dirigir com uso recente de substâncias. Este tipo de informação epidemiológica é fundamental para estratégias de planejamento que visam enfrentar problemas relacionados com o beber e dirigir, uma vez que estas condições psiquiátricas são diagnosticáveis. **Observações Finais:** Os resultados deste estudo podem auxiliar no desenvolvimento de intervenções, programas de tratamento e avaliação psiquiátrica, tanto no Brasil quanto no exterior.

Introduction

According to the World Health Organization, low-and middle-income countries (LAMICs) have higher road traffic fatality rates (21.5 and 19.5 per 100,000 people, respectively) than high-income countries 10.3/100,000 people, making vehicle crashes a serious concern for public health.¹ Although 90% of traffic deaths occur in developing countries, data on risk factors are limited, making it difficult to conduct systematic studies on the characteristics of the drivers who cause these fatal crashes. Because the use of alcohol is associated with 41% of all traffic accidents, understanding the degree to which individuals drive under the influence (DUI) of alcohol or other drugs, as well as whether they have comorbid psychiatric problems,^{2,3} would provide valuable information for policy planning and prevention efforts.

The Brazilian government is struggling to develop adequate policies against DUI based on reliable data. There have been few studies suggesting that the prevalence of drinking and driving among Brazilian drivers is markedly high. Pechansky et al.,⁴ based on an analysis of data from the first National Survey on Household Consumption Patterns of Alcohol⁴ found that approximately 34% of drivers reported drinking and driving in the past year. This percentage is more than twice (13.2%) the percentage of adults who reported alcohol-impaired driving in the U.S.A,⁵ and 85 (0.4%) times higher than rate in Norway.⁶ Brazil has limited information on how many individuals drive after drinking alcohol or using drugs, and there are no published findings related to psychiatric disorders and their relationships to traffic and driving in the country, which may be due to geographical and financial difficulties. The lack of data on this issue is an important void, as driving under the influence of alcohol

or drugs has been associated with a high prevalence of psychiatric conditions in the U.S., and it is reported in several studies to be a factor associated with an increased risk of crash.^{3,7-14} Therefore, “What is the rate of psychiatric disorders among impaired drivers?” is an important research question, particularly in Brazil because the country is passing through a period of interest in the relationship between alcohol and traffic accidents. The government has shown its intention to reorganize public policies to this need, such as by the recent the implementation of the new “dry law” and increased investment in research in the area.

In 2001, an important study of 1,105 DUI offenders⁷ revealed that, among those drivers with a lifetime alcohol-use disorder, 50% of women and 33% of men were diagnosed with at least one psychiatric condition, often major depressive disorder and/or PTSD, as well as drug abuse or dependence. In contrast to the prevalence found from the National Comorbidity Survey (NCS) impaired drivers exhibit a higher-than-expected prevalence of psychiatric disorders. For example, 61% (women) and 70% (men) of respondents demonstrated lifetime alcohol dependence, which is more than twice the respective NCS sample prevalence. The possibility that vehicle crashes and drug use are influenced by psychiatric disorders is supported by the literature, but most studies are conducted with samples of DUI offenders admitted to court-mandated treatment programs, and comparisons with road surveys are often made with different samples of drivers.^{8,9,12,13,15-18} Therefore, the determination of the association between traffic accidents and psychiatric diagnoses or identification of risk factors for these morbidities can determine interventions and post-treatment planning for driving offenders to reduce relapse, consumption and traffic violations.¹⁷

The purpose of this study was to assess the prevalence of psychiatric disorders among drivers who drove under the influence of alcohol and other substances and the association between impaired driving and psychiatric problems among drivers in Brazil utilizing telephone interviews with a contemporary bioethics approach using rolling consent.¹⁹ Specifically, drivers who tested positive for substances were compared to those who did not with regard to mood disorders, PTSD and antisocial personality disorder (ASPD) - selected according to the prevalence in studies with similar samples. Consistent with the international literature, our hypothesis was that psychiatric problems would be more prevalent among drivers who tested positive for substances via a breathalyzer or saliva test.

Methods

Participants

Volunteers who participated in this study were recruited during data collection for a larger study about the prevalence of positive breath for alcohol and other psychoactive substances in professional and private drivers in Brazil.^{20,21} This was a cross-sectional study with a sample selected in blocks in each capital, proportional to the number of drivers in the specific state and stratified by type of vehicle - automobiles, motorcycles, buses and trucks with random selection in proportion to the fleet size of each state. Private and professional drivers who were driving on federal highways which intersect the metropolitan areas of 23 Brazilian state capitals were evaluated.

The sample size had to be calculated considering the limited budget available from the funding agency, and the prevalence of the use of amphetamines (6%, with a 2% margin of error, and a confidence level of 95%) that we supposed to be smaller than the use of alcohol. For that, 542 drivers should be included in each stratum, which would allow for an estimate of alcohol use of 30%, with a margin of error smaller than 1.5%. A minimum number of drivers per state (blocks) was determined. The states with estimated $n < 100$ or those whose n was very close to 100 were consolidated, with a total of 21 states ($n = 100 \times 21 = 2,100$ cases). According to the previous calculation, a total of 1,288 (3,388-2,100) subjects would be divided among the other states with an estimated $n > 100$. The total of drivers of the states with $n > 100$ when added up resulted in 38,322,119 drivers, and the proportions for each state were recalculated based on this figure. Finally, the number of drivers to be approached in each state was estimated for this total of subjects that were a surplus in each state, maintaining an intermediate tercile ($n = 138$), and one single state was responsible for the last third of the sample (São Paulo, $n = 550$). The vehicles were selected proportionally to the number of vehicles in each stratum, according to random number lists generated prior to the data collection.

Data were collected on Fridays and Saturdays from 12 p.m. to 12 a.m. due to economic reasons and logistical details, including security issues. The main objective of the larger study was to estimate risk factors for driving with recent substances use and the prevalence of alcohol and other drugs among Brazilian drivers. During the roadside survey, the presence of alcohol was verified by a breathalyzer Alco-Sensor IV (Intoximeters, Inc), and the presence of

drugs - amphetamine, cocaine, marijuana (tetrahydrocannabinol - THC) and benzodiazepines - was ascertained via a saliva test (Quantisal, Inc). These procedures are less intrusive than other methods and consistent with recommendations found in the literature.^{22,23} For the screening method, the ELISA assay was applied. Confirmatory analyses in oral fluid were performed by Liquid Chromatography Tandem Mass Spectrometry (LC-MS). At the end of the roadside survey interview, each participant was asked to provide his/her name and a telephone number for further contact. Attempts were made to contact all those who accepted to be called. Out of the 2,957 drivers interviewed in the main study, 2,580 (87.25%) agreed to have further contact. After a maximum of 5 attempts, 1,134 individuals had completed the telephone interview and 199 (7.71%) refused to participate at the time of telephone contact. Of the individuals who were not interviewed and did not formally refuse to participate, 300 (11.62%) were scheduled for further contact but did not respond to subsequent calls, and 947 (36.7%) were not found. This resulted in a response rate of 43.95%. Approximately 8% ($n = 94$) of the sample tested positive for alcohol or drugs.

Measures

Demographics - gender, age, type of vehicle, employment and educational level - were obtained through a series of questions developed specifically for the study and were based on questionnaires of the Brazilian Institute of Geography and Statistics (*Instituto Brasileiro de Geografia e Estatísticas* - IBGE).

Psychiatric disorders - current depressive episode, hypomanic/manic episode (grouped for Poisson Regression analysis), PTSD, substance abuse/dependence and ASPD - were measured by the Mini International Neuropsychiatric Interview (MINI), instrument adapted for the Brazilian context.²⁴ The primary author was trained on the use of the Brazilian version of the MINI by one of the key investigators in the country.

Procedures

Telephone interviews were conducted by eight trained interviewers at a call center (through the structure of "VivaVoz", a hotline for information and referrals about drug abuse). Interviewers were instructed to introduce themselves as investigators from the Universidade Federal do Rio Grande do Sul in order to avoid adverse influence in the decision to participate. After being informed of the nature of the survey, drivers who consented had their assessment criteria checked. Two postgraduate level professionals were in charge of real time supervision during interviews. During data collection, monitoring of telephone interviews was conducted by the study coordinator (SF) - with consent from volunteers and through meetings to review logistical problems and assess quality control. Subjects were located and contacted with the minimum information needed to conduct the phone call - only the first name and telephone number were placed on the worksheet that guided the interviewers. Additional information, such as last name and data collected during the roadside study were not related to the contact information and then excluded from the database, in order to preserve participant personal information.

Ethical Issues

Informed consent was obtained in stages with a rolling consent approach. This iterative process prevents the data from being completely discarded if the participant decides to leave the study and does not keep the individual feeling obligated to answer questions until the end of the interview, minimizing the chance of encountering embarrassment. The respondent was asked to consent at every change of section (1- initial consent; 2- between demographics and psychiatric section; and 3- between psychiatric and drug abuse section), so he/she could choose to continue or terminate the interview, therefore validating the information obtained before termination. The study was approved by the Institutional Review Board of Hospital das Clínicas of Porto Alegre.

Statistical analyses

Categorical variables were compared using a series of chi-square tests. In order to identify the association between psychiatric disorders and driving under the influence of alcohol or drugs (dependent variable), a Poisson regression model with robust variance estimator was conducted, whereby prevalence ratios (PR) and confidence intervals (CI) of 95% were estimated. The analysis was adjusted for *gender*, *age*, *income*, *educational level*, and *any substance abuse or dependence* - selected according to information found in the literature.²⁵⁻²⁹ Due to the finding of co-linearity between

income and educational level, this last variable was removed from the model. Data were analyzed using SPSS v.18. (SPSS Inc., Chicago, IL, USA).

Results

The sample was comprised primarily of men (95%) with a median of 36 years of age (P25: 28.0; P75:44.5). Participants and non-participants were compared regarding demographics using data collected via the main study. No statistical differences in demographic information were found between drivers who were positive versus those who were negative regarding alcohol/drug use. The total sample was analyzed according to the type of driver participation to determine whether those denying permission differed from those providing permission in their "use of substances on the road status". Comparing drivers who participated in the main study and did not agree to have further contact (6.9%); who answered the interview (8.3%); were not found (9.8%); and refused to participate (7.0%), the results demonstrated no statistically significant difference ($p = 0.20$). The comparisons between drivers with recent use of alcohol and other drugs and the other participants are shown in Table 1.

Drivers who tested positive for alcohol or drug use had a higher prevalence of psychiatric diagnoses when compared to the other drivers without alcohol and other drugs using bivariate analysis. The prevalence of any type of psychiatric

Table 1 Compared prevalence of sociodemographic characteristics between drivers who tested positive for substance use and other drivers in Brazil

Sociodemographic characteristics	Drivers positive for substance		Other drivers		p
	n = 94	(8.3%)	n = 1,040	(91.7%)	
Total					
Gender	94	-	1,036	-	
Male	91	(96.8)	98.4	(95.0)	0.616
Female	3	(3.2)	52	(5.0)	
Age (years)	94	-	1,034	-	
18 - 29	23	(24.5)	334	(32.3)	0.306
30 - 44	45	(47.9)	442	(42.7)	
45 - 64	23	(24.5)	242	(23.4)	
65 and older	3	(3.2)	16	(1.5)	
Employ	91	-	1,026	-	
Employed	89	(97.8)	984	(95.9)	0.573
Unemployed	2	(2.2)	42	(4.1)	
Educational level	92	-	1,027	-	
Elementary and middle school	27	(29.3)	318	(31.0)	0.925
High school	41	(44.6)	437	(42.6)	
Bachelor's degree or more	24	(26.1)	272	(26.5)	
Income	92	-	1,022	-	
R\$ 0 - 936	52	(56.5)	577	(56.5)	0.686
R\$ 937 - 1,561	17	(18.5)	221	(21.6)	
R\$ 1,562 or more	23	(25.0)	224	(21.9)	

* Non-excluding variable.

Table 2 Association of selected psychiatric disorders and use of substances among Brazilian drivers analyzed by Poisson regression

Psychiatric disorder	Crude PR	(95% CI) [§]	Adjusted PR	(95% CI) [§]	p
Mood Disorders [*]	3.0	(2.1 - 4.3)	2.5	(1.7 - 3.7)	< 0.001
PTSD [†]	15.9	(5.2 - 48.9)	4.5	(1.5 - 13.3)	0.006
ASPD [‡]	6.0	(2.5 - 4.7)	3.1	(1.0 - 9.5)	0.039
Any Disorder	3.1	(2.3 - 4.3)	2.5	(1.8 - 3.6)	< 0.001

^{*} Depression, mania and hypomania.

[†] Posttraumatic stress disorder, not adjusted for gender due to the lack of women presenting this disorder in the study.

[‡] Antisocial personality disorder, not adjusted for gender due to the lack of women presenting this disorder in the study.

[§] Confidence Interval.

disorder was 40.5% among drivers who tested positive and 12.9% among the other drivers ($p < 0.001$). Drivers who were tested positive reported a significantly higher prevalence of psychiatric diagnoses of depression (19.4%), mania (6.5%), hypomania (5.4%), post-traumatic stress disorder (8.6%), antisocial personality (7.8%), and substance or alcohol abuse or dependence (48.1%) when compared to other drivers (3.5%, 2.5%, 2.1%, 0.5%, 1.3% and 18.3%, respectively). The results from the Poisson regression analysis are shown in Table 2.

According to the results of the Poisson regression analysis, there was a statistically significant association between psychiatric disorders and driving under the influence of psychoactive substances. Even after variable adjustment, participants who tested positive for substances were 2.5 times more likely to have psychiatric symptoms.

Discussion

Psychiatric disorders and drivers with recent use of alcohol and other drugs

Results show that psychiatric disorders were clearly associated with the practice of drunk/drugged driving in this sample of Brazilian drivers. Participants who tested positive for substances were more likely to have mood disorder, PTSD, ASPD or any disorder. The prevalence of psychiatric problems among Brazilian drivers testing positive for alcohol or drugs was high compared to Australia (15.5%), Germany (31.1%), Netherlands (23.2%) and U.S. (26.2%) general population,³⁰ although the sampling time chosen - a transition law period in Brazil - makes it difficult to compare the present results with the prevalence of alcohol and drug findings in other countries. It is documented in the literature that drivers under the influence of alcohol and drugs present a greater prevalence of psychiatric disorders including alcohol and other substance abuse.^{7-9,18} In this study, like in the other studies published in international literature, mood disorders, PTSD, substance abuse and dependence were very prevalent psychiatric disorders in the sample of drivers who test positive for substances, indicating the need for greater attention to these specific disorders.^{9,31} What seems to be significant in relation to these results is the important difference between the two types of drivers, even when compared with the findings of other studies previously conducted in the U.S.⁷ If drivers could be properly assessed and receive treatment for potential psychiatric problems, the prevalence of driving under the influence of alcohol and

drugs could eventually decrease. This, however, poses a tricky solution, since a screening of these symptoms and conditions should be made at the preliminary stages of traffic education and licensing for young drivers. Nevertheless, psychiatric testing to get a license or to have a suspended license reinstated seems difficult to administer. Since most of these conditions can be properly ascertained by a focused set of questions, this would imply much further training from examiners than what is typically provided in many LAMIC countries. Consequently, it seems that the mandatory substance use treatment for all DUI offenders would be more feasible. Some important changes in the environment, such as restriction on times for alcohol sales and banishment of alcohol in certain places like gas stations and bars on the roads, may be useful to address drivers' behavior regarding alcohol consumption. These policies must be aligned with adequate enforcement, which is currently limited in Brazil - a situation that can be overcome with additional resources and public support. The secondary focus of attention would be the referral for medical and eventually psychiatric treatment of those cases properly identified after a car crash or traffic violation before allowing them to return to driving.

Although data collection began soon after a new law passed in 2008 prohibiting the consumption of any level of alcohol or other substance by drivers in the country,³² the prevalence of drivers who tested positive for a substance (8%) seems to be high when compared to surveys conducted in Australia (3.7% positive for at least one illicit substance) and Norway (4.5% positive for alcohol or drugs) and similar when compared to Canada (8% for alcohol).^{6,33,34} According to the new Brazilian law, drivers who drink may have to pay a fine, have their license suspended for 12 months or even be arrested, an important penalty, depending on the concentration of alcohol per liter of blood. So, considering the Brazilian socio-political context and surveys performed in other countries, we can say that the prevalence of drivers who drove under the influence of any substance is relatively high and certainly helps maintain the frequency of vehicle crashes in Brazilian roads.

Implications

Prevention of problems on the roads such as drinking and driving is a large policy issue. For a policy to be effective, however, it is necessary to learn the reality and variables involved in order to provide evidence to base interventions and design programs that address the need of the driver and society in the historical and cultural context where they need to be implemented.

To the best of our knowledge, this is the first study of national scope in Brazil and in LAMIC that aims at evaluating psychiatric disorders of substance positive drivers who drove with recent use of alcohol and drugs measured by breathalyzer and saliva test, comparing this result with other drivers from the same sample. Also, we believe it was possible to set a landmark on the use of a contemporary bioethics approach through rolling or scaled consent applied through telephone surveys in Brazil. Another great strength of this study was the identification of the drivers with recent use of alcohol and drugs using objective tests rather than self-report measures of the use of psychoactive substances. Having the objective measures strongly attests to the participants' driving behavior.

In this sense, we believe some aspects were essential to ensure the study's quality. Among them the fact that data were collected in a structured call center, allowing better organization of calls and staff; real time monitoring of data collection, that in turn allowed quick problem solution during interviews, being a determinant factor to increase the quality of the information obtained. Interviewers were acquainted both with the technicalities of the call center as well as with diagnostics procedures.

Limitations

Our findings must be viewed in the context of the study's limitations. First, the sample may not have national representativeness due to the restriction of the days and time of data collection. This was due to limitations and issues involving security, as data collection on highways after midnight was discouraged by police authorities. Second, the time allotted for calls to the drivers who consented to participate was set at a maximum of two weeks after consent, so that all regions could be covered in the same way. Third, MINI is an instrument with a considerable sensitivity (at least 0.77 for the modules used - 0.94 for Major Depressive Episode), a fact that may have overestimated some results (24). Fourth, we have obtained a response rate lower than those presented in face-to-face studies. As shown in an editorial piece published by *Addiction*, 30% is the average response rate in surveys conducted about alcohol in the U.S. general population, which, in this case, may contribute to underestimation of the prevalence of impaired driving and psychiatric problems. However, response rates for other telephone surveys, ranging from 40 to 80%, were not different from the ones produced in this study.³⁵⁻³⁹ On the other hand, the results show homogeneity among groups in the comparison between respondents and non-respondents regarding demographics and substances recent use measured on the roads. This suggests that this similarity is likely to remain in other aspects, which makes drivers comparable despite the differences in sample size for each group.

Further investigation of this matter would perhaps include other psychiatric diagnoses not ascertained in this study, such as generalized anxiety disorder, obsessive-compulsive disorder, attention deficit hyperactivity disorder, among others, since many have already shown they can be associated with drunk/drugged driving. Future studies in Brazil could compare drivers with multiple charges of DUI and those with no record of recurrence, considering specific penalties and

restrictions for each group. Since a recent study has shown that alcohol alone accounted for 47% of the total costs of traffic crashes in the city of Porto Alegre, Brazil, for the year 2009,⁴⁰ another investigation of importance could be the estimation of the impact that evaluations and treatment of psychiatric disorders - and consequent reduction of crashes - would have on the economy of the country, aiming to evaluate the cost-benefit analysis of the investment on public policies directed for the issue. Although costs would be higher at first impression, it is believed that not only could such problems be avoided or at least minimized, but also costs in the end would diminish.

Final Remarks

The type of epidemiological information presented in this paper highlights the fact that driving after recent use of alcohol and other drugs is an important public health problem throughout the world, including in LAMIC, being vital to the planning of broader strategies aimed at reaching such psychiatric disorders and problems related to alcohol and other drugs. This information can be used not only for traffic education purposes, such as advertising campaigns and training courses for suspended drivers but also to design interventions, treatment programs and psychiatric evaluations, both in Brazil and abroad.

Acknowledgements

The authors would like to acknowledge the staff of VivaVoz, mainly Helena Maria Tannhauser Barros, Maristela Ferigolo, Taís de Campos Moreira, Bárbara Diniz, Cássio Machado, Solange Paim, Glaucus Maidana, Ricardo Schneider and Guilherme Foletto.

Disclosures

Sibele Faller

Employment: Center for Drug and Alcohol Research of the Hospital das Clínicas de Porto Alegre, Universidade Federal do Rio Grande do Sul, Brazil.

J. Matthew Webster

Employment: Center on Drug and Alcohol Research, University of Kentucky (UK), Lexington, KY, USA.

Carl G. Leukefeld

Employment: Center on Drug and Alcohol Research, University of Kentucky (UK), Lexington, KY, USA.

Daniela Benzano Bumaguin

Employment: Center for Drug and Alcohol Research of the Hospital das Clínicas de Porto Alegre, Universidade Federal do Rio Grande do Sul, Brazil.

Paulina do Carmo Arruda Vieira Duarte

Employment: Secretaria Nacional de Políticas sobre Drogas (Brazilian Secretariat for Drug Policies), Brazil.

Raquel De Boni

Employment: Center for Drug and Alcohol Research of the Hospital das Clínicas de Porto Alegre, Universidade Federal do Rio Grande do Sul, Brazil.

Flavio Pechansky

Employment: Center for Drug and Alcohol Research of the Hospital das Clínicas de Porto Alegre, Universidade Federal do Rio Grande do Sul, Brazil.

Funding for this study was provided by the National Secretariat for Drug Policies (SENAD) under grant #07-069.

* Modest

** Significant

*** Significant. Amounts given to the author's institution or to a colleague for research in which the author has participation, not directly to the author.

References

1. WHO. Global Status Report on Road Safety. Geneva, 2009.
2. Traffic NH, Administration S. Report No.: DOT HS 810 809. In: Administration NHTS, editor. Washington, DC2006.
3. Freeman J, Maxwell JC, Davey J. Unraveling the complexity of driving while intoxicated: a study into the prevalence of psychiatric and substance abuse comorbidity. *Accid Anal Prev.* 2011;43(1):34-9.
4. Pechansky F, De Boni R, Diemen L, Bumaguin D, Pinsky I, Zaleski M, et al. Highly reported prevalence of drinking and driving in Brazil: data from the first representative household study. *Rev Bras Psiquiatr.* 2009;31(2):125-30.
5. SAMHSA SAaMHSACfBHSAQ-. National Survey on Drug Use and Health Report: State Estimates of Drunk and Drugged Driving. Rockville: 2010.
6. Gjerde H, Normann PT, Pettersen BS, Assum T, Aldrin M, Johansen U, Lena Kristoffersen L, Øiestad EL, Christophersen AS, Mørland J. Prevalence of alcohol and drugs among Norwegian motor vehicle drivers: a roadside survey. *Accid Anal Prev.* 2008;40(5):1765-72.
7. Lapham S, Smith E, C'de Baca J, Chang I, Skipper B, Baum G, Hunt WC. Prevalence of psychiatric disorders among persons convicted of driving while impaired. *Arch Gen Psychiatry.* 2001;58(10):943-9.
8. Lapham S, C'de Baca J, McMillan G, Lapidus J. Psychiatric disorders in a sample of repeat impaired-driving offenders. *J Stud Alcohol.* 2006;67(5):707-13.
9. McMillan G, Timken D, Lapidus J, C'de Baca J, Lapham S, McNeal M. Underdiagnosis of comorbid mental illness in repeat DUI offenders mandated to treatment. *J Subst Abuse Treat.* 2008;34(3):320-5.
10. Sutton L. Assessment of alcohol dependence and other psychiatric disorders: implications for rehabilitation programs of the DUI offender. *Blutalkohol.* 1994;31(1):33-9.
11. McMillan D, Pang M, Wells-Parker E, Anderson B. Behavior and personality traits among DUI arrestees, nonarrested impaired drivers, and nonimpaired drivers. *Int J Addict.* 1991;26(2):227-35.
12. Albanese MJ, Nelson SE, Peller AJ, Shaffer HJ. Bipolar disorder as a risk factor for repeat DUI behavior. *J Affect Disord.* 2010;121(3):253-7.
13. Peller AJ, Najavits LM, Nelson SE, LaBrie RA, Shaffer HJ. PTSD among a treatment sample of repeat DUI offenders. *J Trauma Stress.* 2010;23(4):468-73.
14. Vaughn MG, Define RS, Delisi M, Perron BE, Beaver KM, Fu Q, Howard MO. Sociodemographic, behavioral, and substance use correlates of reckless driving in the United States: Findings from a national sample. *J Psychiatr Res.* 2010.
15. C'de Baca J, Lapham S, Skipper B, Hunt W. Psychiatric disorders of convicted DWI offenders: a comparison among Hispanics, American Indians and non-Hispanic whites. *J Stud Alcohol.* 2004;65(4):419-27.
16. LaPlante DA, Nelson SE, Odegaard SS, LaBrie RA, Shaffer HJ. Substance and psychiatric disorders among men and women repeat driving under the influence offenders who accept a treatment-sentencing option. *J Stud Alcohol Drugs.* 2008;69(2):209-17.
17. Palmer R, Ball S, Rounsaville B, O'Malley S. Concurrent and predictive validity of drug use and psychiatric diagnosis among first-time DWI offenders. *Alcohol Clin Exp Res.* 2007;31(4):619-24.
18. Shaffer HJ, Nelson SE, LaPlante DA, LaBrie RA, Albanese M, Caro G. The epidemiology of psychiatric disorders among repeat DUI offenders accepting a treatment-sentencing option. *J Consult Clin Psychol.* 2007;75(5):795-804.
19. Dutton R. Food and Drug Administration public hearing of the conduct of emergency clinical research: testimony of Dr. Dutton. *Acad Emerg Med.* 2007;14(4):e33-6.
20. Pechansky F, De Boni R, Duarte P, Cubas FP, Benzano D, Von Diemen L, Leukefeld C. Consumo de drogas entre motoristas privados e profissionais do Brasil. In: Pechansky F, De Boni R, Duarte P (org). *Uso de bebidas alcoólicas e outras drogas nas rodovias brasileiras e outros estudos.* SENAD: Brasília-DF; 2010.
21. Pechansky F, De Boni R, Limberger R, Leukefeld CG, Bozzetti M, Hilgert J, et al. Drunk and drugged driving in Brazilian highways. International Council on Alcohol, Drugs and Traffic Safety (ICADTS): Oslo-Norway; 2010.
22. Pil K, Verstraete A. Current developments in drug testing in oral fluid. *Ther Drug Monit.* 2008;30(2):196-202.
23. Walsh J, Verstraete A, Huestis M, Mørland J. Guidelines for research on drugged driving. *Addiction.* 2008;103(8):1258-68.
24. Amorim P. Mini International Neuropsychiatric Interview (MINI): validation of a short structured diagnostic psychiatric interview. *Rev Bras Psiquiatr.* 2000. p. 106-15.
25. Hingson R, Winter M. Epidemiology and consequences of drinking and driving. *Alcohol Res Health.* 2003;27(1):63-78.
26. Kypri K, Stephenson S. Drink-driving and perceptions of legally permissible alcohol use. *Traffic Inj Prev.* 2005;6(3):219-24.
27. Labouvie E, Pinsky I. Substance use and driving: the coexistence of risky and safe behaviors. *Addiction.* 2001;96(3):473-84.
28. NHTSA. NHTSAs-. National survey on drinking and driving behavior. Washington, DC1998.
29. Stevenson M, Palamara P, Rooke M, Richardson K, Baker M, Baumwol J. Drink and drug driving: what's the skipper up to? *Aust N Z J Public Health.* 2001;25(6):511-3.
30. Baumeister H, Härter M. Prevalence of mental disorders based on general population surveys. *Soc Psychiatry Psychiatr Epidemiol.* 2007;42(7):537-46.
31. Oslin D, O'Brien C, Katz I. The disabling nature of comorbid depression among older DUI recipients. *Am J Addict.* 1999;8(2):128-35.
32. Andreuccetti G, Carvalho HB, Cherpitel CJ, Kahn T, Ponce JeC, Leyton V. The right choice on the right moment. *Addiction.* 2010;105(8):1498-9.
33. Davey J, Freeman J. Screening for drugs in oral fluid: drug driving and illicit drug use in a sample of Queensland motorists. *Traffic Inj Prev.* 2009;10(3):231-6.
34. Beirness DJ, Beasley EE. A roadside survey of alcohol and drug use among drivers in British Columbia. *Traffic Inj Prev.* 2010;11(3):215-21.
35. Iredell H, Shaw T, Howat P, James R, Granich J. Introductory postcards: do they increase response rate in a telephone survey of older persons? *Health Educ Res.* 2004;19(2):159-64.
36. Feveile H, Olsen O, Høgh A. A randomized trial of mailed questionnaires versus telephone interviews: response patterns in a survey. *BMC Med Res Methodol.* 2007;7:27.
37. Hocking J, Lim M, Read T, Hellard M. Postal surveys of physicians gave superior response rates over telephone interviews in a randomized trial. *J Clin Epidemiol.* 2006;59(5):521-4.
38. Patten S, Adair C, Williams J, Brant R, Wang J, Casebeer A, Beauséjour P. Assessment of mental health and illness by telephone survey: experience with an Alberta mental health survey. *Chronic Dis Can.* 2006;27(3):99-109.
39. Caetano R. Non-response in alcohol and drug surveys: a research topic in need of further attention. *Addiction.* 2001;96(11):1541-5.
40. Porto Junior SS, Sousa TRV, Correa E, Stampe MZ. Impacto econômico dos acidentes de trânsito relacionados ao uso de substâncias psicoativas. In: Pechansky F, Duarte PCAV, De Boni RB. (org). *Uso de Bebidas Alcoólicas e outras Drogas nas Rodovias Brasileiras e outros Estudos.* Porto Alegre: Digitalcom Comunicação & Design Ltda, 2010, p. 16-19.