

Diffractive photoproduction of radially excited $\psi(2S)$ mesons in photon-Pomeron reactions in PbPb collisions at energies available at the CERN Large Hadron Collider

M. B. Gay Ducati, M. T. Griep, and M. V. T. Machado

High Energy Physics Phenomenology Group, GFPAE IF-UFRGS Caixa Postal 15051, CEP 91501-970, Porto Alegre, RS, Brazil

(Received 3 April 2013; revised manuscript received 25 June 2013; published 29 July 2013)

In this work we investigate the photoproduction of radially excited vector mesons off nuclei in heavy ion relativistic collisions. In particular, we analyze the exclusive photoproduction of $\psi(2S)$ off nuclei, evaluating the coherent and the incoherent contributions to that process. The theoretical framework used in the present analysis is the light-cone dipole formalism and predictions are done for PbPb collisions at the CERN Large Hadron Collider energy of 2.76 TeV. The theoretical uncertainties are analyzed and comparison is also done to the recent ALICE Collaboration data for the $\psi(1S)$ state photoproduction.

DOI: [10.1103/PhysRevC.88.014910](https://doi.org/10.1103/PhysRevC.88.014910)

PACS number(s): 12.38.Bx, 13.60.Hb, 24.85.+p, 25.75.Dw

I. INTRODUCTION

An outstanding feature of diffractive photoproduction of mesons at the high energy regime is the possibility to investigate the Pomeron exchange. In such an energy domain hadrons and photons can be considered as color dipoles in the mixed light cone representation [1]. In particular, their transverse sizes are to be considered frozen during the interaction reaction. Then, the scattering process is characterized by the color dipole cross section representing the interaction of those color dipoles with the target (protons or nuclei). The color evolution of the dipole cross section at small Bjorken- x is given by the solution of a nonlinear evolution equations. It has been known for a long time that dipole sizes of magnitude $r \sim 1/\sqrt{m_V^2 + Q^2}$ (m_V is the vector meson mass) are probed by the $1S$ vector meson production amplitude [1]. As far as heavy quarks are concerned, the sufficiently large mass of quarkonium states makes the amplitude to be perturbatively calculable even at photoproduction region $Q^2 \rightarrow 0$. The diffractive production of the $2S$ radially excited vector mesons, like $\psi(2S)$ and $\Upsilon(2S)$, is especially interesting due to the node effect [2]. It means a strong cancellation of dipole size contributions to the production amplitude from the region above and below the node position in the $2S$ radial wave function [3]. This is the origin of the large suppression of the photoproduction of radially excited vector mesons $2S$ versus $1S$. It is an experimental fact that the ratio $\sigma(\psi')/\sigma(\psi) \simeq 0.2$ at DESY-HERA energies at $Q^2 = 0$ and the ratio is a Q^2 -dependent quantity as the electroproduction cross sections are considered [4]. The question generated intense debate a long time ago [5,6] and it was found for instance in Ref. [7] that the combination of the energy dependence of the dipole cross section and the node of the radial wave function of $2S$ states leads to an anomalous Q^2 and energy dependence of diffractive production of $2S$ vector mesons. In addition, such anomaly appears also in the t dependence of the differential cross section of radially excited $2S$ light vector mesons [8], which is in contradiction with the usual monotonical behavior of corresponding $1S$ states.

Here, we focus on the photoproduction of radially excited vector mesons off nuclei in heavy ion relativistic collisions. In particular, we analyze the exclusive photoproduction of

ψ' off nuclei, $\gamma A \rightarrow \psi(2S)X$, where for coherent scattering one has $X = A$ whereas for incoherent case $X = A^*$ with A^* being an excited state of the A -nucleon system. The theoretical framework used in the present work is the light-cone dipole formalism. In such framework, the $c\bar{c}$ fluctuation of the incoming quasireal photon interacts with the nucleus target via the dipole cross section and the result is projected on the wave function of the observed hadron. In the energy regime we are interested, the dipole cross section depends on the gluon distribution in the target and nuclear shadowing of the gluon distribution is expected to reduce it compared to a proton target. Moreover, theoretically at high energies one expects the transition of the regime described by the linear dynamics, where only the parton emissions are considered, to a new regime where the physical process of recombination of partons becomes important in the parton cascade and the evolution is given by a nonlinear evolution equation (for recent reviews on the topic see Ref. [9]). This regime is characterized by the limitation on the maximum phase-space parton density that can be reached in the hadron wave function, the so-called parton saturation. The transition is specified by a typical scale, the so called saturation scale Q_{sat} [9], which is energy dependent.

Recently, the ALICE Collaboration has measured the diffractive $\psi(1S)$ vector meson production at a relatively large rapidity $y \simeq 3$ [10] and central rapidities [11] as well in the $\sqrt{s} = 2.76$ TeV run, which opens the possibility of investigating small- x physics with heavy nuclei. In addition, the incoherent $\psi(1S)$ cross section has been also measured [11]. This is interesting as the saturation is enhanced for nuclear targets, i.e., $Q_{\text{sat}} \propto A^{1/3}$. The LHCb Collaboration has also measured the cross section in proton-proton collisions at $\sqrt{s} = 7$ TeV of exclusive dimuon final states, including the $\psi(2S)$ state [12]. The ratio at forward rapidity $2.0 \leq \eta_{\mu^\pm} \leq 4.5$ in that case is $\sigma(\psi(2S))/\sigma(\psi(1S)) = 0.19 \pm 0.04$, which is still consistent to the color dipole approach formalism. Therefore, an investigation on the $\psi(2S)$ photoproduction in PbPb collisions at the CERN Large Hadron Collider (LHC) is interesting by itself and timely.

This paper is organized as follows. In the next section we present a brief review of the diffractive photoproduction of vector mesons in electromagnetic reactions in nucleus-nucleus collisions focusing on the PbPb reactions at the LHC energy

regime. In Sec. III we show our predictions for the $\psi(2S)$ photoproduction cross section including the coherent and incoherent contributions. We also check the compatibility with the recent measurements of the $\psi(1S)$ state [10,11]. Moreover, we compare the current results to related approaches available in the literature. Finally, in Sec. IV we summarize our main results and conclusions.

II. PHOTON-POMERON PROCESS IN RELATIVISTIC NUCLEUS-NUCLEUS COLLISIONS

The electromagnetic interaction is dominant in the nucleus-nucleus interaction at large impact parameter and at ultrarelativistic energies. In heavy ion collisions, the heavy nuclei give rise to strong electromagnetic fields due to the coherent action of all protons in the nucleus, which can interact with each other. Accordingly, the total cross section for a given process can be factorized in terms of the equivalent flux of photons of the hadron projectile and the photon-photon or photon-target production cross section [13]. In what follows our main focus shall be in photon-hadron processes which is relevant for the photoproduction of radially excited vector mesons. Considering the requirement that photoproduction is not accompanied by hadronic interaction an analytic expression for the equivalent photon flux of a nuclei can be calculated [13]:

$$\frac{dN_\gamma(\omega)}{d\omega} = \frac{2Z^2\alpha_{em}}{\pi\omega} \left[\xi_R^{AA} K_0(\xi_R^{AA}) K_1(\xi_R^{AA}) - \frac{(\xi_R^{AA})^2}{2} K_1^2(\xi_R^{AA}) - K_0^2(\xi_R^{AA}) \right], \quad (1)$$

where ω is the photon energy, γ_L is the Lorentz boost of a single beam, and $K_0(\xi)$ and $K_1(\xi)$ are the modified Bessel functions. Considering symmetric nuclei having radius R_A , one has $\xi_R^{AA} = 2R_A\omega/\gamma_L$.

The cross section for the photoproduction of ψ' off nuclei in heavy ion relativistic collisions is given by

$$\sigma(AA \rightarrow \psi(2S)X) = \int_{\omega_{min}}^{\infty} d\omega \int dt \frac{dN_\gamma(\omega)}{d\omega} \frac{d\sigma}{dt}(W_{\gamma A}, t), \quad (2)$$

where $\frac{d\sigma}{dt}$ is the differential cross section for the process $\gamma A \rightarrow \psi' X$, $\omega_{min} = M_\psi^2/4\gamma_L m_p$, $W_{\gamma p}^2 = 2\omega\sqrt{S_{NN}}$ and $\sqrt{S_{NN}}$ is the c.m.s. energy of the nucleus-nucleus system. Since photon emission is coherent over the entire nucleus and the photon is colorless we expect that the events to be characterized by $X = A$ and two rapidity gaps in the case of coherent process. In the incoherent process, $X = A^*$ (excited nucleus state) as already mentioned.

The rapidity distribution y for quarkonium photoproduction in nucleus-nucleus collisions can be also computed directly from Eq. (2), by using its relation with the photon energy ω , i.e., $y \propto \ln(2\omega/m_X)$. Explicitly, the rapidity distribution is written as

$$\frac{d\sigma[AA \rightarrow A \otimes \psi(2S) \otimes X]}{dy} = \omega \frac{dN_\gamma(\omega)}{d\omega} \sigma_{\gamma A \rightarrow \psi(2S)X}(\omega), \quad (3)$$

where \otimes represents the presence of a rapidity gap. Consequently, given the photon flux, the rapidity distribution is thus a direct measure of the photoproduction cross section for a given energy.

Let us consider photon-nucleus scattering in the light-cone dipole frame, in which most of the energy is carried by the hadron, while the photon has just enough energy to dissociate into a quark-antiquark pair before the scattering. In this representation the probing projectile fluctuates into a quark-antiquark pair (a dipole) with transverse separation \mathbf{r} long after the interaction, which then scatters off the hadron [1]. In the dipole picture the amplitude for vector meson production off nucleons reads as (see, e.g., Refs. [1,14])

$$\mathcal{A}(x, Q^2, \Delta) = \sum_{h,\bar{h}} \int dz d^2\mathbf{r} \Psi_{h,\bar{h}}^\gamma \mathcal{A}_{q\bar{q}} \Psi_{h,\bar{h}}^{V*}, \quad (4)$$

where $\Psi_{h,\bar{h}}^\gamma(z, \mathbf{r}, Q^2)$ and $\Psi_{h,\bar{h}}^V(z, \mathbf{r})$ are the light-cone wave functions of the photon and of the vector meson, respectively. The quark and antiquark helicities are labeled by h and \bar{h} , variable \mathbf{r} defines the relative transverse separation of the pair (dipole), $z(1-z)$ is the longitudinal momentum fractions of the quark (antiquark). The quantity Δ denotes the transverse momentum lost by the outgoing proton ($t = -\Delta^2$) and x is the Bjorken variable. Moreover, $\mathcal{A}_{q\bar{q}}$ is the elementary amplitude for the scattering of a dipole of size \mathbf{r} on the target. In a compact notation, the nonforward amplitude and the differential cross section for exclusive production of charmonia (or other final state) off a nucleon target, respectively, are given by

$$\mathcal{A}(x, Q^2, \Delta) = \langle \Psi^V | \mathcal{A}_{q\bar{q}}(x, \mathbf{r}, \Delta) | \Psi^\gamma \rangle, \quad (5)$$

$$\frac{d\sigma(s, Q^2)}{dt} = \frac{1}{16\pi} |\mathcal{A}(x, Q^2, \Delta)|^2. \quad (6)$$

In the numerical calculation shown in the next section, the corrections due to the skewedness effect (off-diagonal gluon exchange) and real part of amplitude are also taken into account. Details on the model dependence on these corrections can be found for instance in Ref. [15].

The photon wave functions appearing in Eq. (4) are relatively well known [14]. Concerning the meson wave function, in the current calculation we consider the boosted gaussian wave function:

$$\psi_{\lambda, h\bar{h}}^{nS} = \sqrt{\frac{N_c}{4\pi}} \frac{\sqrt{2}}{z(1-z)} \{ \delta_{h,\bar{h}} \delta_{\lambda, 2h} m_c + i(2h) \delta_{h,-\bar{h}} e^{i\lambda\phi} \times [(1-z)\delta_{\lambda,-2h} + z\delta_{\lambda,2h}] \partial_r \} \phi_{nS}(z, r). \quad (7)$$

Here, $\phi(z, r)$ in the mixed (r, z) representation is obtained by boosting a Schrödinger gaussian wave function in momentum representation, $\Psi(z, \mathbf{k})$. In this case, one obtains the following expression for the 1S state [16]:

$$\phi_{1S}(r, z) = N_T^{(1S)} \left\{ 4z(1-z) \sqrt{2\pi R_{1S}^2} \exp \left[-\frac{m_q^2 R_{1S}^2}{8z(1-z)} \right] \times \exp \left[-\frac{2z(1-z)r^2}{R_{1S}^2} \right] \exp \left[\frac{m_q^2 R_{1S}^2}{2} \right] \right\}, \quad (8)$$

where for the $1S$ ground state vector mesons we determine the parameters R_{1S}^2 and N_T by considering the normalization property of wave functions and the predicted decay widths.

The radial wave function of the $\psi(2S)$ is obtained by the following modification of the $1S$ state [2]:

$$\begin{aligned} \phi_{2S}(r, z) &= N_T^{(2S)} \left\{ 4z(1-z) \sqrt{2\pi R_{2S}^2} \exp \left[-\frac{m_q^2 R_{2S}^2}{8z(1-z)} \right] \right. \\ &\quad \times \exp \left[-\frac{2z(1-z)r^2}{R_{2S}^2} \right] \exp \left[\frac{m_q^2 R_{2S}^2}{2} \right] (1 - \hat{\phi}) \left. \right\}, \\ \hat{\phi} &= \alpha \left[1 + m_q^2 R_{2S}^2 - \frac{m_q^2 R_{2S}^2}{4z(1-z)} + \frac{4z(1-z)}{R_{2S}^2} r^2 \right], \end{aligned} \quad (9)$$

with the new parameter α controlling the position of the node. In addition, the two parameters α and R_{2S} are determined from the orthogonality conditions for the meson wave function. See, for instance, the example for the determination of parameters for the Υ photoproduction in Ref. [17].

At this point some comments are in order. First, we are using a particular choice for the meson wave functions, Eqs. (8) and (9). The boosted gaussian wave function considered here is a simplification of the NNPZ wave function presented in Refs. [1,2]. It has been compared to recent analysis of DESY-HERA data for vector meson exclusive processes. For instance, in Ref. [16] the boosted gaussian wave function was successfully compared to the light mesons and J/ψ production data. In Ref. [17], the production of $\Upsilon(1S)$ and its excited states [$\Upsilon(2S)$, $\Upsilon(3S)$] was investigated and very good agreement with DESY-HERA data was found. Another point to be analyzed is the role played by the node effect to describe the measured ratio $\sigma(\psi')/\sigma(J/\psi)$ in the photoproduction case. Such a ratio is sensitive to the time-scale of the production process. In the dipole approach the interactions occur during the period where the color dipole is compact having a transverse size $r \simeq 1/m_q$ and the production cross section is proportional to the square of the quarkonium wave function at origin, $\sigma \propto |\phi(0)|^2$. On the other hand, further interactions depend on the wave function profile for transverse sizes larger than $r_B = \mathcal{O}(1/\alpha_s m_q)$, the so-called Bohr radius. In exclusive charmonia electroproduction at relatively large Q^2 the dipole size is of order $1/Q^2 \ll r_B$ and the cross section is predicted to be proportional to $|\phi_n(0)|^2$. This leads the ratio to be of order $|\phi_{2S}(0)|^2/|\phi_{1S}(0)|^2 \simeq 0.6$ at large Q^2 whereas the measured value in photoproduction is around 0.16 [4]. It was determined long time ago in Refs. [1,2] that the moderate value of charm mass and the dominant color transparency behavior of dipole cross section $\sigma_{dip} \propto r^2$ imply the amplitude to probe the meson wave function at a transverse size around r_B . This fact reduces the $\psi(2S)$ contribution due to the node in its wave function and correctly describes the measured DESY-HERA ratio. Along these lines, it was explicitly shown in Ref. [6] that at $Q^2 \rightarrow 0$ the leading logarithmic approximation $rl_\perp \ll 1$, which gives the usual $\sigma_{dip} \propto r^2$, is not able to provide alone the correct value for the ratio ψ'/ψ . Here, l_\perp is the exchanged gluon transverse momentum in a two-gluon exchange model. Therefore, important contributions come from the overlap of the large-sized color dipole configurations and the $\psi(2S)$ wave

function. Thus, despite the leading logarithmic approximation to be able to describe the J/ψ production cross section the same is not true for the excited states as the ψ' . This is the reason why we will use a model for the dipole cross section that takes into account the correct behavior for large dipole configurations (the transition hard-soft is given by the saturation scale).

The exclusive $\psi(2S)$ photoproduction off nuclei for coherent and incoherent processes can be simply computed in high energies where the large coherence length $l_c \gg R_A$ is fairly valid. In such case the transverse size of $c\bar{c}$ dipole is frozen by Lorentz effects. The expressions for the coherent and incoherent cross sections are given by [18]

$$\sigma_{coh}^{\gamma A} = \int d^2b |\langle \Psi^V | 1 - \exp \left[-\frac{1}{2} \sigma_{dip}(x, \mathbf{r}) T_A(b) \right] | \Psi^\gamma \rangle|^2, \quad (10)$$

$$\begin{aligned} \sigma_{inc}^{\gamma A} &= \frac{1}{16\pi B_V(s)} \int d^2b T_A(b) |\langle \Psi^V | \sigma_{dip}(x, \mathbf{r}) \\ &\quad \times \exp \left[-\frac{1}{2} \sigma_{dip}(x, \mathbf{r}) T_A(b) \right] | \Psi^\gamma \rangle|^2, \end{aligned} \quad (11)$$

where $T_A(b) = \int dz \rho_A(b, z)$ is the nuclear thickness function given by integration of nuclear density along the trajectory at a given impact parameter b . In addition, B_V is the diffractive slope parameter in the reaction $\gamma^* p \rightarrow \psi p$. Here, we consider the energy dependence of the slope using the Regge motivated expression $B_V(W_{\gamma p}) = b_{el}^V + 2\alpha' \log \frac{W_{\gamma p}^2}{W_0^2}$ with $\alpha' = 0.25 \text{ GeV}^{-2}$ and $W_0 = 95 \text{ GeV}$. It is used the measured slopes [4] for $\psi(1S)$ and $\psi(2S)$ at $W_{\gamma p} = 90 \text{ GeV}$, i.e., $b_{el}^{\psi(1S)} = 4.99 \pm 0.41 \text{ GeV}^{-2}$ and $b_{el}^{\psi(2S)} = 4.31 \pm 0.73 \text{ GeV}^{-2}$, respectively.

The last ingredient is the model for the dipole cross section in Eqs. (10) and (11). In our calculation, we consider the color glass condensate model [19] for $\sigma_{dip}(x, r)$. This model has been tested for a long period against DIS, diffractive DIS, and exclusive production processes in ep collisions. In addition, we allow for its renormalization by the effect of the gluon shadowing phenomenon as the gluon density in nuclei at a small- x region is known to be suppressed compared to a free nucleon. That is, we will take $\sigma_{dip} \rightarrow R_G(x, Q^2, b) \sigma_{dip}$ following studies in Ref. [20]. The factor R_G is the nuclear gluon density ratio. In the present investigation we will use the nuclear ratio from the leading twist theory of nuclear shadowing based on generalization of the Gribov-Glauber multiple scattering formalism as investigated in Ref. [21]. We used the two models available for $R_G(x, Q^2)$ in [21], models 1 and 2, which correspond to higher nuclear shadowing and lower nuclear shadowing, respectively. Such a choice is completely arbitrary and other nuclear gluon ratios available in literature could be considered. It would be also interesting to investigate the effect of using the impact parameter dependent nuclear parton distribution ratios. We discuss about this distinct issues in the next section.

III. RESULTS AND DISCUSSIONS

Let us start by checking the present theoretical approach against the recent data for the $1S$ state measured by the ALICE Collaboration at the energy of 2.76 TeV in PbPb collisions at the LHC [10,11]. In Fig. 1 we present the numerical calculations for the rapidity distribution of coherent $\psi(1S)$ state within the color dipole formalism, Eqs. (3) and (10), using distinct scenarios for the nuclear gluon shadowing. The dot-dashed curve represents the result using $R_G = 1$ and it is consistent with previous calculations using the same formalism [15]. It overestimates the ALICE data on the backward (forward) and mainly in central rapidities. In the backward/forward rapidity case, the overestimation is already expected as a proper threshold factor for $x \rightarrow 1$ was not included in the present calculation. In that kinematical region either a small- x photon scatters off a large- x gluon or vice versa. For instance, for $y \simeq \pm 3$ one gets x as large as 0.02. On the other hand, for central rapidity $y = 0$ one can obtain $x = M_V e^{\pm y} / \sqrt{s_{NN}}$ smaller than 10^{-3} for the nuclear gluon distribution. In such a case, considering $R_G = 1$ the ALICE data [11] is overestimated by a factor 2 or so, as already noticed in the recent study of Ref. [22]. The situation is improved if we consider nuclear shadowing renormalizing the dipole cross section. The reason is that the gluon density in nuclei at small Bjorken x is expected to be suppressed compared to a free nucleon due to interferences. For the ratio of the gluon density, $R_G(x, Q^2 = m_V^2/4)$, we have considered the theoretical evaluation of Ref. [21]. There, two scenarios for the gluon shadowing are investigated: model 1 corresponds to a strong gluon shadowing and model 2 concerns a small nuclear shadowing. The consequence of renormalizing the dipole cross section by gluon shadowing effects is represented by the long-dashed (model 1) and solid (model 2) lines, respectively.




FIG. 1. (Color online) The rapidity distribution of coherent $\psi(1S)$ meson photoproduction at $\sqrt{s} = 2.76$ TeV in PbPb collisions at the LHC. The theoretical curves stand for color dipole formalism using $R_G = 1$ (dot-dashed curve) and two scenarios for the nuclear gluon distribution (solid and long-dashed curves, see text). Data from ALICE collaboration [10,11].




FIG. 2. (Color online) The rapidity distribution of coherent $\psi(2S)$ meson photoproduction at $\sqrt{s} = 2.76$ TeV in PbPb collisions at the LHC. The theoretical curves follow the same notation as in the previous figure.

Clearly, the small shadowing option is preferred in the current analysis. It is worth mentioning that the theoretical uncertainty related to the choice of meson wave function is relatively large. As a prediction at central rapidity, one obtains $\frac{d\sigma}{dy}(y = 0) = 4.95, 1.68,$ and 2.27 mb for a calculation using $R_G = 1,$ models 1 and 2, respectively. Here, a word of caution is needed as we are considering R_G as independent of the impact parameter. It has been long known that a b -dependent ratio could give a smaller suppression compared to that presented in our calculation. For instance, in Ref. [20] the suppression is of order 0.85 for the LHC energy and central rapidity.

In Fig. 2 we show our predictions for the coherent photoproduction of the $\psi(2S)$ state. This is the first estimate in literature for the photoproduction of the $2S$ state in nucleus-nucleus collisions. The theoretical predictions follow the general trend as for the $1S$ state, where the notation for the curves is the same as used in Fig. 1. In particular, for $R_G = 1$ one obtains for central rapidity $\frac{d\sigma}{dy}(y = 0) = 0.71$ mb and the following in the forward/backward region $\frac{d\sigma}{dy}(y = \pm 3) = 0.16$ mb. When introducing the suppression in the dipole cross section due to nuclear shadowing one gets instead $\frac{d\sigma}{dy}(y = 0) = 0.24$ mb and 0.33 mb for models 1 and 2, respectively. At central rapidities, the meson state ratio is evaluated to be $R_\psi^{y=0} = \frac{\sigma_{\psi(2S)}}{\sigma_{\psi(1S)}}(y = 0) = 0.14$ in the case of $R_G = 1$ which is consistent with the ratio measured in CDF, i.e., 0.14 ± 0.05 , on the observation of exclusive charmonium production at 1.96 TeV in $p\bar{p}$ collisions [23]. A similar ratio is obtained using models 1 and 2 at central rapidity as well. As a prediction for the planned LHC run in PbPb mode at 5.5 TeV, we obtain $\frac{d\sigma_{coh}}{dy}(y = 0) = 1.27$ mb and $\frac{d\sigma_{inc}}{dy}(y = 0) = 0.27$ mb for the coherent and incoherent $\psi(2S)$ cross sections (upper bound using $R_G = 1$), respectively.

Finally, in Fig. 3 we show the incoherent contribution to the rapidity distribution for both $\psi(1S)$ (solid line) and $\psi(2S)$




FIG. 3. (Color online) The rapidity distribution of incoherent $\psi(1S)$ (solid line) and $\psi(2S)$ (dashed line) meson photoproduction at $\sqrt{s} = 2.76$ TeV in PbPb collisions at the LHC. Data from ALICE collaboration [11].

(dashed line) meson states. The theoretical estimates are done using Eq. (6) taking into account the corresponding diffractive slope for each meson state as discussed in the previous section. For the $\psi(1S)$ state, the present calculation can be directly compared with those studies presented in Ref. [22]. It was found in [22] that the incoherent cross section $\frac{d\sigma_{\text{inc}}}{dy}$ ranges from 0.5 to 0.7 mb (using IIM dipole cross section) or between 0.7 to 0.9 mb (using fIPsat dipole cross section) at central rapidities, with the uncertainty determined by the distinct meson wave function considered. In our case, we obtained $\frac{d\sigma_{\text{inc}}}{dy}(y=0) = 1.1$ mb using a different expression for the incoherent amplitude, Eq. (11). Our result fairly describes the recent ALICE data [11] for the incoherent cross section at midrapidity, $\frac{d\sigma_{\text{ALICE}}}{dy}(-0.9 < y < 0.9) = 0.98 \pm 0.25$ mb. As a prediction for the $\psi(2S)$ state, we have found $\frac{d\sigma_{\text{inc}}}{dy} = 0.16$ mb for central rapidities. In both cases we have only computed the case for $R_G = 1$. Therefore, this gives an upper bound for the incoherent cross section compared to the calculations of models 1 and 2. We notice that for the incoherent case, the gluon shadowing is weaker than the coherent case and the reduction is around 20% compared to the case of $R_G = 1$. As expected, the incoherent piece is quite smaller compared to the main coherent contribution. As an example of order of

magnitude, the ratio incoherent/coherent is a factor 0.22 for the $1S$ state and 0.23 for the $2S$ state at central rapidity.

IV. SUMMARY

We have investigated the photoproduction of radially excited vector mesons off nuclei in heavy ion relativistic collisions as the $\psi(2S)$ charmonium state. The theoretical framework used in the present analysis is the light-cone dipole formalism and predictions are done for PbPb collisions at the CERN-LHC energy of 2.76 TeV. The effect of suppressing of the dipole cross section due to the gluon shadowing was studied and the results for $R_G = 1$ give the larger cross sections. It was found that the coherent exclusive photoproduction of $\psi(2S)$ off nuclei has an upper bound of order 0.71 mb at $y = 0$ down to 0.10 mb for backward/forward rapidities $y = \pm 3$. The incoherent contribution was also computed and it is a factor 0.2 below the coherent one. A comparison has been done to the recent ALICE collaboration data for the $\psi(1S)$ state photoproduction and the analysis shows that a small nuclear shadowing $R_G(x, Q^2 = \frac{m_V^2}{4})$ is preferred in data description whereas the usual $R_G = 1$ value overestimates the central rapidity cross section by a factor 2. On the other hand, the present theoretical approach fairly describes the ALICE data for the incoherent cross section. Thus, the central rapidity data measured by the ALICE collaboration for the rapidity distribution of the $\psi(1S)$ state is crucial to constrain the nuclear gluon function. The cross section for exclusive quarkonium production is proportional to $[\alpha(Q^2)xg_A(x, Q^2)]^2$ in the leading-order pQCD calculations, evaluated at the relevant scale $Q^2 \approx m_V^2/4$ and at momentum fraction $x \simeq 10^{-3}$ in central rapidities. The theoretical uncertainty is large and it has been investigated in several studies [24,25]. Along these line, the authors of Ref. [26] extract the nuclear suppression factor, $S(x \approx 10^{-3}) = 0.61 \pm 0.064$, using the ALICE data on coherent $\psi(1S)$ and considering the nuclear gluon shadowing predicted by nuclear parton distribution functions and by leading twist nuclear shadowing.

ACKNOWLEDGMENTS

This work was partially financed by the Brazilian funding agencies CNPq and FAPERGS and by the French-Brazilian scientific cooperation Project No. CAPES-COFECUB 744/12. M.V.T.M. thanks Magdalena Malek, Heikki Mäntysaari, and Daniel Tapia Takaki for helpful comments.

- [1] N. N. Nikolaev and B. G. Zakharov, *Phys. Lett. B* **332**, 184 (1994); *Z. Phys. C* **64**, 631 (1994).
 [2] J. Nemchik, N. N. Nikolaev, E. Predazzi, and B. G. Zakharov, *Phys. Lett. B* **374**, 199 (1996).
 [3] J. Nemchik, *Phys. Rev. D* **63**, 074007 (2001).
 [4] C. Adloff *et al.* (H1 Collaboration), *Phys. Lett. B* **541**, 251 (2002).

- [5] P. Hoyer and S. Peigne, *Phys. Rev. D* **61**, 031501 (2000).
 [6] K. Suzuki, A. Hayashigaki, K. Itakura, J. Alam, and T. Hatsuda, *Phys. Rev. D* **62**, 031501 (2000).
 [7] J. Nemchik, N. N. Nikolaev, E. Predazzi, and B. G. Zakharov, *Z. Phys. C* **75**, 71 (1997).
 [8] J. Nemchik, *Eur. Phys. J. C* **18**, 711 (2001).

- [9] F. Gelis, E. Iancu, J. Jalilian-Marian, and R. Venugopalan, *Ann. Rev. Nucl. Part. Sci.* **60**, 463 (2010); H. Weigert, *Prog. Part. Nucl. Phys.* **55**, 461 (2005); J. Jalilian-Marian and Y. V. Kovchegov, *ibid.* **56**, 104 (2006).
- [10] B. Abelev *et al.* (ALICE Collaboration), *Phys. Lett. B* **718**, 1273 (2013).
- [11] E. Abbas *et al.* (ALICE Collaboration), arXiv:1305.1467 [nucl-ex].
- [12] R. Aaij *et al.* (LHCb Collaboration), *J. Phys. G* **40**, 045001 (2013).
- [13] G. Baur, K. Hencken, D. Trautmann, S. Sadovsky, and Y. Kharlov, *Phys. Rep.* **364**, 359 (2002); C. A. Bertulani, S. R. Klein, and J. Nystrand, *Ann. Rev. Nucl. Part. Sci.* **55**, 271 (2005).
- [14] A. C. Caldwell and M. S. Soares, *Nucl. Phys. A* **696**, 125 (2001); H. Kowalski and D. Teaney, *Phys. Rev. D* **68**, 114005 (2003); J. R. Forshaw, R. Sandapen, and G. Shaw, *ibid.* **69**, 094013 (2004); C. Marquet, R. Peschanski, and G. Soyez, *ibid.* **76**, 034011 (2007); H. Kowalski, L. Motyka, and G. Watt, *ibid.* **74**, 074016 (2006).
- [15] V. P. Goncalves and M. V. T. Machado, *Phys. Rev. C* **84**, 011902(R) (2011).
- [16] J. R. Forshaw, R. Sandapen, and G. Shaw, *J. High Energy Phys.* **11** (2006) 025.
- [17] B. E. Cox, J. R. Forshaw, and R. Sandapen, *J. High Energy Phys.* **06** (2009) 034.
- [18] B. Z. Kopeliovich and B. G. Zakharov, *Phys. Rev. D* **44**, 3466 (1991).
- [19] E. Iancu, K. Itakura, and S. Munier, *Phys. Lett. B* **590**, 199 (2004).
- [20] Y.P. Ivanov, B. Z. Kopeliovich, A. V. Tarasov, and J. Hufner, *Phys. Rev. C* **66**, 024903 (2002).
- [21] L. Frankfurt, V. Guzey, and M. Strikman, *Phys. Rep.* **512**, 255 (2012).
- [22] T. Lappi and H. Mantysaari, *Phys. Rev. C* **87**, 032201(R) (2013).
- [23] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **102**, 242001 (2009).
- [24] A. L. Ayala Filho, V. P. Goncalves, and M. T. Griep, *Phys. Rev. C* **78**, 044904 (2008).
- [25] A. Adeluyi and C. A. Bertulani, *Phys. Rev. C* **85**, 044904 (2012).
- [26] V. Guzey, E. Kryshen, M. Strikman, and M. Zhalov, arXiv:1305.1724 [hep-ph].