

Stability and self-organization of planetary systems

Renato Pakter and Yan Levin

Instituto de Física, UFRGS, Caixa Postal 15051, CEP 91501-970, Porto Alegre, RS, Brazil
 (Received 1 November 2017; revised manuscript received 26 February 2018; published 30 April 2018)

We show that stability of planetary systems is intimately connected with their internal order. An arbitrary initial distribution of planets is susceptible to catastrophic events in which planets either collide or are ejected from the planetary system. These instabilities are a fundamental consequence of chaotic dynamics and of Arnold diffusion characteristic of many body gravitational interactions. To ensure stability over astronomical time scale of a *realistic* planetary system—in which planets have masses comparable to those of planets in the solar system—the motion must be quasiperiodic. A dynamical mechanism is proposed which naturally evolves a planetary system to a quasiperiodic state from an arbitrary initial condition. A planetary self-organization predicted by the theory is similar to the one found in our solar system.

DOI: [10.1103/PhysRevE.97.042221](https://doi.org/10.1103/PhysRevE.97.042221)

I. INTRODUCTION

Stability of the solar system has been a subject of great debate since the time that Isaac Newton first wrote his famous law of universal gravitation [1]. Newton realized that the interaction between planets will perturb their orbits from a simple elliptical shape postulated by Kepler. Over millions of years these small perturbations will accumulate leading to catastrophic events such as collisions between planets or ejection of planets from the solar system [2]. Newton's solution to this difficulty was to invoke divine intervention in which God would adjust the planetary orbits to keep the solar system stable. Newton's contemporary, Leibnitz, objected strongly to Newton's theological solution: "Was God an inferior watchmaker," he demanded, "who could not get things right from the beginning?" [3] Over the centuries the question of stability has attracted the attention of both physicists and mathematicians, without a definite solution. An apparently unrelated puzzle is a seeming regularity of spacing of planetary orbits [4]; see Fig. 1. In 1766 Titius noticed that the orbits of then known planets followed a geometric sequence, if a "missing" planet was inserted between Mars and Jupiter [5]. Subsequent discovery of the asteroid belt and of Uranus at the positions predicted by the Titius-Bode "law" gave further credence to the belief that there is a hidden order in the solar system. The modern version of the Titius-Bode law can be written as $r_n = r_0 1.7^n$, with $n = 1$ being Mercury, $n = 2$ Venus, $n = 3$ Earth, etc.; see Fig. 1. The normalization $r_0 = 0.2294$ a.u. was chosen so that $n = 1$ corresponds exactly to the orbit of Mercury in astronomical units (AU) [6].

In this paper we will show that stability of planetary systems is intimately connected with their internal order. The dynamical simulations demonstrate that a generic arrangement of planets is unstable to small perturbations resulting from interplanetary interactions which lead to catastrophic events. We argue that a planetary system will remain stable over astronomical time scales only if its dynamics is quasiperiodic. Indeed various near commensurabilities have been observed for satellites and planets in the solar system [7,8] and already in 1970 Hill

suggested that there is a dynamical origin to the Titius-Bode law [9]. In this paper we will argue that a specific requirement of quasiperiodicity results in a planetary distribution almost identical to the one observed in the solar system. Furthermore, we will provide a dynamical mechanism that leads to spontaneous self-organization of a planetary system into a periodic state. This is different from other approaches used previously to explain mass distribution in the solar system which rely either on statistics or hydrodynamic instabilities [6,10–12]. Even if such approaches can account for geometrical progression of planetary distances, they cannot explain stability of planetary systems. The Titius-Bode law is not a condition of stability, but rather a consequence of self-organization, as will be demonstrated in the present paper. Finally we should mention that according to Kolmogorov-Arnold-Moser (KAM) theorem there is a dense set of initial conditions for a many body gravitational system that lead to stable quasiperiodic trajectories. However, as was already demonstrated by Henon in 1966, the KAM stability only applies to planetary systems with unrealistically small planetary masses of less than 10^{-320} of solar mass [13]. A generic initial condition for a realistic planetary system will, therefore, result in a chaotic dynamics, such that in the infinite time limit planets will either collide or will be ejected from the system.

II. THEORY

The stability of planetary systems is an outstanding problem. Since Newton's gravitational potential is bound from above and is unbound from below [14,15], orbits of planets are in general unstable—some planets can gain enough kinetic energy to escape altogether from the planetary system [16,17], while others will fall into the sun or collide with each other; see Fig. 2. This type of instability driven by chaos and Arnold diffusion [18] is a fundamental characteristic of many body celestial dynamics. On the other hand, it is possible to find very special initial conditions—corresponding to a set of measure zero, since the KAM theorem does not apply to realistic




FIG. 1. Mean radii of planets in our solar system on a semilog scale: (Me) Mercury, (V) Venus, (E) Earth, (Ma) Mars, (Ast) Asteroid belt, (J) Jupiter, (S) Saturn, (U) Uranus, (N) Neptune, and (P) Pluto. Straight line is the modern version of Titius-Bode law.

planetary systems [13]—for which the dynamics of a nonlinear interacting system is purely periodic. These periodic solutions will persist indefinitely, with the relative configuration of planets repeating itself after a synodic period of time T . The fundamental difficulty is to obtain initial conditions which lead to periodic dynamics of a fully interacting gravitational system. Furthermore, since the probability that a planetary system will be “born” with planets at precisely the correct positions is highly improbable, there must be a mechanism that makes a planetary system evolve towards a stable periodic orbit. Motivated by the theories of control of chaos [19,20], we suggest that periodic orbits can be stabilized by energy nonconserving perturbations [21]. Such perturbations could have originated from the interaction of newborn planetesimals with the gas and dust of the protoplanetary disk. The angular frequency of dust particles at a distance r from the star has a

simple Keplerian form [22,23]

$$\omega(r) = \sqrt{\frac{GM}{r^3}}, \quad (1)$$

where M is the star mass and G is Newton’s gravitational constant. For concreteness we will take the star mass to be that of our sun and will measure all the distances in astronomical units and time in Earth years. To simplify the calculations we will suppose that all the planetary orbits are restricted to the ecliptic plane and all planets have the same mass m . To speed up the simulations we will take the planetary mass m to be a few times that of Jupiter. We will see, however, that our conclusions do not depend on the planetary mass as long as it is much smaller than the star mass.

Newton’s equations of motion for the coordinates x and y of a planet i are

$$\begin{aligned} m\ddot{x}_i &= -\frac{GMmx_i}{r_i^3} - \sum_j \frac{Gm^2(x_i - x_j)}{r_{ij}^3} - f_i^\theta \frac{y_i}{r_i}, \\ m\ddot{y}_i &= -\frac{GMmy_i}{r_i^3} - \sum_j \frac{Gm^2(y_i - y_j)}{r_{ij}^3} + f_i^\theta \frac{x_i}{r_i}, \end{aligned} \quad (2)$$

where $r_i = \sqrt{x_i^2 + y_i^2}$ is the distance of i th planet from the star and $r_{ij} = \sqrt{(x_i - x_j)^2 + (y_i - y_j)^2}$ is the separation between planet i and planet j . The purely angular force $\mathbf{f}_i^\theta = f_i^\theta \hat{\theta}$, where $\hat{\theta}$ is the unit angular vector, is responsible for the interaction of a planet i with the residual dust of the protoplanetary disk. Here we will use a simple phenomenological expression for such nonconservative force: if the angular velocity of a planet is lower than the Keplerian velocity of the surrounding dust in the same orbit, the planet will gain energy from dust; if the planetary velocity is larger than the velocity of dust, it will lose energy. The simplest possible mathematical expression with this characteristic is

$$f_i^\theta = -\beta(r_i\omega(r_i) - v_i^\theta) \left[L - m \sum_i v_i^\theta r_i \right], \quad (3)$$

where β is a small phenomenological constant which controls the interaction between dust and planets, $v_i^{\text{dust}} = r_i\omega(r_i)$ is the angular velocity of dust at the location of planet i , and v_i^θ is the angular velocity of the planet i ,

$$v_i^\theta = \frac{\dot{y}_i x_i - y_i \dot{x}_i}{r_i}. \quad (4)$$

The expression Eq. (3) is analogous to viscous dissipation of an object in a rotating fluid. The term in square brackets of Eq. (3) is included so that the nonconservative force “turns off” when the net planetary angular momentum reaches a predetermined value L . This is designed to model a progressive depletion of dust or gas from the protoplanetary disk which will result in a continuously decreasing value of f^θ . In our simulations we used $\beta = 10^{-3} - 10^{-2} \text{ AU}^{-2}$. To integrate the equations numerically we employed a Runge-Kutta algorithm with adaptive time step that uses embedded fifth order and sixth order Runge-Kutta estimates to compute the solution and the relative error [24]. To speed up the simulation and avoid numerical instabilities the singular form of the Newton’s




FIG. 2. Radial coordinate evolution of eight planets of mass $0.005M$ originally distributed uniformly between 1 and 8 AU in their respective Keplerian orbits. After a short time we start seeing catastrophic events (the almost vertical trajectories) in which planets begin to be ejected from the planetary system. The resulting planetary system remains with only two planets.




FIG. 3. Temporal evolution of radial coordinates of three planets under the action of Eq. (3). The dynamics is very complex with two of the planets switching their relative order in the sequence.




FIG. 4. Temporal evolution of the radial coordinates of the three planet system of Fig. 3 after a self-organized periodic state has been established.

gravitational potential between the planets is regularized to

$$V(r) = -\frac{(2d^3 - 2dr^2 + r^3)Gm^2}{d^4} \quad \text{for } r \leq d, \tag{5}$$

$$V(r) = -\frac{Gm^2}{r} \quad \text{for } r > d,$$

where d is an arbitrary short distance cutoff whose precise value does not affect our conclusions. In our simulations we used d on the order of 10^{-3} a.u. With this modification, we do not need to introduce any specific “collision model,” since the integrator will have enough resolution to deal with the fast dynamics resulting from catastrophic planetary collisions. Indeed, the dynamics of planets interacting through Eq. (5) can result in planetary collision in which two or more trajectories merge into one. We shall call such events “planetary coalescence,” which are analogous to nonelastic collisions. The advantage of regularized Newton potential Eq. (5) is that such singular events can be handled using the adaptive step size Runge-Kutta integrator without any numerical instabilities.

We stress that our goal is to find the simplest possible mechanism that can drive a planetary system towards a stable quasiperiodic state. During the formation of the solar system, more complicated gravitational, electromagnetic, and collisional processes have certainly taken place. Here, however, our aim is to provide a proof of concept that energy nonconserving perturbation can drive a planetary system into a self-organized quasiperiodic state. In the simulations we observe that the angular force given by Eq. (3) leads to a self-organized periodic state if $\beta \rightarrow 0$ and $t \rightarrow \infty$. In practice, however, the simulation time is finite, so that β cannot be too small. In Fig. 3 we show the dynamical evolution described by Eqs. (2) of a system with three planets, initially placed in Keplerian orbits uniformly distributed from 1 AU to 3 AU.

We see that the dynamics is very complex, but after $t = 10^4$ yr, this planetary system relaxes to a periodic state with synodic period $T = 1.6$ yr, in which adjacent planetary orbits exhibit a perfect synchronization, with the time between two consecutive perihelions—anomalous period—increasing in the ratio of 2:1; see Fig. 4. The mean orbital distance of each

planet in this synchronized state follows the Titius-Bode law $r_n \sim 1.69^n$; see Fig. 5.

In Fig. 6 we show a periodic orbit to which a planetary system with four planets evolves. From the figure we see that the orbits of planets once again exhibit a perfect synchronization, with the anomalous period of adjacent planets increasing in the ratio of 2:1. To make this observation more quantitative we define a radial and an angular winding number:

$$\omega_i^r = \lim_{t \rightarrow \infty} \frac{N_{\text{peri}}(t)}{t}, \tag{6}$$

$$\omega_i^\theta = \lim_{t \rightarrow \infty} \frac{N_\theta(t)}{t},$$

where $N_{\text{peri}}(t)$ is the number of times that the orbit of planet i passes through a perihelion during a time interval t , and $N_\theta(t)$ is the number of times that the planet i completes a full rotation around the star. It is important to keep in mind that because of the interplanetary gravitational attraction the dynamics is very complicated, with the orbit of a planet precessing around the star. The planetary year, therefore, will not be equal to the anomalous period, amount of time between two consecutive




FIG. 5. Mean radial distance from the star of three planets for different initial conditions (symbols), after the self-organized periodic state is established. In all cases the final planetary distribution follows the Titius-Bode law $r_n \sim 1.69^n$, shown by parallel lines.




FIG. 6. Temporal evolution of radial coordinates of planets in a self-organized periodic planetary system with four planets. Note that the orbits of adjacent planets exhibit a perfect 2:1 synchronization.

perihelions. This dichotomy is clearly demonstrated by the normalized radial and angular winding numbers (winding number of planet i divided by the winding number of the outermost planet), see Table I, which shows a perfect 2:1 synchronization of anomalistic periods of adjacent orbits, but no synchronization of rotational periods. Furthermore, Fig. 7 shows that the mean planetary distances from the star in this self-organized planetary system follow a geometric progression—Titius-Bode law— $r_n \sim 1.67^n$, very similar to the one observed in our solar system. The synodic period for this planetary system is $T = 2.2$ yr.

We next study a planetary system which starts with nine planets of mass $m = 0.002M$ uniformly distributed from 1 AU to 9 AU. After a period of dynamical evolution this system once again self-organized into a complex periodic structure. During the evolution, planets 1 and 2 collided producing a new planet of mass $2m$. This planet, in turn, formed a binary with the planet 3. Planets 4 and 5 also collided forming a new planet of mass $2m$, as did planets 6 and 7. The resulting planetary system contains six planets with synodic period $T = 6.5$ yr. In spite of a distinct mass distribution and a very complex dynamics, the planetary distances are found to, once again, follow a geometric progression $r_n \sim 1.63^n$; see Fig. 8. Furthermore, the radial winding numbers indicate a perfect 2:1 synchronization between anomalistic periods, while the orbital periods do not show any clear structure; see Table II. Finally, we should note that it is very difficult to obtain large planetary systems, even if we start with a very large number of planetesimals; very quickly most of them will either fall into the sun, coalesce, or will be ejected. The final planetary system will have only a small number of planets. In spite of a diligent effort we were

TABLE I. Normalized radial and angular winding numbers of a four planet system.

Planet	1	2	3	4
ω_r	1.000	2.000	4.000	8.000
ω_θ	1.000	2.275	4.827	9.929




FIG. 7. Mean radial distance of four planets from the star, which agrees precisely with the Titius-Bode law, the solid line.

not able to find a stable planetary system with more than six planets.

III. CONCLUSIONS

We have shown that stability of planetary systems is intimately connected with the orbital arrangement of planets. An arbitrary initial distribution of planets is susceptible to catastrophic events in which planets are ejected from the planetary system or collide with each other. These catastrophic events are an unavoidable consequence of chaotic dynamics and of Arnold diffusion characteristic of celestial mechanics in the $t \rightarrow \infty$ limit. We note that even if the planets are placed at radial positions consistent with the Titius-Bode law, a planetary system will still, in general, be unstable unless the orbits of planets are properly synchronized and the dynamics is periodic. In this paper we presented a mechanism which leads to self-organization of a planetary system into a stable periodic state. The mechanism proposed is probably not unique and should rather be viewed as a proof of concept which demonstrates that energy nonconserving perturbations can drive a planetary system into a self-organized periodic state




FIG. 8. Mean radial planetary distance from the star in a self-organized system which originally contained nine planets. Various planets have collided forming new planets of mass $2m$, so that only five distinct radial positions appear in the plot. These follow the Titius-Bode distribution.

TABLE II. Radial and angular winding numbers of a self-organized system with nine planets.

Planet	1	2	3	4	5	6	7	8	9
ω_r	1.000	1.000	1.000	2.000	2.000	4.000	4.000	8.000	16.000
ω_θ	1.000	1.000	1.000	2.146	2.146	4.439	4.439	9.024	18.195

from an *arbitrary* initial condition. In such a state anomalistic periods between all planets are synchronized, while orbital periods do not indicate any synchronous structure. When the anomalistic periods between the radially adjacent planets are synchronized in the 2:1 ratio, the mean orbital distance is found to follow a geometric progression, $r_n \sim 1.7^n$; the same as the one observed in our solar system. In principle, however, there is no *a priori* reason why all the planets should have 2:1 synchronization, and indeed other synchronized states are possible. For such planetary systems the Titius-Bode law will not be valid. We stress again that it is the anomalistic, and not the orbital periods which show synchronous behavior in the self-organized state. Indeed, if orbital periods would be synchronized, the explanation of the Titius-Bode law would be quite straightforward. For small planetary masses, the planetary year is related to the length of the semimajor axis

through Kepler's law $T^2 \sim a^3$; if the planetary years would be locked in 2:1 resonance, the ratio of a semimajor axis would then follow a geometric progression $r_n \sim 2^{2n/3} \approx 1.5874^n$, which is very similar to the observed Titius-Bode law. However, Tables I and II show that there is no synchrony of orbital periods and only anomalistic periods that are synchronized. Therefore, this simple argument cannot be used to account for the orbital structure inside the self-organized state observed in our simulations.

ACKNOWLEDGMENTS

This work was partially supported by the CNPq, National Institute of Science and Technology Complex Fluids INCT-FCx, and by the US-AFOSR under Grant No. FA9550-16-1-0280.

-
- [1] J. Laskar, Is the solar system stable? in *Chaos*, Progress in Mathematical Physics, edited by B. Duplantier, S. Nonnenmacher, and V. Rivasseau, Vol. 66 (Birkhäuser, Basel, 2013), pp. 239–270.
 - [2] I. Newton, *Opticks: Or, A Treatise of the Reflections, Refractions, Inflexions and Colours of Light*, 4th ed. (Dover Publications Inc., New York, 1952).
 - [3] G. S. Morson, *Narrative, Philosophy and Life* (Springer, Boston, 2014), p. 123.
 - [4] W. Hayes and S. Tremain, *Icarus* **135**, 549 (1998).
 - [5] M. M. Nieto, *The Titius-Bode Law of Planetary Distances* (Pergamon, London, 1972).
 - [6] F. Graner and B. Dubrulle, *Astron. Astrophys.* **282**, 269 (1994).
 - [7] P. Goldreich, *Mon. Not. R. Astron. Soc.* **130**, 159 (1964).
 - [8] S. J. Peale, *Annu. Rev. Astron. Astrophys.* **14**, 215 (1976).
 - [9] J. G. Hill, *Nature (London)* **225**, 840 (1970).
 - [10] J. Laskar, *Phys. Rev. Lett.* **84**, 3240 (2000).
 - [11] C. Hernández-Mena, *Mon. Not. R. Astron. Soc.* **412**, 95 (2011).
 - [12] J. Laskar and A. C. Petit, *Astron. Astrophys.* **605**, A72 (2017).
 - [13] J. Laskar, Michel henon and the stability of the solar system, [arXiv:1411.4930](https://arxiv.org/abs/1411.4930).
 - [14] Y. Levin, R. Pakter, F. B. Rizzato, T. N. Telles, and F. P. C. Benetti, *Phys. Rep.* **535**, 1 (2014).
 - [15] F. P. C. Benetti, A. C. Ribeiro-Teixeira, R. Pakter, and Y. Levin, *Phys. Rev. Lett.* **113**, 100602 (2014).
 - [16] J. E. Chambers, G. W. Wetherill, and A. P. Boss, *Icarus* **119**, 261 (1996).
 - [17] B. A. Conway and T. J. Elsner, *Long-Term Dynamical Behavior of Natural and Artificial N-body Systems* (Kluwer Academic, Amsterdam, 1988).
 - [18] V. I. Arnold, *Sov. Math.* **5**, 581 (1964).
 - [19] E. Ott, C. Grebogi, and J. A. Yorke, *Phys. Rev. Lett.* **64**, 1196 (1990).
 - [20] K. Pyragas, *Phys. Lett. A* **170**, 421 (1992).
 - [21] M. Joyce, J. Morand, and P. Viot, *Phys. Rev. E* **93**, 052129 (2016).
 - [22] N. I. Shakura and R. A. Sunyaev, *Astron. Astrophys.* **24**, 337 (1973).
 - [23] D. Lynden-Bell and J. E. Pringle, *Mon. Not. R. Astron. Soc.* **168**, 603 (1974).
 - [24] W. H. Press, B. P. Flannery, S. A. Teukolsky, and W. T. Vetterling, *Numerical Recipes in C: The Art of Scientific Computing* (Cambridge University Press, Cambridge, UK, 1988).