

REPRODUCTIVE PARAMETERS AND LONGEVITY OF
Gryon gallardoi (BRETHES) (HYMENOPTERA: SCELIONIDAE)
PARASITIZING *Spartocera dentiventris* (BERG)
(HEMIPTERA: COREIDAE) EGGS

CANTO-SILVA, C. R.¹, ROMANOWSKI, H. P.¹ and REDAELLI, L. R.^{1,2}

¹PPG Biologia Animal, UFRGS, Av. Bento Gonçalves, 9500, CEP 91501-970, Porto Alegre, RS, Brazil

²Departamento de Fitossanidade, UFRGS, Av. Bento Gonçalves, 7712, CEP 91540-000, Porto Alegre, RS, Brazil

Correspondence to: Celson Roberto Canto Silva, Universidade Estadual do Rio Grande do Sul – UERGS,
Unidade Cachoeira do Sul, Rua Sete de Setembro, 1040, Centro, Cachoeira do Sul, RS, Brazil,
e-mail: cantosilva@ig.com.br

Received September 30, 2003 – Accepted December 19, 2003 – Distributed February 28, 2006

(With 3 figures)

ABSTRACT

The fecundity, daily progeny and longevity of *Gryon gallardoi* (Brethes) (Hym.; Scelionidae) were determined under laboratory conditions, using *Spartocera dentiventris* (Berg) (Hem.; Coreidae) eggs as host. Nineteen *G. gallardoi* females and 34 males were reared at 25 ± 1 °C, with a 12 h photophase, fed on a 10% aqueous honey solution and provided with 25-30 *S. dentiventris* eggs daily. The average pre-oviposition period was 1.3 ± 0.35 days, although some females began laying from the day of emergence. On average, oviposition lasted for 10.1 ± 1.74 days, reaching a peak on the second day, with 67.5 ± 11.29 eggs laid. The post-oviposition period was short (2.4 ± 0.48 days). *G. gallardoi* females lived significantly longer than males: 13.7 ± 1.94 and 10.6 ± 1.78 days, respectively. The overall sex ratio was 0.79. The results reported here on the reproductive capability of the species suggest it may have a good potential as an agent for the control of *S. dentiventris*.

Keywords: insecta, tobacco, gray-tobacco-bug, parasitoid.

RESUMO

Parâmetros reprodutivos e longevidade de *Gryon gallardoi* (Brethes) (Hym.; Scelionidae) parasitando ovos de *Spartocera dentiventris* (Berg) (Hem.; Coreidae)

A fecundidade, a produção diária de prole e a longevidade de *Gryon gallardoi* (Brethes) (Hym.; Scelionidae) foram determinadas em condições de laboratório, utilizando-se ovos de *Spartocera dentiventris* (Berg) (Hem.; Coreidae) como hospedeiro. Dezenove fêmeas e 34 machos de *G. gallardoi* foram acompanhados ao longo de sua vida a 25 ± 1 °C, com fotofase de 12 h, recebendo solução aquosa de mel a 10% como alimento e grupos de 25 a 30 ovos de *S. dentiventris* para serem parasitados. O período médio de pré-oviposição foi $1,3 \pm 0,35$ dias, sendo que fêmeas foram capazes de ovipositar já no dia de sua emergência. Foi observado um período médio de oviposição de $10,1 \pm 1,74$ dias, com o pico de oviposição no segundo dia, sendo depositados ao longo dos dias uma média de $67,5 \pm 11,29$ ovos. O período de pós-oviposição foi curto ($2,4 \pm 0,48$ dias). Fêmeas de *G. gallardoi* foram significativamente mais longevas que os machos, vivendo, respectivamente, $13,7 \pm 1,94$ e $10,6 \pm 1,78$ dias. A razão sexual total observada foi de 0,79. Os resultados aqui registrados sobre a capacidade reprodutiva da espécie sugerem que esta apresenta um bom potencial como um agente para o controle de *S. dentiventris*.

Palavras-chave: insecta, fumo, percevejo-cinzento-do-fumo, parasitóide.

INTRODUCTION

The gray-tobacco-bug, *Spartocera dentiventris* (Berg) (Hem.: Coreidae), is a species associated with tobacco crops in Rio Grande do Sul, Brazil. The species cause losses resulting from the curling and withering of the attacked leaves (Costa, 1941; Schaefer & Panizzi, 2000). Parasitism is the main mortality factor acting on eggs of the bug and may reach rates close to 50% in the field (Santos *et al.*, 2001). Among the species of parasitoids responsible for such losses, *Gryon gallardoi* (Brethes) (Hym.: Scelionidae) is the most abundant, suggesting it may be a useful biological control agent.

Knowledge of the reproductive capacity of natural enemies is fundamental for the evaluation of a biological control agent's potential (Jervis & Kidd, 1996). Moreover, studies of the factors that influence this reproductive capacity are equally important. The relation between longevity and potential fecundity indirectly influences the actual fecundity of individuals, since the females may die before laying all their eggs. Parameters may be generated by studying the longevity and fecundity of species, which make it possible not only to compare the effectiveness of different natural enemies but also to build models that allow for more sophisticated analyses of this interaction.

The genus *Gryon* is composed of species associated mostly to coreid bugs, many of which are pests of important crops in the northern hemisphere, Africa and Asia (Nechols *et al.*, 1989; Asante *et al.*, 2000; Romeis *et al.*, 2000). In Brazil, *G. gallardoi* is cited as a natural enemy of coreids, attacking several cultivated and non-cultivated plants (Loiacono, 1980; Becker & Prato, 1982; De-Souza & Amaral-Filho, 1999). However, little is known about its role in the control of these species. Many biological and ecological studies of these species have been carried out recently in an attempt to understand the interaction between *S. dentiventris* and *G. gallardoi* on tobacco and to evaluate the use of this natural enemy as a control agent for the bug (Caldas *et al.*, 1999; Canto-Silva, 2003; Wiedemann *et al.*, 2003). Thus, the present work aimed to assess the reproductive biology and the longevity of *G. gallardoi* parasitizing *S. dentiventris* eggs.

MATERIAL AND METHODS

The present study was developed in the "Departamento de Fitossanidade, Faculdade de Agronomia, Universidade Federal do Rio Grande do Sul (UFRGS)", Porto Alegre (30° 01' S 51° 13' W), RS, Brazil, from November 2000 to February 2001. *G. gallardoi* was reared in the laboratory from *S. dentiventris* parasitized eggs collected from a tobacco crop in the field. The parasitoids were kept in 500 ml transparent plastic bottles containing *S. dentiventris* eggs, sealed with cotton wool and kept at 25 ± 1 °C, with a 12 h photophase. The parasitoids were fed daily *ad libitum* on a 10% aqueous honey solution. Non-parasitized *S. dentiventris* eggs were obtained from an experimental tobacco crop (type Virginia, var. k326), where *S. dentiventris* females were kept on the plants in voile fabric cages for the exclusion of parasitism.

In the laboratory, an experiment was carried out under the same breeding conditions. Newly emerged *G. gallardoi* couples (< 4 h old) were confined in 500 mL transparent plastic bottles containing food and about 25-30 freshly laid *S. dentiventris* eggs (< one day old). These eggs were kept on the part of the plant where they had been laid in order to minimize manipulation. The eggs were removed and substituted for a new group of non-parasitized eggs daily until the parasitoid female died. The eggs exposed to parasitism were then reared under the same conditions until their fate could be determined. When parasitoids emerged, they were sexed and counted. Eggs still remaining after a week elapsed since the last emergence were dissected to check their content.

For the analysis of fecundity and longevity, only females that had effectively produced offspring were considered. Individuals whose death was caused by accidental factors – such as injury due to manipulation or drowning in drops of the solution used as food – were excluded from the analysis. Individuals were kept in couples all the time so that, whenever a parasitoid died, it was replaced by other of the same sex. All handling of parasitoids was conducted with the use of CO₂. Based on these criteria, the results obtained correspond to the follow-up of 19 *G. gallardoi* females and 34 males.

The findings of male and female longevity were compared by the Mann-Whitney test (Sokal

& Rohlf, 1981) and the averages are presented together with their standard error.

RESULTS AND DISCUSSION

The average longevities of *G. gallardoi* males and females were 10.6 ± 1.78 and 13.7 ± 1.94 days, respectively (Table 1). The values recorded were highly variable for both sexes, with modes on 1 to 5 days for males and on 6 to 10 days for females (Fig. 1). Females lived significantly longer than males ($Z(u) = 2.15$; $p = 0.0314$). Maximum longevities recorded were 40 and 32 days for males

and females, respectively (Table 1). *G. gallardoi* longevity was lower than described for other species of the genus. Asante *et al.* (2000), studying the efficiency of *Gryon fulviventris* (Crawford), a parasitoid of the eggs of the coreid *Clavigralla tomentosicollis* Stal in Nigeria, found an average longevity of 34.29 ± 0.5 days (laboratory conditions: 27 ± 1 °C, 50-80% relative humidity, 50% aqueous honey solution diet, no hosts offered). Studies carried out in the central region of the USA with *Gryon pennsylvanicum* (Ashmead), an egg parasitoid of *Anasa tristis* DeGeer (Hemiptera: Coreidae), reported an average longevity of

TABLE 1

Reproductive parameters and longevity of *Gryon gallardoi* parasitizing *Spartocera dentiventris* eggs, reared at 25 ± 1 °C and a 12 h photophase, fed on a 10% aqueous honey solution.

Parameter	n	Mean \pm SE	Range
Male longevity (days)	34	10.6 ± 1.78	2 – 40
Female longevity (days)	19	13.7 ± 1.94	5 – 32
% of eggs parasitised/female	19	19.3 ± 1.81	2.86 – 35.92
Pre-oviposition period (days)	19	1.3 ± 0.35	0 – 6
Oviposition period (days)	19	10.1 ± 1.74	1 – 24
Post-oviposition period (days)	19	2.4 ± 0.48	0 – 6
Progeny/female	19	67.5 ± 11.29	8 – 173
Male progeny/female (%)	19	20.3 ± 1.49	12.5 – 38.18
Overall Sex ratio	1260	0.79	-




Fig. 1 — Percent survival of male (black blocks) and female (white blocks) *Gryon gallardoi* parasitizing *Spartocera dentiventris* eggs, reared in the laboratory at 25 ± 1 °C and a 12 h photophase, fed on a 10% aqueous honey solution and provided with hosts.

39.2 ± 12.4 days (laboratory conditions: 27 ± 1 °C, 75% relative humidity, 40% aqueous honey solution diet, hosts offered) (Nechols *et al.*, 1989). An even higher value was obtained by Romeis *et al.* (2000) for *Gryon clavigrallae* Mineo parasitizing *Clavigralla scutellaris* Spinola and *C. gibosa* (Westwood) (Hemiptera: Coreidae) in India: 55.0 ± 3.30 days for females and 28.8 ± 5.34 days for males (laboratory conditions: 25 ± 1 °C, 50 ± 10% relative humidity, non-diluted honey diet, no hosts offered). However, the lower longevity of *G. gallardoi* when compared with the above-listed species must be considered carefully since experimental conditions vary substantially between studies.

On average, approximately 19% of the eggs offered to the females during their lives were parasitized (Table 1). *G. gallardoi* females can start laying on the day they emerge, although only 32% of them did so. An average pre-oviposition period of 1.3 ± 0.35 days was observed (Table 1). On average, females laid for 10.1 ± 1.74 days about 67.5 ± 11.29 eggs (Table 1), distributed throughout most of the observed oviposition period. *G. gallardoi* fecundity is similar to that recorded for *G. pennsylvanicum* (71.8 ± 23.1 eggs; Vogt & Nechols, 1993) and higher than that found

for *G. clavigrallae* (56.4 ± 4.4 eggs; Romeis *et al.*, 2000). When compared with the fecundity of other Scelionidae, it lies on an intermediate level. For *Trissolcus basalis* (Wollaston), a classical *Nezara viridula* (L.) (Pentatomidae) biological control agent, Awan *et al.* (1990) reported 233.7 eggs in California, USA, while Corrêa-Ferreira & Zamataro (1989) obtained 250.4 ± 52.2 eggs in southern Brazil. On the other hand, studies on the scelionids *Telonomus podisi* Ashmead, *Trissolcus euschisti* (Ashmead) (Yergan, 1982) and *Telonomus cristatus* Johnson (Orr & Boethel, 1990) parasitizing *Podisus maculiventris* (Say) (Hem.; Pentatomidae) eggs showed reproductive potentials lower than the one obtained for *G. gallardoi* (39.6 ± 4.50, 63.7 ± 6.90 and 32.2 ± 1.80 eggs, respectively)

Parasitoid fecundity can reflect the abundance or ease with which hosts can be found (Price, 1974); hence, the above mentioned differences in fecundity may relate to the average size of the groups of eggs or egg clusters of the attacked hosts. In fact, both *P. maculiventris* and *S. dentiventris* lay groups with few eggs, *i.e.*, on average 18.7 (Orr *et al.*, 1986) and 23.9 eggs, respectively (Caldas *et al.*, 2000). In contrast, *N. viridula* parasitized by *T. basalis* lays clusters of approximately 100 eggs on average (Costa, 1996).




Fig. 2 — Influence of *Gryon gallardoi* age on the production of male (black circles) and female (white circles) progeny and female survival proportion (%) under laboratory-controlled conditions (25 ± 1 °C and 12 h photophase, fed on a 10% aqueous honey solution).

The production of female offspring per day peaked on the second day of life (7.1 ± 1.76 eggs, thereafter declining with the females' age (Fig. 2). This pattern, with an oviposition peak on the second or third day of life, is widespread not only among species of *Gryon* (Nechols *et al.*, 1989; Romeis *et al.*, 2000), but also among other genera of the Scelionidae (Powell & Shepard, 1982; Yergan, 1982; Corrêa-Ferreira & Zamataro, 1989). Male offspring were relatively constant along the parasitoid life, increasing slightly towards the end of the oviposition period of the long-living females (Fig. 2). The same finding was reported by Corrêa-Ferreira & Zamataro (1989) for *T. basalis* females over 4 days (Australian breed) or 7 days old (native breed).

An analysis of the pooled offspring indicated that all the females produced an excess of females: only $20.3 \pm 1.49\%$ of the offspring was male and the overall sexual ratio was 0.79 (Table 1).

Longer-living females oviposited more. Regression of the number of offspring versus female longevity showed a significantly positive relation ($y = 4.9464x - 0.1061$; $r^2 = 0.68$; $p = 0.0001$; Fig. 3), suggesting a high reproductive potential for the species, though strongly counteracted by the accomplished longevity. The maximum oviposition period recorded was 28 days (173 eggs). This long oviposition period associated with a short post-oviposition period (2.4 ± 0.48 days on average, Table 1), suggests a reproductive strategy typical of synovigenic parasitoids, *i.e.*, oocytes developing gradually along the lifespan. One may speculate that the long oviposition period of *G. gallardoi* is an adaptation to the similarly long *S. dentiventris*

oviposition period (approximately 48 days; Caldas *et al.*, 1999). Such a strategy would enable the parasitoid to have its reproduction synchronized with the seasonal availability of its host.

Comparisons of biological parameters obtained from laboratory studies are constrained by the variety of experimental conditions of the studies. The results reported here indicate that the magnitude of the species' fecundity is similar to that observed for other species used in biological control. The abundance of *G. gallardoi* in the system studied may be explained, at least in part, by such fecundity. This pattern indicates the good potential of the species as a biocontrol agent of *S. dentiventris* populations. The actual efficiency of *G. gallardoi* control depends on several factors that can act upon its fecundity in the field and on its effectiveness to locate and parasitize its host. Such aspects have been also investigated and the resulting analysis will be the subject of forthcoming papers.

Acknowledgments — The authors thank the Departamento de Fitossanidade, Faculdade de Agronomia - Universidade Federal do Rio Grande do Sul for the use of its facilities in the development of this study. Thanks are also due to Dra. Beatriz S. Corrêa-Ferreira for information and bibliography. *Contribution* n° 416 of the Department of Zoology, Post-graduate Program in Animal Biology, Instituto de Biociências, UFRGS.

REFERENCES

- ASANTE, S. K., JACKAI, E. N. & TAMO, M., 2000, Efficiency of *Gryon fulviventris* (Hymenoptera: Scelionidae) as an egg parasitoid of *Clavigralla tomentosicollis* (Hemiptera: Coreidae) in northern Nigeria. *Environ. Entomol.*, 29(4): 815-821.




Fig. 3 — Female longevity and progeny of *Gryon gallardoi* on *Spartocera dentiventris* eggs in the laboratory, at 25 ± 1 °C and a 12 h photophase, fed on a 10% aqueous honey solution.

- AWAN, M. S., WILSON, L. T. & HOFFMANN, M. P., 1990, Comparative biology of three geographic populations of *Trissolcus basalisi* (Hym.: Scelionidae). *Environ. Entomol.*, 19(2): 387-392.
- BECKER, M. & PRATO, M. D., 1982, Natalidade and natural mortality of *Spartocera lativentris* Stal, 1870 (Heteroptera: Coreidae) in the egg stage. *An. Soc. Entomol. Brasil*, 11: 269-281.
- CALDAS, B-H. C., REDAELLI, L. R. & DIEFENBACH, L. M. G., 1999, Parâmetros Reprodutivos de *Corecoris dentiventris* Berg (Hemiptera, Coreidae) em Cultura de Fumo (*Nicotiana tabacum*). *An. Soc. Entomol. Brasil*, 28(4): 595-600.
- CALDAS, B-H. C., REDAELLI, L. R. & L. M. G. DIEFENBACH, 2000, Biology of *Corecoris dentiventris* Berg, 1884 (Hemiptera: Coreidae) in tobacco culture (*Nicotiana tabacum*). *Rev. Bras. Biol.*, 60: 173-178.
- CANTO-SILVA, C. R., 2003, Parâmetros bioecológicos de *Gryon gallardoi* (Brethes) (Hymenoptera: Scelionidae) e modelagem da dinâmica espaço-temporal da sua interação com *Spartocera dentiventris* (Berg) (Hemiptera: Coreidae) através da simulação de múltiplos agentes. Tese de doutorado, Universidade Federal do Rio Grande do Sul, Programa de Pós-Graduação em Biologia Animal, Instituto de Biociências, Porto Alegre, 210p.
- CORRÊA-FERREIRA, B. S. & ZAMATARO, C. E. O., 1989, Capacidade reprodutiva e longevidade dos parasitoides de ovos *Trissolcus basalisi* (Wollaston) e *Trissolcus mitsukurii* Ashmead (Hymenoptera: Scelionidae). *Rev. Brasil. Biol.*, 49(2): 621-626.
- COSTA, J. A., 1996, *Cultura da soja*. Ed. Ivo Manica e José Antônio Costa, Brasil, 233p.
- COSTA, R. G., 1941, *Pragas de plantas cultivadas no Rio Grande do Sul*. Porto Alegre, Secretaria da Agricultura, 136p.
- DE-SOUZA, C. E. P. & AMARAL-FILHO, B. F., 1999, Ocorrência natural de parasitoides de *Leptoglossus zonatus* (Dallas) (Heteroptera: Coreidae). *An. Soc. Entomol. Brasil*, 28(4): 757-759.
- JERVIS, M. & KIDD, N., 1996, *Insect Natural Enemies: practical approaches to their study and evaluation*. Chapman & Hall, London, 491p.
- LOIACONO, M. S., 1980, Nota sobre tres Scelionidos parasitoides de hemipteros de la República argentina y Brasil (Hymenoptera-proctotrupoidea). *Rev. Soc. Ent. Argentina*, 39 (3-4): 173-178.
- NECHOLS, J. R., TRACY, J. L. & VOGT, E. A., 1989, Comparative ecological studies of indigenous egg parasitoids (Hymenoptera: Scelionidae, Encyrtidae) of the squash bug, *Anasa tristis* (Hemiptera: Coreidae). *J. Kansas Entomol. Soc.*, 62(2): 177-188.
- ORR, D. B. & BOETHEL, D. J., 1990, Reproductive Potential of *Telenomus cristatus* and *T. podisi* (Hymenoptera: Scelionidae), Two Egg Parasitoids of Pentatomids (Heteroptera). *Ann. Entomol. Soc. Am.*, 83(5): 902-905.
- ORR, D. B., RUSSIN, J. S., BOETHEL, D. J. & JONES, W. A., 1986, Stink bug (Hemiptera: Pentatomidae) egg parasitism in Louisiana soybean. *Environ. Entomol.*, 15: 1250-1254.
- POWELL, J. E. & SHEPARD, M., 1982, Biology of Australian and United States strains of *Trissolcus basalisi*, a parasitoid of green vegetable bug, *Nezara viridula*. *Aust. J. Ecol.*, 7: 181-186.
- PRICE, P. W., 1974, Strategies for egg production. *Evolution*, 28: 76-84.
- ROMEIS, J., SHANOWER, T. G. & MADHURI, K., 2000, Biology and field performance of *Gryon clavigrallae* (Hymenoptera: Scelionidae), an egg parasitoid of *Clavigralla* spp. (Hemiptera: Coreidae) in India. *Bull. Entomol. Res.*, 90: 253-263.
- SANTOS, R. S. S., REDAELLI, L. R. & DIEFENBACH, L. M. G., 2001, Ocorrência de Parasitismo em Ovos de *Spartocera dentiventris* (Berg) (Hemiptera: Coreidae) em Cultura de Fumo. *Neotrop. Entomol.*, 30(4): 731-733.
- SCHAEFER, C. W. & PANIZZI, A. R., 2000, *Heteroptera of Economic Importance*. CRC Press, Boca Raton, Florida, USA, 856p.
- SOKAL, R. R. & ROHLF, F. J., 1981, *Biometry*. 2nd ed., New York, Freeman, 859p.
- VOGT, E. A. & NECHOLS, J. R., 1993, The influence of host deprivation and host source on the reproductive biology and longevity of the squash bug egg parasitoid *Gryon pennsylvanicum* (Ashmead) (Hymenoptera: Scelionidae). *Biol. control*, 3: 148-154.
- WIEDEMANN, L. M., CANTO-SILVA, C. R., ROMANOWSKI, H. P. & REDAELLI, L. R., 2003, Oviposition behaviour of *Gryon gallardoi* (Hym.: Scelionidae) on eggs of *Spartocera dentiventris* (Hem.: Coreidae). *Braz. J. Biol.*, 63(1): 133-139.
- YERGAN, K. V., 1982, Reproductive capability and longevity of the parasitic wasps *Telenomus podisi* and *Trissolcus euschisti*. *Ann. Entomol. Soc. Am.*, 75(2): 181-183.