



Antimicrobial resistance and genetic relationships of enterococci from siblings and non-siblings *Heliconius erato phyllis* caterpillars

Rosana Huff¹, Rebeca Inhoque Pereira², Caroline Pissetti³, Aldo Mellender de Araújo⁴, Pedro Alves d'Azevedo², Jeverson Frazzon⁵ and Ana Paula Guedes Frazzon¹

¹Institute of Basic Health Sciences, Department of Microbiology, Immunology and Parasitology, Federal University of Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil

²Basic Health Sciences, Department of Microbiology, Health Sciences Federal University, Porto Alegre, Rio Grande do Sul, Brazil

³Department of Veterinary Preventive Medicine, Federal University of Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil

⁴Institute of Biosciences, Genetic Department, Federal University of Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil

⁵Food Science Institute, Federal University of Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil

ABSTRACT

Background. Studies evaluating bacteria in insects can provide information about host–microorganism–environment interactions. The gut microbial community has a profound effect on different physiological functions of insects. *Enterococcus* spp. are part of the gut community in humans and other animals, as well as in insects. The presence and antimicrobial resistance profile of enterococci are well studied in different animals; however, data for *Heliconius erato phyllis* (Lepidoptera: Nymphalidae) do not yet exist. Therefore, the aims of this study were to evaluate the distribution of enterococcal species, their antimicrobial resistance profile and virulence genes, and the genetic relationships between enterococci isolated from fecal samples from sibling and non-sibling *H. erato phyllis* caterpillars collected from different sites in South Brazil.

Methods. Three *H. erato phyllis* females were captured (two from a forest fragment and one from an urban area), and kept individually in open-air insectaries. Eggs were collected and caterpillars (siblings and non-siblings) were fed daily with *Passiflora suberosa* leaves. Fecal samples ($n = 12$) were collected from fifth-instar caterpillars, inoculated in selective medium, and 15 bacterial colonies were randomly selected from each sample. Enterococci were identified by PCR and MALDI-TOF, analyzed by disk diffusion antimicrobial susceptibility tests, and screened for resistance and virulence genes by PCR. The genetic relationships between the strains were determined using pulsed-field gel electrophoresis (PFGE).

Results. A total of 178 enterococci strains were identified: *E. casseliflavus* (74.15%; $n = 132$), *E. mundtii* (21.34%; $n = 38$), *E. faecalis* (1.12%; $n = 2$) and *Enterococcus* sp. (3.37%; $n = 6$). High rates of resistance to rifampicin (56%) and erythromycin (31%) were observed; 120 (67.41%) of the isolates showed resistance to at least one antibiotic and six (3.37%) were multidrug-resistant. None of the erythromycin-resistant strains was positive for the *erm*(B) and *msrC* genes. The virulence genes *esp*, *ace*, and *gelE*

Submitted 23 September 2019

Accepted 27 January 2020

Published 27 February 2020

Corresponding author

Ana Paula Guedes

Frazzon, ana.frazzon@ufrgs.br

Academic editor

Joseph Gillespie

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.8647

© Copyright
2020 Huff et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

were observed in 35%, 7%, and 1% of the strains, respectively. PFGE separated the enterococci into 22 patterns, four being composed of strains from sibling caterpillars. **Conclusion.** *Enterococcus casseliflavus* was the dominant species in fecal samples of fifth-instar caterpillars. Resistant enterococci strains may be related to environmental pollution or the resistome. The PFGE analysis showed genetic relationships between some strains, suggesting that the enterococci isolated from fecal samples of the sibling caterpillars might have come from common sources, e.g., via diet (herbivory) and/or vertical transmission (through the egg surface). Further studies will be conducted to better understand the role of *Enterococcus* in the microbial community of the gastrointestinal tract of these insects, and the mechanisms involved in acquisition and maintenance of enterococci.

Subjects Ecology, Microbiology, Molecular Biology

Keywords Lepidoptera, *Enterococcus casseliflavus*, Molecular typing, Antimicrobial profile, Enterococcal species

INTRODUCTION

Heliconius (Lepidoptera: Nymphalidae) represents a widespread genus of butterflies distributed throughout tropical and subtropical regions, from South America to the southern United States (Brown, 1981; Merrill et al., 2015). Adults feed on pollen as well as nectar (Gilbert, 1972; Merrill et al., 2015), and this food supply provides a better use of resources and a reduction in competition since the same flower can provide nutrients for both *Heliconius* and other butterflies that exclusively use nectar as food. This diet is rich in amino acids that allow adult females to oviposit on a daily basis during their lives (Gilbert, 1972). *Heliconius erato phyllis* is a subspecies commonly found in forests and urban environments from northeastern Brazil to northern Argentina. Adult females are monandrous, i.e., they mate only once, and lay their eggs individually in the apical meristem of a host plant, to minimize potential cannibalization of eggs by first-instar caterpillars (De Nardin, Da Silva & Araújo, 2016). Several species of passionflower vines, including *Passiflora suberosa*, *P. misera* and *P. capsularis*, are host plants for oviposition and feeding of caterpillars. These *Passiflora* species, commonly found in Southern Brazil where this study was conducted, have cyanogenic glycosides, which are assimilated by the caterpillars to make them unpalatable to potential predators. However, adults of *H. erato phyllis* are also able to synthesize cyanogenic glycosides and transfer them to their eggs (Hay-Roe & Nation, 2007).

It has been recognized for a long time that microorganisms play key roles in various physiological functions of animal hosts. The gut microbial community promotes an especially diverse range of benefits for insects, e.g., by improving nutrition via synthesis of vitamins and/or establishment of metabolic pathways, actively participating in degradation of xenobiotic compounds, and contributing to the defense against invading pathogens and immune system modulation (Douglas, 2015; Shao et al., 2017). The presence of microorganisms in the gastrointestinal (GI) tract of insects can be explained by environmental bacteria ingested with food and/or acquired by maternal transfer (Engel

& Moran, 2013). A growing number of studies have addressed the importance of the microbiota in the GI tract of insects (Engel & Moran, 2013; Chen et al., 2016; Douglas, 2018), and *Enterococcus* is one of the most frequent bacterial genera present in the gut microbiota at different life stages of Lepidoptera (Brinkmann, Martens & Tebbe, 2008; Chen et al., 2016; Snyman et al., 2016; van Shooten et al. 2018; Allonsius et al., 2019).

The *Enterococcus* genus is often found in the GI tract of humans and animals, as well as in the guts of insects (Shao et al., 2017). Hammer, McMillan & Fierer (2014) reported that *Enterococcus* was the most abundant genus found in immature stages and adults of *H. erato* from Panama. Furthermore, it has also been reported in insects of other orders, such as Coleoptera (Kim et al., 2017), Hymenoptera (Audisio et al., 2011), and Diptera (Ghosh et al., 2014). A characteristic of this genus is its intrinsic resistance to several antimicrobial agents and a great ability to transfer and acquire resistant genes (Hollenbeck & Rice, 2012). Despite the environmental resistome, the intense use of antimicrobials and anthropogenic activities, such as animal husbandry, agronomic practices, and wastewater treatment, play an important role in the emergence and spread of resistant-enterococci and/or antibiotic resistance genes in the environment, especially in soil, water, wastewater, and food (Gothwal & Shashidhar, 2014; Singer et al., 2016).

Antimicrobial resistance is one of the most serious public health problems, because of the spread of resistant bacteria leading to persistent infections, which are difficult to treat, and contamination of natural environments (Watkins & Bonomo, 2016; Ferri et al., 2017; Aslam et al., 2018). Insects have a wide distribution and can move freely between different environments; they may play an important role as reservoirs of drug-resistant strains and as their disseminators between animals and humans, especially when in contact with organic waste, livestock and their surrounding environment, and hospital facilities (Zurek & Ghosh, 2014; Mohammed et al., 2016; Schaumburg et al., 2016; Zhang et al., 2017; Onwugamba et al., 2018). In relation to insects carrying antibiotic-resistant bacterial strains, studies have identified flies (Ahmad et al., 2011; Usui et al., 2015; Mohammed et al., 2016; Schaumburg et al., 2016; Zhang et al., 2017; Onwugamba et al., 2018) and cockroaches (Ahmad et al., 2011; Pai, 2013; Moges et al., 2016) as hosts of extended-spectrum beta-lactamase- and carbapenemase-producing Enterobacteriaceae, vancomycin-resistant *E. faecium* (VRE), and methicillin-resistant *Staphylococcus aureus* (MRSA). Insects collected from food establishments and in association with stored products were also found to be colonized by antimicrobial-resistant bacteria (Macovei & Zurek, 2006; Channaiah et al., 2010; Mohammed et al., 2016). Despite their importance, few studies have addressed the concern of insects carrying resistant enterococci (Allen et al., 2009; Channaiah et al., 2010; Ahmad et al., 2011; Lowe & Romney, 2011).

As previously mentioned, the *Enterococcus* genus is often found in the GI tract in Lepidoptera (Hammer, McMillan & Fierer, 2014; Chen et al., 2016; Snyman et al., 2016; van Shooten et al. 2018; Allonsius et al., 2019), and it may play a fundamental role in the development and regulation of bacterial communities in these insects (Chen et al., 2016; Shao et al., 2017). The identification of enterococcal strains and their resistance profile in insects is an important aspect that must be addressed for host–microorganism–environment interactions. To our knowledge, there have been no studies to date evaluating

enterococci in *H. erato phyllis*. The aims of our research were to analyze enterococcal species distribution, their antimicrobial resistance profile and virulence genes, and the genetic relationships between enterococci isolated from fecal samples from sibling and non-sibling *H. erato phyllis* caterpillars collected from different sites in South Brazil.

MATERIALS & METHODS

Sample collection

The fecal samples used in the present study were collected from fifth instar caterpillars. The caterpillars were sourced from three different populations of *Heliconius erato phyllis* butterflies and consisted of sibling from the same female. The *H. erato phyllis* females (HE) were captured with entomological nets in Rio Grande do Sul, South Brazil. The first female (HEAB2) was collected in a forest fragment located in Águas Belas Agronomical Station (30°02'18.1"S; 51°01'23.0"W), the second female (HEV2) from a population in an intense urban area in Viamão (30°09'40.5"S; 50°55'01.5"W) and the third female (HES2) in a forest fragment located in São Francisco de Paula (29°26'34.1"S; 50°36'48.8"W).

Butterflies were kept individually in open-air insectaries with dimensions of 2.3 m × 3 m × 3 m (width, length, height) approximately. Insectaries had many plants for simulation of natural conditions, including *P. suberosa*, used by females for oviposition. The butterflies were fed daily with a mixture containing water, honey and pollen.

A total of 12 eggs were collected (five from HEAB2, five from HEV2 and two from HES2) with the assistance of a paintbrush. The eggs were transported to the laboratory, and caterpillars were grown individually in cylindrical plastic pots. Immatures were fed exclusively with *P. suberosa* leaves (Fig. 1). Fecal samples were collected from each caterpillar individually after 48 h of molting to the fifth instar ($n = 12$), with the aid of a disposable plastic spoon, stored in 1.5 mL microtubes and maintained at $-80\text{ }^{\circ}\text{C}$ until processing. The oviposition dates are shown in Table S1.

This study was carried out in accordance with the recommendations of Chico Mendes Institute for Biodiversity Conservation (ICMbio). The protocol was approved by Information Authorization System in Biodiversity (SISBIO) number 33404-1. This study has the Council for the Management of Genetic Patrimony - CGEN - under the Ministry of Environment number A720680.

Isolation and identification of Enterococci

Isolation and identification of enterococci were performed as previously described in Santestevan *et al.* (2015), with modifications. One milligram of fecal sample was transferred to 10 mL of saline 0.85% and incubated at $37\text{ }^{\circ}\text{C}$ for 24 h. One mL was inoculated in nine mL of Azide Dextrose Broth selective medium (Himedia, Mumbai, India) and incubated for 24 h at $37\text{ }^{\circ}\text{C}$. Aliquots (one mL) were placed in nine mL of saline 0.85%, and initial samples were further diluted 10-fold. From dilution 10^{-5} and 10^{-6} , 100 μL was inoculated in brain heart infusion (BHI) agar plates (Himedia, Mumbai, India) supplemented with 6.5% NaCl, incubated for 48 h at $37\text{ }^{\circ}\text{C}$.

Fifteen colonies were randomly selected from each fecal sample. Phenotypic criteria (size/volume, shape, color, gram staining, catalase production and bile aesculin reaction)




Figure 1 Illustration of *Heliconius erato phyllis* butterflies and caterpillars used in the study. (A) Female HEAB2 and offspring (6, 7, 10, 11, and 14); (B) female HEV2 and offspring (9, 18, 26, 27, and 29); (C) female HES2 and offspring (3 and 17). Illustration credit: Lucas de Oliveira Einsfeld and Gabriella Oliveira de Araújo.

Full-size  DOI: [10.7717/peerj.8647/fig-1](https://doi.org/10.7717/peerj.8647/fig-1)

were used to separate the enterococci group and the non-enterococcal strains. Selected pure colonies were stored at -20°C in a 10% (w/v) solution of skim milk (Difco, Sparks, MD, USA) and 10% (v/v) glycerol (Neon Comercial Ltda, São Paulo, SP, BR).

Genomic DNA was extracted by physicochemical method as previously described by *Depardieu, Perichon & Courvalin (2004)*. Polymerase chain reaction (PCR) assay was carried out using genus-specific primers targeting the *tuf* gene, which encodes elongation factor EF-Tu (*Ke et al., 1999*) (*Table 1*). Amplifications were performed in a total volume of 25 μL containing: 100 ng of template DNA, 1 X reaction buffer (Ludwig Biotechnology), 0.4 μM of each primer (Ludwig Biotechnology), 1.5 mM MgCl_2 , 200 μM of dNTP (Ludwig Biotechnology), 1 U *Taq* DNA polymerase (Ludwig Biotechnology), and MilliQ water. Amplification was carried out in a conventional thermocycler (Applied Biosystems 2720 Thermal Cycler) according to the following program: initial denaturation at 95°C for 3 min, followed by 35 cycles of 95°C for 30 s, 54°C for 30 s and 72°C for 1 min, and a final extension at 72°C for 7 min. The PCR products were visualized using electrophoresis on 1.5% (w/v) agarose gel, stained with SYBR[®] Safe DNA Gel, and visualized on a photocomputer. *E. faecalis* ATCC 29212 was used as positive control.

Characterization of enterococci species

Isolates were screened with the species-specific PCR assay for *Enterococcus faecalis*, *Enterococcus faecium*, *Enterococcus casseliflavus* and *Enterococcus mundtii* (*Cheng et al., 1997*; *Jackson, Fedorka-Cray & Barrett, 2004*; *Sedgley et al., 2005*). The primers and annealing temperature used are listed in *Table 1*. Amplifications were prepared as described to *tuf* gene. PCR amplification was performed in the conventional thermocycler (Applied Biosystems 2720 Thermal Cycler) according to the following program: 94°C for 5 min followed by 35 cycles of 94°C for 1 min, appropriate annealing temperature for each species for 1 min, extension at 72°C for 1 min, and a final extension at 72°C for 5 min.

Table 1 Primers used in the PCRs carried out in this study.

Primer	Nucleotide sequence (5'-3')	AT ^a (°C)	Amplicon (bp)	Reference
Genus				
tuf-F	TACTGACAAACCATTCATGATG	54	112	<i>Ke et al. (1999)</i>
tuf-R	AACTTCGTCACCAACGGGAAC			
<i>E. faecalis</i>				
E16s-F	CCGAGTGCTTGCACTCAATTGG	66	138	<i>Sedgley et al. (2005)</i>
E16s-R	CTCTTATGCCATGCGGCATAAAC			
<i>E. faecium</i>				
EM1A-F	TTGAGGGACACCAGATTGACG	62	658	<i>Cheng et al. (1997)</i>
EM1B-R	TATGACAGCGACTCCGATTCC			
<i>E. casseliflavus</i>				
CA1	TCCTGAATTAGGTGAAAAAAC	59	288	<i>Jackson, Fedorka-Cray & Barrett (2004)</i>
CA2	GCTAGTTTACCGTCTTTAACG			
<i>E. mundtii</i>				
MU1-F	CAGACATGGATGCTATTCCATCT	60	94	<i>Jackson, Fedorka-Cray & Barrett (2004)</i>
MU2-R	GCCATGATTTTCCAGAAGAATG			
<i>16s rRNA</i>				
8F	AGAGTTTGATCCTGGCTCAG	60	1,514	<i>Coenye et al. (1999)</i>
519F	CAGCAGCCGCGGTAATAC			
926R	CCGTCAATTCCCTTTGAGTT			
1522R	AAGGAGGTGATCCAGCCGCA			
Erythromycin				
erm(B)_F	GAAAAGGTA CTCAACCAAATA	52	639	<i>Sutcliffe, Tait-Kamradt & Wondrack (1996)</i>
erm(B)_R	AGTAACGGTACTTAAATTGTTTAC			
mnrC_3	AAGGAATCCTTCTCTCTCCG	52	343	<i>Werner, Hildebrandt & Witte (2001)</i>
mnrC_4	GTAAACAAAATCGTTCCCG			
Gelatinase				
gelE_TE9	ACCCCGTATCATTGGTTT	50	419	<i>Eaton & Gasson (2001)</i>
gelE_TE10	ACGCATTGCTTTTCCATC			
Cytolysin				
cylA_TE17	TGGATGATAGTGATAGGAAGT	59	517	<i>Eaton & Gasson (2001)</i>
cylA_TE18	TCTACAGTAAATCTTTTCGTCA			
Adhesion				
ace1_F	AAAGTAGAATTAGATCCACAC	59	320	<i>Mannu et al. (2003)</i>
ace2_R	TCTATCACATTCGGTTGCG			
Biofilm				
ESP46	TTACCAAGATGGTTCTGTAGGCAC	60	913	<i>Shankar et al. (1999)</i>
ESP47	CCAAGTATACTTAGCATCTTTTGG			
Aggregation				
agg_TE3	AAGAAAAAGAAGTAGACCAAC	60	1,553	<i>Eaton & Gasson (2001)</i>
agg_TE4	AAACGGCAAGACAAGTAAATA			

Notes.^aAT, annealing temperature.

The DNA fragment amplified by PCR was analyzed in 1.5% (w/v) agarose gels stained with SYBR[®] Safe DNA Gel, and visualized on a photocomputer.

Strains that were not identified by PCR reactions were submitted to matrix-assisted laser desorption and ionization time-of-flight technique (MALDI-TOF) applied to *Enterococcus* sp., according to [Sauget et al. \(2017\)](#).

Isolates classified as *Enterococcus* sp. were identified by Sanger sequence analysis. The PCR product of 16S rRNA gene, using the 8F and R1522 primers ([Coenye et al., 1999](#)) ([Table 1](#)), was purified with Illustra[™] GFX[™] PCR DNA and gel band purification kit (GE Healthcare, Buckinghamshire, UK). To perform Sanger sequencing two additional primers, 519F and 926R, were used ([Coenye et al., 1999](#)) ([Table 1](#)). Sequencing was performed with the ABI PRISM[®] BigDye[®] Primer Cycle Sequencing Ready Reaction Kit in an ABI PRISM[®] 3100 Genetic Analyzer (Applied Biosystems[®]), according to the manufacturer's protocol. The sequence obtained was compared to nucleotide sequences of reference enterococci strains deposited in GenBank.

Antimicrobial susceptibility testing

Antimicrobial susceptibilities were determined by Kirby-Bauer disk diffusion method recommended by the Clinical and Laboratory Standards Institute ([CLSI, 2016](#)). Eleven antibiotics used in clinical and veterinary medicine were tested: ampicillin 10 µg (AMP), vancomycin 30 µg (VAN), erythromycin 15 µg (ERY), tetracycline 30 µg (TET), ciprofloxacin 5 µg (CIP), norfloxacin 10 µg (NOR), nitrofurantoin 300 µg (NIT), rifampicin 5 µg (RIF), chloramphenicol 30 µg (CHL), gentamicin 120 µg (GEN) and streptomycin 300 µg (STR).

Intermediate and resistant strains were considered in a single category and classified as resistant. Strains showing resistance to three or more unrelated antibiotics were considered as multidrug-resistant (MDR) ([Schwarz et al., 2010](#)).

Detection of resistance and virulence genes

Erythromycin-resistant strains were tested by PCR for the presence of resistance encoding genes more commonly associated to clinical and environmental enterococci: *erm* (B), which encodes a ribosomal methylase that mediates MLSB resistance; and *msr* C, which encodes for a macrolide and streptogtamin B efflux pump. The presence of virulence associated genes *gelE* (gelatinase enzyme), *cylA* (activator of cytolysin), *ace* (accessory colonization factor), *esp* (associated to biofilm formation) and *agg* (aggregation substance) was determined by PCR in all enterococcal isolates. The amplifications were performed as described in [Prichula et al. \(2016\)](#). Amplifications were prepared as described to *tuf* gene. PCR amplification was performed in the conventional thermocycler (Applied Biosystems 2720 Thermal Cycler) according to the following program: 94 °C for 3 min followed by 35 cycles of 94 °C for 1 min, appropriate annealing temperature for each resistance or virulence gene for 1 min, extension at 72 °C for 1 min, followed by final extension at 72 °C for 5 min. The DNA fragment amplified by PCR was analyzed in 1.5% (w/v) agarose gels stained with SYBR[®] Safe DNA Gel, and visualized on a photocomputer. The sequences of the primers and annealing temperature are described in [Table 1](#).

Molecular typing of Enterococci by *Pulsed-field gel electrophoresis* (PFGE)

Eight six enterococci strains isolated from siblings and non-sibling caterpillars were selected for PFGE analysis according to the following criteria: maternal origin (females HEAB2, HEV2 or HES2), hatched larvae, enterococcal species and antimicrobial profile. Chromosomal DNA extraction and electrophoresis conditions were prepared according to [Murray et al. \(1990\)](#) and [Saeedi et al. \(2002\)](#). The restriction enzyme used was *Sma* I (Invitrogen®). The electrophoresis was carried out using a clamped homogeneous electric field (CHEF-DR1I device; Bio-Rad Laboratories, Richmond, Calif.), with ramped pulse times recommended by [Saeedi et al. \(2002\)](#) at 11 °C. Lambda Ladder PFG Marker (New England Biolabs) was used. The gels were stained with ethidium bromide (0.5 µg/mL for 20 min). The PFGE patterns were interpreted using the program GelCompar II v. 11 6.6, with 1.0% of tolerance, and the percentage of similarity was estimated using the Dice coefficient. The pulsotypes were clustered using the unweighted pair group whit arithmetic averages (UPGMA). A dendrogram was generated to examine the relatedness of PFGE patterns for selected isolates, and cutoff level of 80% applied to this dendrogram ([Tenover et al., 1995](#)).

Statistical Analysis

Simpson's index of diversity (D) was calculated to assess the differentiation of enterococci species among the caterpillars from different maternal origins ([Hunter & Gaston, 1988](#)).

RESULTS

Enterococci species present in fecal samples of *Heliconius erato phyllis* caterpillars

A total of 178 strains of *Enterococcus* were isolated from fecal samples from fifth-instar caterpillars ([Table 2](#)). *Enterococcus casseliflavus* was the most common species identified (74.15%; $n = 132$), followed by *E. mundtii* (21.34%; $n = 38$) and *E. faecalis* (1.12%; $n = 2$). Six strains (3.37%) could not be identified to species level.

Differences in the composition of enterococci were detected between the three groups of caterpillars, as shown in [Table 2](#). The Simpson's diversity index was different between the three populations, with higher diversity of enterococci species from fecal samples of caterpillars from HES2 ($1-D = 0.68$), followed by HEV2 ($1-D = 0.49$) and HEAB2 ($1-D = 0.27$).

Antimicrobial susceptibility

One hundred and twenty (67.41%) enterococci were resistant to at least one evaluated antimicrobial agent. The frequency of antibiotic-susceptible strains is shown in [Table 3](#). The rifampicin-resistance phenotype was the most commonly observed (56%; $n = 100$), followed by erythromycin (31%; $n = 55$). Eight strains (4%) were resistant to norfloxacin and five (3%) to ciprofloxacin. All investigated strains were susceptible to ampicillin, vancomycin, tetracycline, nitrofurantoin, chloramphenicol, gentamicin, and streptomycin.

Single- (SR), double- (DR), and multiple-drug resistance (MDR) were observed in 67% ($n = 80$), 28% ($n = 34$), and 5% ($n = 6$) of strains, respectively, of the 120 resistant strains.

Table 2 Distribution of *Enterococcus* sp. isolated from fecal samples of *Heliconius erato phyllis* caterpillars.

Species	Number (%) of enterococci strains isolated of caterpillars from ^a :			
	HEAB2	HEV2	HES2	Total (%)
<i>E. faecalis</i>	2 (2.04)	0	0	2 (1.12)
<i>E. casseliflavus</i>	83 (84.69)	42 (65.62)	7 (43.75)	132 (74.15)
<i>E. mundtii</i>	13 (13.26)	19 (29.68)	6 (37.50)	38 (21.34)
<i>Enterococcus</i> sp.	0	3 (4.68)	3 (18.75)	6 (3.37)
Total	98 (100)	64 (100)	16 (100)	178 (100)

Notes.

^aHEAB2, female from Águas Belas; HEV2, female from Viamão; HES2, female from São Francisco de Paula.

Table 3 Antibiotic resistance profiles in enterococci isolated from fecal samples of *Heliconius erato phyllis* caterpillars.

Female ^a	Species (n)	Number (%) of resistant strains ^b				Profiles ^c		
		ERY	CIP	NOR	RIF	SR	DR	MDR
HEAB2	<i>E. faecalis</i> (2)	2 (100)	0	0	2 (100)	0	2 (100)	0
	<i>E. casseliflavus</i> (83)	25 (30)	4 (5)	8 (10)	70 (84)	48 (58)	21 (25)	5 (6)
	<i>E. mundtii</i> (13)	0	0	0	0	0	0	0
HEV2	<i>E. casseliflavus</i> (42)	27 (64)	0	0	23 (55)	28 (67)	11 (26)	0
	<i>E. mundtii</i> (19)	0	0	0	0	0	0	0
	<i>Enterococcus</i> sp. (3)	0	0	0	3 (100)	3 (100)	0	0
HES2	<i>E. casseliflavus</i> (7)	0	0	0	0	0	0	0
	<i>E. mundtii</i> (6)	0	0	0	0	0	0	0
	<i>Enterococcus</i> sp. (3)	1 (33)	1 (33)	0	2 (67)	1 (33)	0	1 (33)
Total (178)		55 (31)	5 (3)	8 (4)	100 (56)	80 (45)	34 (19)	6 (3)

Notes.

^aHEAB2, female from Águas Belas; HEV2, female from Viamão; HES2, female from São Francisco de Paula.

^bAntibiotics: ERY, erythromycin; CIP, ciprofloxacin; NOR, norfloxacin; RIF, rifampicin.

^cProfiles: SR, single resistant; DR, double resistant; MDR, multidrug resistant.

Determinates of resistance and virulence

None of the 55 erythromycin-resistant strains was positive for *erm*(B) and *msr*C genes. The presence of virulence genes was evaluated in all strains, and the *esp* gene was detected in 35.39% ($n = 63$), *ace* in 6.74% ($n = 12$) and *gelE* in 1.12% ($n = 2$). No strain was positive for the *clyA* and *agg* genes.

Genetic relationships between enterococci isolated from sibling and non-sibling caterpillars in the fifth-instar

Of the 178 strains isolated, 86 (*E. casseliflavus*, $n = 58$; *E. mundtii*, $n = 23$; *E. faecalis*, $n = 2$; and *Enterococcus* sp., $n = 3$) were chosen for PFGE (Table S1). From the sibling caterpillars, numbered 6, 7, 10, 11 and 14, of the HEAB2 female were picked *E. casseliflavus* ($n = 32$), *E. mundtii* ($n = 8$) and *E. faecalis* ($n = 2$). All strains were isolated from caterpillars hatched closely in time and fed the same batch of *P. suberosa* leaves. From the offspring of the HEV2 female (sibling caterpillars 9, 18, 26, 27, and 29) *E. casseliflavus* ($n = 19$), *E. mundtii* ($n = 9$) and *Enterococcus* sp. ($n = 3$) were selected. These isolates were recovered from siblings that hatched at different times and were fed different batches of leaves. From the offspring

of the HES2 female (sibling caterpillars 3 and 17) *E. casseliflavus* ($n = 7$) and *E. mundtii* ($n = 6$) were selected.

The hierarchical relationship between enterococci selected from sibling and non-sibling caterpillars showed 22 patterns (15 patterns and 7 single strain-[singleton]; Fig. 2). Four patterns generated by PFGE indicated a genetic relationship between strains isolated from sibling caterpillars (P7, P8, P9, and P13); 11 were composed of strains isolated from the same caterpillars (P1, P2, P3, P4, P5, P6, P10, P11, P12, P14, and P15). No genetic relationships were observed for strains isolated from non-siblings.

The band patterns for *E. casseliflavus* ($n = 32$) isolates from sibling caterpillars (6, 7, 10, 11 and 14) of the HEAB2 female showed six PFGE patterns (P5, P7, P8, P9, P11 and P12) and three singletons. Three PFGE patterns (P7, P8, and P9) included 18 of the 32 strains that were isolated from sibling caterpillars 6, 7, 10, and/or 11, with low levels of genetic variability. P5 and P11 each contained two isolates, and P12 with eight isolates showed genetic variation; the remaining three *E. casseliflavus* isolates were singletons and represented unique PFGE patterns. All *E. mundtii* isolates from caterpillar 14 were genetically closely related and were clustered into one pattern (P10), as were the two *E. faecalis* (P3) isolates from caterpillar 6. These results demonstrate that strains may originate from a single lineage.

Seven different PFGE patterns (P4, P6, P13, P15, and three singletons) were obtained from sibling caterpillars 9, 18, 26, 27, and 29 of the HEV2 female. The analysis of the fragment profiles of the *E. mundtii* ($n = 9$) and *Enterococcus* sp. ($n = 1$) strains isolated from caterpillars 18, 26, 27, and 29 demonstrated a genetic relationship between them (P13). The 19 *E. casseliflavus* strains showed four PFGE patterns: P4, P6, and P15, each containing six, three, and nine isolates, respectively, and one singleton. These distinct patterns are suggestive of genetic events in these strains.

The seven *E. casseliflavus* strains from caterpillars 3 and 17 (offspring of the HES2 female) showed two distinct patterns (P1 and P2) with 100% of genetic similarity between them. P1 contained four isolates and P2 had three isolates. Of the six *E. mundtii* isolates from caterpillar 3, five showed 100% similarity and were clustered in the P14 pattern, suggesting that these strains may be progeny from a single lineage. One strain had distinct and unrelated PFGE by the criteria of Tenover et al. (1995). In addition, most of the patterns were shared by isolates with the same antimicrobial profile.

DISCUSSION

An increasing number of studies have aimed to investigate the microbial communities in the GI tract of insects. In the Lepidoptera, a high abundance of *Enterococcus* has been found, both in immatures and adults, raising questions about the role of these bacteria in invertebrates and their importance in maintaining the health of individuals (Hammer, McMillan & Fierer, 2014; Holt et al., 2015; Chen et al., 2016; Snyman et al., 2016; Shao et al., 2017; Van Schooten et al., 2018; Allonsius et al., 2019). To our knowledge, only Hammer, McMillan & Fierer (2014) and, more recently, Van Schooten et al. (2018) have addressed the microbial communities in the *Heliconius* GI tract, and have reported that the genus *Enterococcus* is dominant in microbial samples from different *Heliconius* species.




Figure 2 Dendrogram of enterococci isolated from fecal samples of *Heliconius erato phyllis* caterpillars. HEAB2, *H. erato phyllis* from Águas Belas; HEV2, *H. erato phyllis* from Viamao; HES2, *H. erato phyllis* from São Francisco de Paula. Antibiotics: RIF, rifampicin; ERY, erythromycin; NOR, norfloxacin; CIP, ciprofloxacin; S, susceptible; R, resistant.

Full-size DOI: 10.7717/peerj.8647/fig-2

Enterococcus is associated with the environment and a wide range of organisms, including plants (Müller *et al.*, 2001; Byappanahalli *et al.*, 2012; Sánchez Valenzuela *et al.*, 2012). In the present study, the most abundant *Enterococcus* species in the feces of *H. erato phyllis* caterpillars was *E. casseliflavus*. The diet of insects constitutes an additional source of microorganisms in the GI tract. Enterococci in the GI tract of caterpillars can play an important role in the protection against other pathogens, since these bacteria are able to produce lactic acid (causing a decrease in pH) and enterocins (peptides with antimicrobial activity). Since *E. casseliflavus* is frequently a part of microbial communities on plants (Byappanahalli *et al.*, 2012; Micallef *et al.*, 2013; Ong *et al.*, 2014), the predominance of this species in fecal samples of *H. erato phyllis* caterpillars could be explained by the plant diet of juveniles, resulting in the dominance of this species. As noted by Chen *et al.* (2016), enterococci stably colonize the larval gut of *Spodoptera littoralis* (Lepidoptera: Noctuidae), and may act beneficially in caterpillars by promoting nutrient supplementation (metabolism of terpenoids and polyketides) and host defense (production of antimicrobials by *E. mundtii*). However, further studies are needed to understand the role of these bacteria as members of the GI tract in *H. erato phyllis* caterpillars.

Resistant enterococcal strains isolated from fecal samples of caterpillars are a matter of concern, since these larvae had not been exposed to antibiotics. Studies that highlight the resistance profile of enterococci isolated from insects in natural environments are scarce. Channaiah *et al.* (2010) described enterococci isolated from insect pests of stored products; they were resistant to tetracycline, streptomycin, erythromycin, kanamycin, ciprofloxacin, ampicillin, and chloramphenicol; this suggests that these animals can be potential vectors in disseminating antibiotic-resistant strains. Ahmad *et al.* (2011) reported MDR enterococci isolated from house flies and cockroaches in a confined swine production environment, and have suggested that these insects may be potential vectors and/or reservoirs of resistant enterococci. Despite the occurrence and spread of resistant strains intensifying due to the use of antimicrobials, the isolation of resistant enterococci in the present study could be related to the resistance that occurs naturally in the environment (environmental resistome) and/or to anthropogenic activities (contamination of the environment) (Martínez, 2008; Allen *et al.*, 2010).

The most widespread mechanism of resistance to macrolides in enterococci is mediated by the *erm* and *msr* C genes (Aarestrup *et al.*, 2000; Santestevan *et al.*, 2015; Prichula *et al.*, 2016). Nevertheless, none of these genes was detected in the present study. It is possible that these strains harbored other erythromycin-resistance genes, such as *erm* D, E, and F, and other efflux pump genes such as *msr* A. A low percentage of virulence genes was detected in enterococci of *H. erato phyllis* caterpillars. Although these genes are related to pathogenicity of clinical enterococcal strains, their presence in strains in fecal samples from caterpillars may be associated with the maintenance of cells of the GI tract, and consequently with microorganism and host interactions.

From the analysis of the PFGE fingerprint, *E. casseliflavus*, *E. faecalis*, *E. mundtii*, and *Enterococcus* sp. isolated from fecal samples of sibling and non-sibling caterpillars demonstrated unrelated or related patterns based on maternal origin. The unrelated patterns found in P1, P2, P3, P4, P5, P6, P10, P11, P12, P14, and P15 demonstrated genetic

diversity among these strains. The genetic variation in these strains may be associated with genetic events, such as mobile elements or mutation, a common characteristic of enterococci (Lebreton, Willems & Gilmore, 2014). The related patterns observed among strains isolated from sibling caterpillars (P7, P8, P9, and P13) may be associated with a common source, e.g., diet (herbivory) and/or vertical transmission (through the egg surface). Since the fecal samples used in the present study were collected from fifth-instar caterpillars, the last stage before the pupa, the results present here may suggest vertical transmission of enterococci that are being replaced from the diet. Plants are a food source for bacteria present in the GI tract of insects; these bacteria improve the quality of diets poor in nutrients and take part in development and maturation of the immune system to protect the host against pathogenic microorganisms (Dillon & Dillon, 2004; Engel & Moran, 2013). Therefore, it is likely that herbivory provides an abundant supply of enterococci throughout the larval stage of *H. erato phyllis*. Considering that *Passiflora* leaves are the only food of the caterpillars, those leaves could be the source of *Enterococcus* sp. in their GI tracts.

Besides the diet, vertical transmission from the female to her offspring can also be a source of bacteria. Some studies have described the mechanism for vertical transmission of bacteria in different species of Lepidoptera (Brinkmann, Martens & Tebbe, 2008; Chen et al., 2016; Teh et al., 2016; Shao et al., 2017). The caterpillar hatches by chewing a hole in the chorion and emerging through it from the egg. Brinkmann, Martens & Tebbe (2008) reported that the *Enterococcus* spp. present in the gut of the larvae of *Manduca sexta* (Lepidoptera: Sphingidae) were acquired via ingestion of their eggshell, demonstrating maternal transmission of microorganisms. Chen et al. (2016) analyzed the composition and activity of microbiota in the moth *Spodoptera littoralis*, which is an agricultural pest; they found that enterococci associated with adult females were also in the egg mass, and further colonized the larval gut of individuals, suggesting vertical transmission of these bacteria. Teh et al. (2016) showed the route of transmission of *E. mundtii* in *S. littoralis* when administered *in vivo*; the authors reported the presence of *E. mundtii* at all life stages of this insect. In addition, they established the presence of these bacteria in oocytes and the muscle tissue in the first-instar larvae of the second-generation offspring, highlighting again the vertical transmission of enterococci in this lepidopteran. The enterococci isolated in the present study may be linked to herbivory; however, although our analysis does not include adult females and the bacterial communities of *Passiflora* leaves, we do not rule out the possibility that enterococci could also be transmitted from the female to the offspring through the surface of the egg, as previously demonstrated for other species of Lepidoptera.

CONCLUSIONS

Enterococcus casseliflavus was the dominant enterococcal species isolated in fecal samples of the fifth-instar caterpillars of *H. erato phyllis*. Resistant strains present in the caterpillars studied may be related to the environmental resistome and/or anthropogenic activity. In addition, the presence of MDR enterococci may be an indication of contamination

of the environmental by antibiotics. The results obtained by PFGE analysis suggest that the enterococci isolated from fecal samples of sibling caterpillars might have come from common sources, e.g., the diet (herbivory) and/or vertical transmission (through the egg surface). Further studies will be conducted to better understand the role of *Enterococcus* in the GI tract microbial community of *H. erato phyllis* butterflies, and the mechanisms involved in acquisition and maintenance of these bacteria. In addition, the data obtained can be used in future comparative analyses of the microbiota present in adult *H. erato phyllis* females and their offspring, to confirm the occurrence of vertical transmission of *Enterococcus* sp. in this model organism.

ACKNOWLEDGEMENTS

We thank Lucas de Oliveira Einsfeld and Gabriella Oliveira de Araújo for the drawing.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico do Brasil, CNPq - #302574/2017-4, #401714/2016-0 and #303603/2015-1 and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Conselho Nacional de Desenvolvimento Científico e Tecnológico do Brasil.
CNPq: #302574/2017-4, #401714/2016-0 and #303603/2015-1.
Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Rosana Huff conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Rebeca Inhoque Pereira conceived and designed the experiments, performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Caroline Pissetti performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Aldo Mellender de Araújo conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Pedro Alves d'Azevedo and Jeverson Frazzon analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

- Ana Paula Guedes Frazzon conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

This study was carried out in accordance with the recommendations of Chico Mendes Institute for Biodiversity Conservation (ICMbio).

The protocol was approved by Information Authorization System in Biodiversity (SISBIO) number 33404-1. This study has the approval of the Council for the Management of Genetic Patrimony - CGEN - under the Ministry of Environment number A720680.

Data Availability

The following information was supplied regarding data availability:

The raw data and electrophoresis gels are available in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8647#supplemental-information>.

REFERENCES

- Aarestrup FM, Agero Y, Gerner-Smidt P, Madsen M, Jensen LB. 2000. Comparison of antimicrobial resistance phenotypes and resistance genes in *Enterococcus faecalis* and *Enterococcus faecium* from humans in the community, broilers, and pigs in Denmark. *Diagn. Microbiol Infect Dis* 37(2):127–137 DOI 10.1016/S0732-8893(00)00130-9.
- Ahmad A, Ghosh A, Schal C, Zurek L. 2011. Insects in confined swine operations carry a large antibiotic resistant and potentially virulent enterococcal community. *BMC Microbiology* 11(1):23 DOI 10.1186/1471-2180-11-23.
- Allen HK, Cloud-Hansen KA, Wolinski JM, Guan C, Greene S, Lu S, Boeyink M, Broderick NA, Raffa KF, Handelsman J. 2009. Resident microbiota of the gypsy moth midgut harbors antibiotic resistance determinants. *DNA and Cell Biology* 28(3):109–117 DOI 10.1089/dna.2008.0812.
- Allen HK, Donato J, Wang HH, Cloud-Hansen KA, Davies J, Handelsman J. 2010. Call of the wild: antibiotic resistance genes in natural environments. *Nature Reviews Microbiology* 8(4):251–259 DOI 10.1038/nrmicro2312.
- Allonsius CN, Van Beeck W, De Boeck I, Wittouck S, Lebeer S. 2019. The microbiome of the invertebrate model host *Galleria mellonella* is dominated by *Enterococcus*. *BMC Animal Microbiome* 1:7 DOI 10.1186/s42523-019-0010-6.
- Aslam B, Wang W, Arshad MI, Khurshid M, Muzammil S, Rasool MH, Nisar NA, Alvi RF, Aslam MA, Qamar MU, Salamat MKF, Baloch Z. 2018. Antibiotic resistance: a rundown of a global crisis. *Infection and Drug Resistance* 11:1645–1658 DOI 10.2147/IDR.S173867.

- Audisio MC, Torres MJ, Sabaté DC, Ibareguren C, Apella MC. 2011.** Properties of different lactic acid bacteria isolated from *Apis mellifera* L. bee-gut. *Microbiological Research* **166**(1):1–13 DOI [10.1016/j.micres.2010.01.003](https://doi.org/10.1016/j.micres.2010.01.003).
- Brinkmann N, Martens R, Tebbe CC. 2008.** Origin and diversity of metabolically active gut bacteria from laboratory-bred larvae of *Manduca sexta* (Sphingidae, Lepidoptera, Insecta). *Applied and Environmental Microbiology* **74**(23):7189–7196 DOI [10.1128/AEM.01464-08](https://doi.org/10.1128/AEM.01464-08).
- Brown KS. 1981.** The biology of *Heliconius* and related genera. *Annual Review of Entomology* **26**:427–456 DOI [10.1146/annurev.en.26.010181.002235](https://doi.org/10.1146/annurev.en.26.010181.002235).
- Byappanahalli MN, Nevers MB, Korajkic A, Staley ZR, Harwood VJ. 2012.** Enterococci in the environment. *Microbiology and Molecular Biology Reviews* **76**(4):685–706 DOI [10.1128/MMBR.00023-12](https://doi.org/10.1128/MMBR.00023-12).
- Channaiah LH, Subramanyam B, McKinney LJ, Zurek L. 2010.** Stored-product insects carry antibiotic-resistant and potentially virulent enterococci. *FEMS Microbiology Ecology* **74**(2):464–471 DOI [10.1111/j.1574-6941.2010.00949.x](https://doi.org/10.1111/j.1574-6941.2010.00949.x).
- Chen B, Teh B, Sun C, Hu S, Lu X, Boland W, Shao Y. 2016.** Biodiversity and activity of the gut microbiota across the life history of the insect herbivore *Spodoptera littoralis*. *Scientific Reports* **6**:29505 DOI [10.1038/srep29505](https://doi.org/10.1038/srep29505).
- Cheng S, McCleskey FK, Gress MJ, Petroziello JM, Liu R, Namdari H, Beninga K, Salmen A, Delvecchio VG. 1997.** A PCR assay for identification of *Enterococcus faecium*. *Journal of Clinical Microbiology* **35**(5):1248–1250 DOI [10.1128/JCM.35.5.1248-1250.1997](https://doi.org/10.1128/JCM.35.5.1248-1250.1997).
- Clinical and Laboratory Standards Institute (CLSI). 2016.** *Performance standards for antimicrobial susceptibility testing—twelve-six edition 100S*. Wayne: CLSI.
- Coenye T, Falsen E, Vancanneyt M, Hoste B, Govan JRW, Kersters K, Vandamme P. 1999.** Classification of *Alcaligenes faecalis*-like isolates from the environment and human clinical samples as *Ralstonia gilardii* sp. nov. *International Journal of Systematic Bacteriology* **49**(2):405–413 DOI [10.1099/00207713-49-2-405](https://doi.org/10.1099/00207713-49-2-405).
- De Nardin J, Da Silva L, Araújo AM. 2016.** Kin recognition in a butterfly: inferences about its heritability. *Ethol Ecol Evol* **29**(3):1–11.
- Depardieu F, Perichon B, Courvalin P. 2004.** Detection of the *van* Alphabet and identification of enterococci and staphylococci at the species level by multiplex PCR. *Journal of Clinical Microbiology* **42**(12):5857–5860 DOI [10.1128/JCM.42.12.5857-5860.2004](https://doi.org/10.1128/JCM.42.12.5857-5860.2004).
- Dillon RJ, Dillon VM. 2004.** The gut bacteria of insects: nonpathogenic interactions. *Annual Review of Entomology* **49**:71–92 DOI [10.1146/annurev.ento.49.061802.123416](https://doi.org/10.1146/annurev.ento.49.061802.123416).
- Douglas AE. 2015.** Multiorganismal insects: diversity and function of resident microorganisms. *Annual Review of Entomology* **60**:17–34 DOI [10.1146/annurev-ento-010814-020822](https://doi.org/10.1146/annurev-ento-010814-020822).
- Douglas AE. 2018.** Gut microbes alter fly walking activity. *Nature* **563**:331–333 DOI [10.1038/d41586-018-07080-y](https://doi.org/10.1038/d41586-018-07080-y).
- Eaton TJ, Gasson MJ. 2001.** Molecular screening of *Enterococcus* virulence determinants and potential for genetic exchange between food and medical isolates. *Applied and Environmental Microbiology* **67**(4):1628–1635 DOI [10.1128/AEM.67.4.1628-1635.2001](https://doi.org/10.1128/AEM.67.4.1628-1635.2001).

- Engel P, Moran NA. 2013.** The gut microbiota of insects—diversity in structure and function. *FEMS Microbiology Reviews* **37**(5):699–735 DOI [10.1111/1574-6976.12025](https://doi.org/10.1111/1574-6976.12025).
- Ferri M, Ranucci E, Romagnoli P, Giaccone V. 2017.** Antimicrobial resistance: a global emerging threat to public health systems. *Critical Reviews in Food Science and Nutrition* **57**(13):2857–2876 DOI [10.1080/10408398.2015.1077192](https://doi.org/10.1080/10408398.2015.1077192).
- Ghosh A, Akhtar M, Holderman C, Zurek L. 2014.** Significance and survival of enterococci during the house fly development. *Journal of Medical Entomology* **51**(1):63–67 DOI [10.1603/ME13161](https://doi.org/10.1603/ME13161).
- Gilbert LE. 1972.** Pollen feeding and reproductive biology of *Heliconius* butterflies. *Proceedings of the National Academy of Sciences of the United States of America* **69**(6):1403–1407 DOI [10.1073/pnas.69.6.1403](https://doi.org/10.1073/pnas.69.6.1403).
- Gothwal R, Shashidhar T. 2014.** Antibiotic pollution in the environment: a review. *Clean* **43**:479–489.
- Hammer TJ, McMillan WO, Fierer N. 2014.** Metamorphosis of a butterfly-associated bacterial community. *PLOS ONE* **9**:e86995 DOI [10.1371/journal.pone.0086995](https://doi.org/10.1371/journal.pone.0086995).
- Hay-Roe MM, Nation J. 2007.** Spectrum of cyanide toxicity and allocation in *Heliconius erato* and *Passiflora* host plants. *Journal of Chemical Ecology* **33**(2):319–329 DOI [10.1007/s10886-006-9234-5](https://doi.org/10.1007/s10886-006-9234-5).
- Hollenbeck BL, Rice LB. 2012.** Intrinsic and acquired resistance mechanisms in enterococcus. *Virulence* **3**(5):421–433 DOI [10.4161/viru.21282](https://doi.org/10.4161/viru.21282).
- Holt JF, Kiedrowski MR, Frank KL, Du J, Guan C, Broderick NA, Dunny GM, Handelsman J. 2015.** *Enterococcus faecalis* 6-Phosphogluconolactonase is required for both commensal and pathogenic interactions with *Manduca sexta*. *Infection and Immunity* **83**(1):396–404 DOI [10.1128/IAI.02442-14](https://doi.org/10.1128/IAI.02442-14).
- Hunter PR, Gaston MA. 1988.** Numerical index of the discriminatory ability of typing systems: an application of Simpson's index of diversity. *Journal of Clinical Microbiology* **26**(11):2465–2466 DOI [10.1128/JCM.26.11.2465-2466.1988](https://doi.org/10.1128/JCM.26.11.2465-2466.1988).
- Jackson CR, Fedorka-Cray PJ, Barrett JB. 2004.** Use of a genus- and species-specific multiplex PCR for identification of Enterococci. *Journal of Clinical Microbiology* **42**:3558–3565 DOI [10.1128/JCM.42.8.3558-3565.2004](https://doi.org/10.1128/JCM.42.8.3558-3565.2004).
- Ke D, Picard FJ, Martineau F, Menard C, Roy PH, Ouellette M, Bergeron MG. 1999.** Development of a PCR assay for rapid detection of enterococci. *Journal of Clinical Microbiology* **37**(11):3497–3503 DOI [10.1128/JCM.37.11.3497-3503.1999](https://doi.org/10.1128/JCM.37.11.3497-3503.1999).
- Kim JM, Choi MY, Kim JW, Lee SA, Ahn JH, Song J, Kim SH, Weon HY. 2017.** Effects of diet type, developmental stage, and gut compartment in the gut bacterial communities of two Cerambycidae species (Coleoptera). *Journal of Microbiology* **55**(1):21–30 DOI [10.1007/s12275-017-6561-x](https://doi.org/10.1007/s12275-017-6561-x).
- Lebreton F, Willems RJL, Gilmore MS. 2014.** Enterococcus diversity, origins in nature, and gut colonization. In: Gilmore MS, Clewell DB, Ike Y, Shankar N, eds. *Enterococci: from commensals to leading causes of drug resistant infection*. Massachusetts: Eye and Ear Infirmary, 1–82.
- Lowe CF, Romney MG. 2011.** Bedbugs as vectors for drug-resistant bacteria. *Emerging Infectious Diseases* **17**(6):1132–1134 DOI [10.3201/eid1706101978](https://doi.org/10.3201/eid1706101978).

- Macovei L, Zurek L. 2006.** Ecology of antibiotic resistance genes: characterization of enterococci from houseflies collected in food settings. *Applied and Environmental Microbiology* **72(6)**:4028–4035 DOI [10.1128/AEM.00034-06](https://doi.org/10.1128/AEM.00034-06).
- Mannu L, Paba A, Daga E, Comunian R, Zanetti S, Duprè I, Sechi LA. 2003.** Comparison of the incidence of virulence determinants and antibiotic resistance between *Enterococcus faecium* strains of dairy, animal and clinical origin. *International Journal of Food Microbiology* **88**:291–304 DOI [10.1016/S0168-1605\(03\)00191-0](https://doi.org/10.1016/S0168-1605(03)00191-0).
- Martínez JL. 2008.** Antibiotics and antibiotic resistance genes in natural environments. *Science* **321(5887)**:365–367 DOI [10.1126/science.1159483](https://doi.org/10.1126/science.1159483).
- Merril RM, Dasmahapatra KK, Davey JW, Dell’Aglío DD, Hanly JJ, Huber B, Jiggins CD, Joron M, Kozak KM, Llauren V, Martin SH, Montgomery SH, Morris J, Nadeau NJ, Pinharanda AL, Rosser N, Thompson MJ, Vanjari S, Wallbank RWR, Yu Q. 2015.** The diversification of *Heliconius* butterflies: what have we learned in 150 years? *Journal of Evolutionary Biology* **28(8)**:1417–1438 DOI [10.1111/jeb.12672](https://doi.org/10.1111/jeb.12672).
- Micallef SA, Goldstein RER, George A, Ewing L, Tall BD, Boyer MS, Joseph SW, Sapkota AR. 2013.** Diversity, distribution and antibiotic resistance of *Enterococcus* spp. recovered from tomatoes, leaves, water and soil on U.S. Mid-Atlantic farms. *Food Microbiology* **36(2)**:465–474 DOI [10.1016/j.fm.2013.04.016](https://doi.org/10.1016/j.fm.2013.04.016).
- Moges F, Eshetie S, Endris M, Huruy K, Muluye D, Feleke T, G/Silassie F, Ayalew G, Nagappan R. 2016.** Cockroaches as a source of high bacterial pathogens with multidrug resistant strains in Gondar town, Ethiopia. *BioMed Research International* **2016**:2825056 DOI [10.1155/2016/2825056](https://doi.org/10.1155/2016/2825056).
- Mohammed AN, Abdel-Latef GK, Abdel-Azeem NM, El-Dakhly KM. 2016.** Ecological study on antimicrobial-resistant zoonotic bacteria transmitted by flies in cattle farms. *Parasitology Research* **115(10)**:3889–3896 DOI [10.1007/s00436-016-5154-7](https://doi.org/10.1007/s00436-016-5154-7).
- Müller T, Ulrich A, Ott EM, Müller M. 2001.** Identification of plant-associated enterococci. *Journal of Applied Microbiology* **91(2)**:268–278 DOI [10.1046/j.1365-2672.2001.01373.x](https://doi.org/10.1046/j.1365-2672.2001.01373.x).
- Murray BE, Singh KV, Heath JD, Sharma BR, Weinstock GM. 1990.** Comparison of genomic DNAs of different enterococcal isolates using restriction endonucleases with infrequent recognition sites. *Journal of Clinical Microbiology* **28(9)**:2059–2063 DOI [10.1128/JCM.28.9.2059-2063.1990](https://doi.org/10.1128/JCM.28.9.2059-2063.1990).
- Ong YY, Tan WS, Mohamad R, Sieo CC, Tey BT. 2014.** Biochemical and molecular identification of *Enterococcus* spp. from red pitaya. *Process Biochemistry* **49(4)**:563–568 DOI [10.1016/j.procbio.2014.01.019](https://doi.org/10.1016/j.procbio.2014.01.019).
- Onwugamba FC, Fitzgerald JR, Rochon K, Guardabassi L, Alabi A, Kühne S, Grobusche MP, Schaumburg F. 2018.** The role of ‘Filth Flies’ in the spread of antimicrobial resistance. *Travel Medicine and Infectious Disease* **22**:8–17 DOI [10.1016/j.tmaid.2018.02.007](https://doi.org/10.1016/j.tmaid.2018.02.007).
- Pai H. 2013.** Multidrug resistant bacteria isolated from cockroaches in long-term care facilities and nursing homes. *Acta Tropica* **125(1)**:18–22 DOI [10.1016/j.actatropica.2012.08.016](https://doi.org/10.1016/j.actatropica.2012.08.016).

- Prichula J, Pereira RI, Wachholz GR, Cardoso LA, Tolfo NC, Santestevan NA, Medeiros AW, Tavares M, Frazzon J, d'Azevedo PA, Frazzon APG. 2016. Resistance to antimicrobial agents among enterococci isolated from fecal samples of wild marine species in the southern coast of Brazil. *Marine Pollution Bulletin* **105**(1):51–57 DOI [10.1016/j.marpolbul.2016.02.071](https://doi.org/10.1016/j.marpolbul.2016.02.071).
- Saeedi B, Hällgren A, Jonasson J, Nilsson LE, Hanberger H, Isaksson B. 2002. Modified pulsed-field gel electrophoresis protocol for typing of enterococci. *APMIS* **110**(12):869–874 DOI [10.1034/j.1600-0463.2002.1101205.x](https://doi.org/10.1034/j.1600-0463.2002.1101205.x).
- Sánchez Valenzuela A, Benomar N, Abriouel H, Pérez Pulido R, Martínez Cañamero M, Gálvez A. 2012. Characterization of *Enterococcus faecalis* and *Enterococcus faecium* from wild flowers. *Antonie Van Leeuwenhoek* **101**(4):701–711 DOI [10.1007/s10482-011-9684-9](https://doi.org/10.1007/s10482-011-9684-9).
- Santestevan NA, De Angelis DZ, Prichula J, Pereira RI, Wachholz GR, Cardoso LA, De Moura TM, Medeiros AW, De Amorin DB, Tavares M, d'Azevedo PA, Franco AC, Frazzon J, Frazzon APG. 2015. Antimicrobial resistance and virulence factor gene profiles of *Enterococcus* spp. isolates from wild *Arctocephalus australis* (South American fur seal) and *Arctocephalus tropicalis* (Subantarctic fur seal). *World Journal of Microbiology and Biotechnology* **31**(12):1935–1946 DOI [10.1007/s11274-015-1938-7](https://doi.org/10.1007/s11274-015-1938-7).
- Sauget M, Valot B, Bertrand X, Hocquet D. 2017. Can MALDI-TOF mass spectrometry reasonably type bacteria? *Trends in Microbiology* **25**(6):447–455 DOI [10.1016/j.tim.2016.12.006](https://doi.org/10.1016/j.tim.2016.12.006).
- Schaumburg F, Onwugamba FC, Akulenko R, Peters G, Mellmann A, Köck R, Becker K. 2016. A geospatial analysis of flies and the spread of antimicrobial resistant bacteria. *International Journal of Medical Microbiology* **306**(7):566–571 DOI [10.1016/j.ijmm.2016.06.002](https://doi.org/10.1016/j.ijmm.2016.06.002).
- Schwarz S, Silley P, Simjee S, Woodford N, Van Duijkeren E, Johnson AP, Gastra W. 2010. Editorial: assessing the antimicrobial susceptibility of bacteria obtained from animals. *Journal of Antimicrobial Chemotherapy* **65**(4):601–604 DOI [10.1093/jac/dkq037](https://doi.org/10.1093/jac/dkq037).
- Sedgley C, Nagel A, Shelburne CE, Clewell DB, Appelbe O, Molander A. 2005. Quantitative real-time PCR detection of oral *Enterococcus faecalis* in humans. *Archives of Oral Biology* **50**(6):575–583 DOI [10.1016/j.archoralbio.2004.10.017](https://doi.org/10.1016/j.archoralbio.2004.10.017).
- Shankar V, Baghdayan AS, Huycke MM, Lindahl G, Gilmore MS. 1999. Infection-derived *Enterococcus faecalis* strains are enriched in *esp*, a gene encoding a novel surface protein. *Infection and Immunity* **67**(1):193–200 DOI [10.1128/IAI.67.1.193-200.1999](https://doi.org/10.1128/IAI.67.1.193-200.1999).
- Shao Y, Chen B, Sun C, Ishida K, Hertweck C, Boland W. 2017. Symbiont-derived antimicrobials contribute to the control of the Lepidopteran gut microbiota. *Cell Chemical Biology* **24**(1):66–75 DOI [10.1016/j.chembiol.2016.11.015](https://doi.org/10.1016/j.chembiol.2016.11.015).
- Singer AC, Shaw H, Rhodes V, Hart A. 2016. Review of antimicrobial resistance in the environment and its relevance to environmental regulators. *Frontiers in Microbiology* **7**:1728.

- Snyman M, Gupta AK, Bezuidenhout CC, Claassens S, Van den Berg J. 2016.** Gut microbiota of *Busseola fusca* (Lepidoptera: Noctuidae). *World Journal of Microbiology and Biotechnology* **32**(7):115 DOI [10.1007/s11274-016-2066-8](https://doi.org/10.1007/s11274-016-2066-8).
- Sutcliffe J, Tait-Kamradt A, Wondrack L. 1996.** *Streptococcus pneumoniae* and *Streptococcus pyogenes* resistant to macrolides but sensitive to clindamycin: a common resistance pattern mediated by an efflux system. *Antimicrobial Agents and Chemotherapy* **40**(8):1817–1824 DOI [10.1128/AAC.40.8.1817](https://doi.org/10.1128/AAC.40.8.1817).
- Teh BS, Apel J, Shao Y, Boland W. 2016.** Colonization of the intestinal tract of the polyphagous pest *Spodoptera littoralis* with the GFP-tagged indigenous gut bacterium *Enterococcus mundtii*. *Frontiers in Microbiology* **14**:1–12.
- Tenover FC, Arbeit RD, Goering RV, Mickelsen PA, Murray BE, Persing DH, Swaminathan B. 1995.** Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *Journal of Clinical Microbiology* **33**(9):2233–2239 DOI [10.1128/JCM.33.9.2233-2239.1995](https://doi.org/10.1128/JCM.33.9.2233-2239.1995).
- Usui M, Shirakawa T, Fukuda A, Tamura Y. 2015.** The role of flies in disseminating plasmids with antimicrobial-resistance genes between farms. *Microbial Drug Resistance* **21**(5):562–569 DOI [10.1089/mdr.2015.0033](https://doi.org/10.1089/mdr.2015.0033).
- Van Schooten B, Godoy-Vitorino F, McMillan WO, Papa R. 2018.** Conserved microbiota among young *Heliconius* butterfly species. *PeerJ* **6**:e5502 DOI [10.7717/peerj.5502](https://doi.org/10.7717/peerj.5502).
- Watkins RR, Bonomo RA. 2016.** Overview: global and local impact of antibiotic resistance. *Infectious Disease Clinics of North America* **30**:313–322 DOI [10.1016/j.idc.2016.02.001](https://doi.org/10.1016/j.idc.2016.02.001).
- Werner G, Hildebrandt B, Witte W. 2001.** The newly described *msrC* gene is not equally distributed among all isolates of *Enterococcus faecium*. *Antimicrobial Agents and Chemotherapy* **45**(12):3672–3673 DOI [10.1128/AAC.45.12.3672-3673.2001](https://doi.org/10.1128/AAC.45.12.3672-3673.2001).
- Zhang J, Wang J, Chen L, Yassin AK, Kelly P, Butaye P, Li J, Gong J, Cattley R, Qi K, Wanga C. 2017.** Housefly (*Musca Domestica*) and Blow Fly (*Protophormia Terraenovae*) as vectors of bacteria carrying colistin resistance genes. *Applied and Environmental Microbiology* **84**(1):e01736–17 DOI [10.1128/AEM.01736-17](https://doi.org/10.1128/AEM.01736-17).
- Zurek L, Ghosh A. 2014.** Insects represent a link between food animal farms and the urban environment for antibiotic resistance traits. *Applied and Environmental Microbiology* **80**(12):3562–3567 DOI [10.1128/AEM.00600-14](https://doi.org/10.1128/AEM.00600-14).