# ABSTRACTS





## 2nd Workshop on Actualistic Taphonomy

Rio Grande do Sul, Brazil

19-21 July 2021 – Online

Centro de Estudos Costeiros, Limnológicos e Marinhos, CECLIMAR, UFRGS Litoral Norte

Edited by Matias N. Ritter, Fernando Erthal, Rodrigo S. Horodyski

**ABSTRACTS VOLUME** 

https://www.ufrgs.br/taas/

doi: 10.5281/zenodo.5114543



#### AN ONLINE EVENT HELD BY







#### **ORGANIZING COMITTEE**

#### Prof. Matias Ritter

Departamento Interdisciplinar, Campus Litoral Norte, UFRGS

#### Prof. Fernando Erthal

Departamento de Paleontologia e Estratigrafia, Instituto de Geociências, UFRGS

#### Prof. Rodrigo Scalise Horodyski

Programa de Pós-Graduação em Geologia da Universidade do Vale do Rio dos Sinos, UNISINOS

#### **COLLABORATORS**

Anna Assumpção, Julia Ribeiro, Laura Porto Hornung, Luísa Crauss, Valentina Santos

#### SCIENTIFIC REFEREES

Claudio G. de Francesco

Fernando Erthal

Gabriela Hassan

Matias N. Ritter

Rodrigo S. Horodyski

Sabrina C. Rodrigues



#### **SUPPORT**



Sociedade Brasileira de Paleontologia

#### FINANCIAL SUPPORT

The 2nd TAAS is funded by CAPES (88887.470844/2019-00), and CNPq (403577/2019-5). Institutional support is provide by the Centro de Estudos Costeiros, Limnológicos e Marinhos (CECLIMAR), Campus Litoral Norte da UFRGS; Programa de Pós-Graduação em Geociências (PPGGeo); Programa de Pós-Graduação em Geologia da Universidade do Vale do Rio dos Sinos (Unisinos).







### PARASITIC LIKE-TRACES ON MARINE MOLLUSKS: NEW SOUNDS FROM SOUTHERN BRAZIL

Valentina Silva dos Santos<sup>1</sup>, Fernando Erthal<sup>1</sup>, Felipe Caron<sup>2</sup>, Sabrina Coelho Rodrigues<sup>3</sup> & Matias do Nascimento Ritter<sup>2</sup>

<sup>1</sup>Departamento de Paleontologia e Estratigrafia, Instituto de Geociências, UFRGS, Porto Alegre, RS. <sup>2</sup>Centro de Estudos Costeiros, Limnológicos e Marinhos, Campus Litoral Norte, Universidade Federal do Rio Grande do Sul, Av. Tramandaí, 976 - Imbé, RS. <sup>3</sup>Instituto de Ciências Exatas e Naturais do Pontal, Universidade Federal de Uberlândia, Rua Vinte, n1600, Ituiutaba, MG.

valensantoss@gmail.com, fernando.erthal@ufrgs.br, felipe.caron@ufrgs.br, sabrina.rodrigues@ufu.br, matias.ritter@ufrgs.br

Parasite traces on invertebrate fossil remains are relatively uncommon, despite its high biological information value regarding ecological interactions. Although these traces are not properly preservable, parasites are ubiquitous in the biosphere and their diversity and abundance is higher rather than predators. However, there is a vast unknowledge about the spatial and environmental variation of parasitic traces derived from the fossil record so far. The digenean trematode, for example, has a complex life cycle with three hosts, infesting mollusks bivalves in the second stage, being an intermediate host. Despite parasites be formed by a soft-bodied, they can produce diagnostic traces and preservable on bivalves, in the inner of the valves thus inducing de growth of oval-shapes modifying locally the shell geochemical composition. It has been observed that these deformities present in the structures, indicate that they were made when the host was still alive. This work aims to evaluate the prevalence of parasite like-traces caused by Digenean trematodes in bivalves' mollusks. To this purpose, 27 standardized samples of 0.05 m<sup>3</sup> of sedimentary material and mollusks shells were gathered, along about 135 km of coast, between the counties of Torres and Palmares do Sul, southern Brazil, simulating retrograding and prograding settings in the Quaternary fossil record. Twelve species of mollusks were identified, while the genus *Donax* displayed a frequency of 88.49 %, the most abundant species being the only one that displayed parasite like-traces. Prevalence values were significantly similar (Mann-Whitney-Wilcoxon test after sine arc transformation, p=0.956) in samples associated with retrograding (0.55) and prograding settings (0.59). The pattern found for studies carried out in the Mediterranean Sea — high prevalence of trematodes infesting bivalve in retrograding sectors —, was not identified here for the Southern Brazilian coast, which highlights the importance of carrying out more studies on the subject. [CNPq 422766/2018-6]