

**UNIVERSIDADE FEDERAL DO RIO GRANDE DO SUL
INSTITUTO DE BIOCÊNCIAS
PROGRAMA DE PÓS GRADUAÇÃO EM BOTÂNICA
ÊNFASE EM SISTEMÁTICA, EVOLUÇÃO E ECOLOGIA DE
ALGAS, PLANTAS E FUNGOS**

ANA PAULA VESTENA CASSOL

Diversidade de microalgas em sistema de cultivo de arroz orgânico e convencional com herbicidas: bentos, plâncton e epifítton

Porto Alegre

2022

ANA PAULA VESTENA CASSOL

Diversidade de microalgas em sistema de cultivo de arroz orgânico e convencional com herbicidas: bentos, plâncton e epifítton

Tese apresentada como requisito parcial para obtenção do título de Doutora em Botânica com ênfase em sistemática, evolução e ecologia de algas, plantas e fungos na Universidade Federal do Rio Grande do Sul.

Orientador(a): Dra. Lezilda Carvalho Torgan

Porto Alegre

2022

*Observar a natureza é encantador.
Compreender a natureza é fascinante.*

Às mulheres que me antecederam e não tiveram oportunidade de estudar.

AGRADECIMENTOS

À Dra. Lezilda Carvalho Torgan pela orientação na tese e pelos incríveis ensinamentos sobre diatomáceas. Muito Obrigada.

À Dra. Maria Angélica Oliveira pelos primeiros ensinamentos muito carinhosos sobre as algas, a ciência e posteriormente pela orientação no trabalho de graduação e mestrado. Muito obrigada.

Aos funcionários do Museu de Ciências Naturais da extinta Fundação Zoobotânica do Rio Grande do Sul (MCN-FZB/RS), pela infraestrutura utilizada indispensável para a realização deste estudo. Às amigas que construímos ao longo desse percurso. Rosana, Silvia, Laura, Jan, Julia, Ana, Vanessa e especialmente Vera e Ingrid. Foram pessoas incríveis e parceiras de ciência, café e chopp. Vera e Ingrid são mulheres cientistas, fortes e acolhedoras cujos ensinamentos levarei para a vida. Vocês são inspiradoras. Muito obrigada.

À Dra. Luciana de Souza Cardoso e Dra. Elisabeth Cabral pela realização dos estágios em docência. Muito Obrigada.

À Universidade Federal do Rio Grande do Sul e ao Programa de Pós-Graduação em Botânica pelo letramento científico, pela construção de um ambiente que promove a diversidade cultural, social e de pensamentos. A universidade pública e gratuita é fundamental em um país tão desigual como o nosso. Muito Obrigada.

À Coordenação de Aperfeiçoamento de Pessoal de Nível Superior-CAPES pela bolsa de doutorado concedida.

Aos meus pais que torceram, lá de Alegrete, pela realização do meu sonho, além de auxiliar na coleta de dados deste projeto. Muito Obrigada.

Às minhas amigas e amigos Suelen, Keyla, Liane, Mariana, Mariane, Tatiane, Juliana, Lurdes, Raquel, Kauana, Ana Paula, Daniéli, Maria Carolina, Francine, Eliza, Gabriele, Gabriela, Maurício, Ítalo que torceram por mim. Muito Obrigada.

Às minhas colegas de laboratório Gisele, Carol, Nicole, Daniela e o colega Eduardo pelas parcerias de conteúdos limnológicos. Muito Obrigada.

À Dra. Dávia Talgatti que gentilmente apresentou a técnica do “trapping method”, essencial para a realização deste estudo. Muito Obrigada.

À minha professora de inglês, Ana Júlia, que pacientemente guiou meus estudos no idioma. Muito Obrigada.

À Valdete Tavares presidente da associação dos produtores de arroz orgânico de Manoel Viana que foi muito solícita e gentilmente acolheu a ideia deste projeto. Muito Obrigada.

Ao meu amigo e colega Vinícius Bueno, cujo apoio e parceria foi fundamental no doutorado, principalmente no período de maior isolamento na pandemia. Muito Obrigada.

Ao Julian Stocker, meu companheiro neste último ano de doutorado, foi parceria e presença importante da vida. Muito Obrigada.

À minha versão no ano de 2017 que escreveu para a Dra. Lezilda solicitando orientação na tese, fez a prova e foi aprovada para iniciar os estudos em 2018. Migrar sozinha para a capital e entrar em uma universidade totalmente desconhecida foi desafiador. Na verdade, foi apavorante. O sonho em ser doutora na UFRGS foi a motivação. Ainda, exatamente na metade do curso, chegou a pandemia da COVID-19. Foram 7 meses afastada do trabalho presencial e muitas incertezas, mais do que as usuais. Foi novamente apavorante. Houve ajuda psicológica para o enfrentamento dos desafios da vida e da orientação. O desafio foi interno, pessoal. Sou a primeira geração universitária da família e estou muito grata por ter concluído este trabalho.

RESUMO

Agroecossistemas, como a lavoura de arroz, passam por intensas e rápidas transformações devido ao uso de fertilizantes e agrotóxicos que podem causar efeitos deletérios em organismos não alvo, como as microalgas. Dessa forma, o presente estudo teve como objetivo avaliar a variação da riqueza, composição e abundância de espécies de diatomáceas do sedimento e das microalgas encontradas no plâncton e epifítton da lavoura de arroz orgânico e convencional, com ênfase nos impactos dos herbicidas clomazone e penoxsulam. As amostragens foram efetuadas em campo em duas lavouras de arroz irrigado no estado do Rio Grande do Sul. Amostras de solo para análise das diatomáceas foram coletadas em três unidades amostrais e em quatro períodos: um dia antes da aplicação dos herbicidas e três datas posteriores. Amostras de água e a raspagem dos colmos do arroz foram realizadas em dois períodos após a aplicação. Foram verificados pH, temperatura, turbidez, condutividade elétrica e a concentração residual dos herbicidas no solo e na água. Variações da riqueza, composição e abundância das comunidades foram verificadas após a aplicação dos herbicidas. Menor riqueza de espécies no sistema de cultivo convencional foi observada na comunidade de diatomáceas do sedimento e epifítton que, provavelmente, estiveram maior tempo em contato com os herbicidas. Diferenças na composição de espécies entre os sistemas de cultivo foram mais evidentes na comunidade fitoplancônica e no sedimento. Redução e aumento na abundância de algumas espécies de diatomáceas foram verificadas no sedimento além de formas teratológicas em *Hantzschia amphioxys* f. *capitata*, *Luticola intermedia*, *L. cristinae*, *Navicula rostellata*, *Stauroneis lapponica* e especialmente em *Stauroneis reichardtii*. Espécies de diatomáceas foram registradas como características de cada sistema de cultivo. Na lavoura de arroz convencional destacaram-se em abundância as espécies: *Caloneis fontinalis*, *Hantzschia amphioxys* var. *amphioxys* f. *capitata*, *Luticola cristinae*, *Luticola fuhrmanni*, *Sellaphora* sp. (grupo pupula), *Nitzschia amphibia*, *Nitzschia perminuta*, *Nitzschia palea*, *Nitzschia brevissima* var. *brevissima*, *Neidium* sp2, *Pinnularia brebissonii*, *Pinnularia* aff *schimanskii*, *Pinnularia dubitabilis*, *Pinnularia borealis*, *Stauroneis borrichii*, *Stauroneis lapponica* e *Stauroneis reichardtii*. Na lavoura de arroz orgânico, destacaram-se as espécies: *Craticula* sp1, *Eunotia didyma* var. *gibbosa*, *Eunotia botuliformis*, *Eunotia longicamelus*, *Eunotia intermedia*, *Eunotia pseudosudetica*, *Frustulia guayanensis* subsp. *ecuadoriana*, *Frustulia saxonica*, *Hantzschia* sp3, *Humidophila contenta*, *Navicula leptostriata*, *Pinnularia rumrichae*,

Pinnularia subcapitata e *Surirella tenuissima*. Considerando a comunidade epifítica, Cyanophyceae e Chlorophyceae foram mais representativas na lavoura convencional e na lavoura orgânica, respectivamente. No fitoplâncton, Bacillariophyceae e Chlorophyceae foram mais representativas na lavoura convencional e na lavoura orgânica, respectivamente. O resultado da análise de indVal mostrou *Nitzschia* cf. *vixnegligenda* e *Stauroneis reichardtii* como espécies indicadoras para o fitoplâncton na lavoura convencional e *Radiococcus planctonicus* na lavoura orgânica. Em síntese, fitoplâncton, epifíton e as diatomáceas bentônicas mostraram diferença nos atributos quando comparados os sistemas de cultivo. A assembleia de diatomáceas terrestres evidenciou mais nitidamente alterações da riqueza, composição e no aparecimento de teratologias, como provável efeito do uso dos herbicidas.

Palavras-chave: diatomáceas terrestres, fitoplâncton, epifíton, arroz orgânico, arroz convencional, herbicidas.

ABSTRACT

Agroecosystems, as rice fields, are under intense transformations due to the use of fertilizers and pesticides. The agrochemicals may affect non-target organisms as microalgae. Thus, the present study aimed evaluate the richness, composition and abundance of the phytoplankton, epiphyton and in soil diatoms from organic and conventional rice fields highlighting the impacts of clomazone and penoxsulam herbicides. The sampling were conducted in two irrigated rice fields in Rio Grande do Sul state. Diatom soil samples were collected from three sample units in four periods: one day before the herbicides application and one, twelve and thirty-five days after the herbicides application. Phytoplankton and epiphyton samples were collected in the last two periods. Were obtained pH, temperature, turbidity, electrical conductivity and the residual herbicides concentration in the water and soil. Alterations in the richness, species composition and relative abundance were observed after the herbicides application. Low richness was observed in soil diatoms and in epiphyton community from the conventional rice field that probably were longer in contact with herbicides. Differences on the species composition between the organic and conventional rice fields were observed in the phytoplankton and in soil diatoms. Fluctuations in the relative abundance of some diatom species were recorded, besides the teratological forms in *Hantzschia amphioxys* f. *capitata*, *Luticola intermedia*, *L. cristinae*, *Navicula rostellata*, *Stauroneis lapponica* and especially in *Stauroneis reichardtii*. Some diatoms species were registered as characteristic from the each system. In the conventional rice field, highlighted in abundance the species: *Caloneis fontinalis*, *Hantzschia amphioxys* var. *amphioxys* f. *capitata*, *Luticola cristinae*, *Luticola fuhrmanni*, *Sellaphora* sp. (pupula group), *Nitzschia amphibia*, *Nitzschia perminuta*, *Nitzschia palea*, *Nitzschia brevissima* var. *brevissima*, *Neidium* sp2, *Pinnularia brebissonii*, *Pinnularia* aff *schimanskii*, *Pinnularia dubitabilis*, *Pinnularia borealis*, *Stauroneis borrichii*, *Stauroneis lapponica* e *Stauroneis reichardtii*. In the organic rice field, highlighted in abundance the species: *Craticula* sp1, *Eunotia didyma* var. *gibbosa*, *Eunotia botuliformis*, *Eunotia longicamelus*, *Eunotia intermedia*, *Eunotia pseudosudetica*, *Frustulia guayanensis* subsp. *ecuadoriana*, *Frustulia saxonica*, *Hantzschia* sp3, *Humidophila contenta*, *Navicula leptostriata*, *Pinnularia rumrichae*, *Pinnularia subcapitata* e *Surirella tenuissima*. Considering the epiphytic community, Cyanophyceae and Chlorophyceae were the most representative class in the conventional and organic rice field, respectively. In the phytoplankton, Bacillariophyceae and

Chlorophyceae were the most representative class in the conventional and organic rice field, respectively. The indVal analysis shown *Nitzschia cf. vixnegligenda* and *Stauroneis reichardtii* as indicators species to the phytoplankton in the conventional rice field and *Radiococcus planctonicus* in the organic rice field. In summary, phytoplankton, epiphyton and benthic diatoms showed differences in the attributes when compared the cultivation systems. The alterations in the richness, composition and teratologies were more pronounced in soil diatoms assemblage suggesting effect of the herbicides application.

Keywords: soil diatom, phytoplankton, epiphyton, organic rice field, conventional rice field, herbicide.

SUMÁRIO

INTRODUÇÃO	12
APRESENTAÇÃO	20
CAPÍTULO I	21
Alteration in the soil diatom assemblage after the herbicides application in the rice field.....	21
Abstract	
Introduction	
Materials and method	
Results	
Discussion	
References	
CAPÍTULO II.....	21
Soil diatoms from organic rice field: a little-known assemblage.....	22
Abstract	
Introduction	
Material and Methods.....	
Results	
Discussion	
References	
CAPÍTULO III	23
Do phytoplankton and epiphyton communities differ between organic and conventional rice fields?	23
Abstract	23
Introduction	24
Materials and methods	25
Results	28
Discussion	37
References	42
CONSIDERAÇÕES FINAIS.....	48

INTRODUÇÃO

A intensificação do uso dos recursos naturais desperta para uma discussão sobre a pressão antropogênica nos ecossistemas aquáticos. Recentemente, doze principais ameaças à biodiversidade em ambientes continentais foram identificadas por Reid *et al.* (2018). Mudanças climáticas, invasões biológicas, doenças infecciosas, contaminantes emergentes como medicamentos, cosméticos e agrotóxicos, florações de algas tóxicas, construção de hidrelétricas e nanomateriais fazem parte desta lista. Além disso, poluição por microplástico, poluição sonora e luminosa, aumento da salinização, declínio das concentrações de cálcio e o efeito cumulativo desses múltiplos estressores na biota aquática também foram citados. Dessa forma, destaca-se a importância de analisar os impactos gerados por esses fatores a fim de identificar práticas de gestão e conservação nos ambientes continentais.

Organismos fotossintéticos como as microalgas, desempenham um papel fundamental no metabolismo dos ecossistemas aquáticos participando das etapas de produção, consumo e decomposição (Wetzel 1996). Possuem uma grande diversidade de formas de vida e ocupam uma diversidade de habitat, podendo viver suspensas na coluna d'água, aderidas a substratos ou sobre os sedimentos (Wehr & Sheath 2003).

Dentre os principais grupos, as diatomáceas são particularmente conhecidas como fonte de informação sobre a qualidade ecológica de ambientes de água doce, através do conhecimento detalhado da ecologia e taxonomia dos táxons. Apesar da elevada abundância em ambientes aquáticos, algumas espécies capazes de ocupar os ambientes terrestres e enfrentar os desafios do ressecamento. Segundo Round *et al.* (1990) a classe Bacillariophyceae é um importante componente do microfitobento, pois contribui para a consolidação do sedimento por produção extracelular de substâncias de coesão e desempenham uma função essencial nos ciclos biogeoquímicos, que ocorrem na interface sedimento/água (ou sedimento/ar). Além disso, a capacidade de movimento no sedimento permite que as diatomáceas busquem por condições mais favoráveis a sua colonização, em direção a luz ou umidade (Lund 1945).

Considerando os agrotóxicos como um fator de risco para a segurança dos ecossistemas, faz-se necessária a avaliação das suas interações sobre os organismos e o ambiente. No Rio grande do Sul, a lavoura de arroz é produzida de forma convencional com uso de fertilizantes e agrotóxicos e de forma orgânica nos assentamentos rurais, ambos dispostos em áreas do bioma pampa.

A utilização dos agrotóxicos no sistema convencional promove elevados níveis de produção, contudo, uma série de desequilíbrios ambientais podem ser observados na microbiota do solo, na água e em organismos não-alvo (Guo *et al.* 2018). Evidências sugerem que o cultivo orgânico e suas práticas associadas beneficiam a riqueza e abundância de alguns animais e plantas nos agroecossistemas quando comparados aos cultivos convencionais (Katayama *et al.* 2019).

Em laboratório, experimentos com mesocosmos permitiram observar efeitos adversos na comunidade zooplanctônica em concentrações reais de herbicidas aplicados na lavoura (Reimche *et al.* 2015). De maneira semelhante, a comunidade fitoplanctônica responde a presença dos herbicidas imazapir e imazapique, na qual as formas coloniais como *Aphanocapsa*, *Eudorina*, *Pandorina* e *Spherozystis* foram caracterizadas como espécies sensíveis. Ainda nessa investigação, *Trachelomonas*, *Scenedesmus*, *Spirogyra* e *Anabaena* parecem relacionar-se de forma positiva com a presença desses compostos (Reck *et al.* 2018). Alterações nucleares e na ornamentação de frústulas de diatomáceas bentônicas também já foram observadas em laboratório, causadas por herbicida conhecido como genotóxico (Debenest *et al.* 2008). Além de modificações morfológicas, também foram identificadas respostas fisiológicas em espécies de microalgas. Herbicidas inibidores de fotossistemas (PSII) parecem favorecer espécies heterotróficas facultativas (Debenest *et al.* 2009; Larras *et al.* 2012).

Estudos em campo também sugerem alterações na biota aquática. A comunidade fitoplanctônica analisada em um lago na China com a presença de agrotóxicos, forneceu indicativos de que Bacillariophyta e Cryptophyta mostram afinidade para absorção de pesticidas organoclorados, já Cyanophyta e Chlorophyta mostram afinidades a compostos hidrocarbonetos aromáticos policíclicos (Zhao *et al.* 2015). Amostras de perifíton em rios da França em áreas agrícolas indicam um padrão de composição da comunidade de diatomáceas influenciada pela entrada de agrotóxicos e nutrientes (Morin *et al.* 2009). Em outras categorias taxonômicas, como em peixes e girinos, alterações toxicológicas através da avaliação de biomarcadores de estresse oxidativo também foram detectadas (Nunes *et al.* 2015; Santos *et al.* 2015).

Neste estudo, os herbicidas avaliados clomazone e penoxsulam, são amplamente utilizados na lavoura de arroz do sul do Brasil para controle de outras plantas. São compostos químicos moderadamente persistentes no solo variando em relação a diferentes condições. Fatores como sorção, dessorção, volatilização, degradação química e biológica vão interferir diretamente no comportamento desses produtos no ambiente

(Silva *et al.* 2012). Além das lavouras, a presença de clomazone e penoxsulam já foi detectada em águas superficiais de bacias hidrográficas no sul do Brasil (Marchesan *et al.* 2007; Primel *et al.* 2005, Silva *et al.* 2009).

O herbicida penoxsulam é um composto que inibe a acetolactato sintase (ALS), uma enzima responsável pela formação de proteínas com cadeia de aminoácidos ramificados envolvida no crescimento celular vegetal. Análises do residual do herbicida obtidos em campo, demonstraram que penoxsulam tem um período de meia vida entre 30 e 135 dias (Senseman 2007). Cabe ressaltar que a fotodegradação aquosa do composto ocorre rapidamente no ambiente, no entanto os produtos resultantes podem permanecer por longos períodos e podem ser difíceis de detectar (Jabusch & Tjeerdema 2006). Estresse oxidativo em organismos não-alvo como lagostins (Costa *et al.* 2018) mexilhões (Patetsini *et al.* 2013) e peixes (Cattaneo *et al.* 2010, 2012) foram encontrados em condições experimentais.

O herbicida clomazone pertencente ao grupo químico das isoxazolidinonas, cujo mecanismo de ação é inibir a biossíntese de carotenoides (Senseman 2007). Clomazone foi o herbicida mais frequente nas águas dos rios Vacacaí e Vacacai-mirim entre 2000 e 2003 na depressão central do RS, durante o período de cultivo do arroz irrigado (Marchesan *et al.* 2007). A persistência do herbicida já foi detectada por até 130 dias nas águas de lavoura de arroz (Zanella *et al.* 2002). O tempo de meia vida apresenta grande variação, sendo observado entre 11 e 126 dias (Du *et al.* 2018) e entre 8 e 32 dias (Noldin *et al.* 2001). A estabilização da lâmina d'água, através da irrigação contínua, acelera o processo de degradação do clomazone (Schreiber *et al.* 2017). Respostas toxicológicas em organismos não-alvo expostos ao clomazone foram demonstrados através de experimentos em girinos (Oliveira *et al.* 2016) em embriões de Zebrafish (Stevanovic *et al.* 2017) e em peixes (Murussi *et al.* 2015).

A lâmina d'água, formada pela prática de cultivo do arroz irrigado, é um importante ambiente temporário para diversas espécies de bactérias, algas, plantas, invertebrados e vertebrados, considerado um sítio de elevada dinâmica de nutrientes e biodiversidade (Roger *et al.* 1991). Nesse agroecossistema, além das algas planctônicas e as que crescem aderidas aos colmos da planta de arroz, fazem parte do ambiente as algas do sedimento da lavoura.

Estudos apontam as classes Chlorophyceae, Euglenophyceae e Bacillariophyceae como as mais representativas na comunidade fitoplânctônica e epifítica (Furtado & Lucca

2003; Alves da Silva & Tamanaha 2007; Reck *et al.* 2018; Sartori *et al.* 2011; Cassol *et al.* 2013). No entanto, a diversidade de diatomáceas no solo é pouco conhecida.

A composição de diatomáceas em solo de lavoura de arroz foi investigada somente em países asiáticos. Ao total, 92 espécies de diatomáceas foram registradas em lavouras de arroz convencional no Laos (Ohtsuka & Fujita 2001) e 53 espécies no Japão (Negoro & Higashino 1986). Entretanto, em solo de cultivo orgânico no Japão, foram registradas 104 espécies de diatomáceas (Fujita & Ohtsuka 2005).

Solos cultivados passam por intensas transformações devido aos processos físicos e químicos durante o ciclo de cultivo. Tais fatores podem ser selecionadores de espécies de diatomáceas em solo (Stanek-Tarkowska *et al.* 2017). Solos de formações florestais criaram um micro-habitat mais estável para as diatomáceas quando comparadas a variação temporal relacionada a outras práticas agrícolas como solos de lavoura e pastagem (Foets *et al.* 2020).

Tendo em vista esses aspectos, o estudo apresenta riqueza, abundância relativa e composição de espécies de diatomáceas do sedimento e das microalgas encontradas no plâncton e epifíton, em sistema de cultivo de arroz orgânico e convencional com uso dos herbicidas penoxsulam e clomazone.

Referências

- Alves da Silva SM, Tamanaha MS. 2008. Ocorrência de Euglenophyceae pigmentadas em rizipiscicultura na região do Vale do Itajaí, SC, Sul do Brasil. *Acta Botanica Brasilica* 22: 145-163.
- Cassol APV, Oliveira MA, Figueiredo MCS, Luz DS, Sartori GMS, Marchesan E. 2013. Microalgas em Cultura de arroz: influência de diferentes manejos de adubação em áreas com residual de herbicidas (imidazolinonas). *Iheringia Série Botânica* 68: 261-271.
- Cattaneo R, Clasen B, Loro VL, Menezes CC, Moraes B, Santi A, Toni C, Avila LA, Zanella R. 2010. Toxicological responses of *Cyprinus carpio* exposed to the herbicide penoxsulam in rice field conditions. *Journal of Applied Toxicology*, 31: 626–632.
- Cattaneo R, Moraes BS, Loro VL, Pretto A, MenezesC, Sartori GMS, Clasen B, Avila LA, Marchesan E, Zanella R. 2012. Tissue Biochemical Alterations of *Cyprinus carpio* exposed to Commercial Herbicide Containing Clomazone Under Rice-

- Field Conditions. *Archives of Environmental Contamination and Toxicology*. 62: 97–106.
- Costa R, Pereira JL, Santos MA, Pacheco M, Guilherme S. 2018. The role of contamination history and gender on the genotoxic responses of the crayfish *Procambarus clarkii* to a penoxsulam-based herbicide. *Ecotoxicology*, 27:908–918.
- Debenest T, Silvestre J, Coste M, Delmas F, Pinelli E. 2008. Herbicide effects on freshwater benthic diatoms: Induction of nucleus alterations and silica cell wall abnormalities. *Aquatic Toxicology*, 88(1): 88–94.
- Debenest T, Pinelli E, Coste M, Silvestre J, Mazzell N, Madigou C, Delmas F. 2009. Sensitivity of freshwater periphytic diatoms to agricultural herbicides. *Aquatic Toxicology*, 93: 11–17.
- Du P, Wu X, Xu J, Dong F, Liu X, Zhang Y, Zheng Y. 2018. Clomazone influence soil microbial community and soil nitrogen cycling. *Science of The Total Environment* 644: 475–485.
- Foets J, Wetzel CE, Teuling AJ, Pfister L. 2020a. Temporal and spatial variability of terrestrial diatoms at the catchment scale: controls on communities. *PeerJ* 8:e8296 DOI 10.7717/peerj.8296.
- Furtado RD, Luca SJ. 2003. Técnicas de cultivo de arroz irrigado: relação com a qualidade de água, protozoários e diversidade fitoplanctônica. *Revista Brasileira de Engenharia Agrícola e Ambiental* 7: 165–172.
- Fujita Y, Ohtsuka T. 2005. Diatoms from paddy fields in northern Laos. *Diatom* 21:71–89.
- Guo S, Xiong W, Xu H, Hang X, Liu H, Xu L, Li R, Shena Q. 2018. Continuous application of different fertilizers induces distinct bulk and rhizosphere soil protist communities. *European Journal of Soil Biology*, 88: 8–14.
- Jabusch TW, Tjeerdema RS. 2006. Photodegradation of Penoxsulam. *Journal of Agricultural and Food Chemistry*, 54(16): 5958–5961.
- Katayama N, Osada Y, Mashiko M. et al. 2019. Organic farming and associated management practices benefit multiple wildlife taxa: A large-scale field study in rice paddy landscapes. *Journal of Applied Ecology*, 56:1970–1981.
- Larras F, Bouchez A, Rimet F, Montuelle B. 2012. Using Bioassays and Species Sensitivity Distributions to Assess Herbicide Toxicity towards Benthic Diatoms. *PLoS ONE*, 7(8): e44458.

- Marchesan E, Zanella R, Avila LA, Camargo ER, Machado SLO, Macedo VRM. 2007. Rice herbicide monitoring in two Brazilian rivers during the rice growing season. *Scientia Agricola*, 64 (2): 131–137.
- Morin S, Bottin M, Mazzella N, Macary F, Delmas F, Winterton P, Coste M. 2009. Linking diatom community structure to pesticide input as evaluated through a spatial contamination potential (Phytopixal): A case study in the Neste river system (South-West France). *Aquatic Toxicology*, 94(1): 28–39.
- Morussi CR, Costa M, Menezes C, Leitemperger J, Guerra L, Lopes T, Severo E, Zanella R, Loro VL. 2015. Integrated Assessment of Biomarker Response in Carp (*Cyprinus carpio*) and Silver Catfish (*Rhamdia quelen*) Exposed to Clomazone. *Archives of Environmental Contaminant and Toxicology*, 68:646–654.
- Negoro K, Higashino M. 1986. Diatom vegetation of paddy fields in Japan. Report 1. Diatom vegetation of paddy fields in the vicinity of Sakurai City, Nara Prefecture. *Diatom* 2: 1-8.
- Noldin JA, Hermes LC, Fay EF, Eberhardt DS, Rossi MA. 2001. Persistência do herbicida clomazone no solo e na água quando aplicado na cultura do arroz irrigado, sistema pré-germinado. *Planta Daninha*, 19(3): 401–408.
- Nunes M, Silva FW, Costa-Silva D, Wallau GL, Posser T, Franco JL. 2015. Assessment of Water Pollution Signs in the Brazilian Pampa Biome Using Stress Biomarkers in Fish (*Astyanax* sp.). *Journal of Ecosystems*, 2015: 1–7.
- Ohtsuka T, Fujita Y. 2001. The diatom flora and its seasonal changes in a paddy field in Central Japan. *Nova Hedwigia* 73: 97-128.
- Oliveira CR, Fraceto LF, Rizzi GM, Salla RF, Abdalla FC, Costa MJ, Silva-Zacarin ECM. 2016. Hepatic effects of the clomazone herbicide in both its free form and associated with chitosan-alginate nanoparticles in bullfrog tadpoles. *Chemosphere*, 149: 304 – 313.
- Patetsini E, Dimitriadis VK, Kaloyianni M. 2013. Biomarkers in marine mussels, *Mytilus galloprovincialis*, exposed to environmentally relevant levels of the pesticides, chlorpyrifos and penoxsulam. *Aquatic Toxicology*, 126: 338– 345.
- Primel EG, Zanella R, MHS Kurz, Gonçalves FF Machado SLO, Marchezan E. 2005. Poluição das águas por herbicidas utilizados no cultivo do arroz irrigado na região central do estado do Rio Grande do Sul, Brasil: predição teórica e monitoramento. *Química Nova*, 28(4): 605–609.

- Reck L, Reimche GB, Alves CR, Oliveita MA, Machado SL. 2018. Efeito Dos Herbicidas Imazapir e Imazapique na Comunidade Fitoplanctônica em Lavoura de Arroz Irrigado. *Iheringia Série Botânica*, 73: 298–307.
- Reid AJ, Carlson AK, Creed IF, Eliason EJ, Gell PA, Johnson PTJ, Kidd KA, MacCormack TJ, Olden JD, Ormerod SJ, Smol JP, Taylor WW, Tockner K, Vermaire JC, Dudgeon D, Cooke SJ. 2018. Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biological Reviews*, 1-25.
- Reimche GB, Machado SLO, Oliveira MA, Zanella R, Dressler VL, Flores EMM, Gonçalves FF, Donato FF, Nunes MAG. 2015. Imazethapyr and imazapic, bispyribac-sodium and penoxsulam: Zooplankton and dissipation in subtropical rice paddy water. *Science of The Total Environment*, 514: 68–76.
- Roger PA, Heong KL, Teng PS. 1991. Biodiversity and sustainability of wetland rice production: role and potential of microorganisms and invertebrates. In: Hawksworth D.L. (ed.) *The Biodiversity of Microorganisms and Invertebrates: Its Role in Sustainable Agriculture*. Wallingford, CAB International. pp. 117–136.
- Round FE, Crawford RM, Mann DG. 1990. *The Diatoms. Biology and Morphology of the Genera*. Cambridge University Press, Cambridge.
- Santos T, Melo R, Costa-Silva DG, Nunes M, Rodrigues NR, Franco JL. 2015. Assessment of water pollution in the Brazilian Pampa biome by means of stress biomarkers in tadpoles of the leaf frog *Phyllomedusa iheringii* (Anura: Hylidae). *PeerJ*, 3: e1016.
- Sartori GMS, Marchesan E, Luz DS, Cassol APV, Figueiredo MCS, Oliveira MA, Silveira MVE, Ferreira RB. 2011. Manejo da adubação e seus efeitos na ocorrência de algas e na produtividade de arroz irrigado em áreas com residual de imidazolinonas. *Ciência Rural*, 8:1323-1330.
- Schreiber F, Scherner, A. Massey, JHS, Zanella Ro, Avila LA. 2017. Dissipation of Clomazone, Imazapyr, and Imazapic Herbicides in Paddy Water under Two Rice Flood Management Regimes. *Weed Technology*, 31(2): 330–340.
- Senseman SA. 2007. *Herbicide handbook*. Lawrence, KS: Weed Science Society Of America.
- Silva DRO, Avila LA, Agostinetto D, Magro TD, Oliveira E, Zanella R, Noldin JÁ. 2009. Monitoramento de agrotóxicos em águas superficiais de regiões orizícolas no sul do Brasil. *Ciência Rural*, 39(9): 2383–2389.

- Silva MSS, Cocenza DS, Rosa AH, Fraceto LF. 2012. Efeito da associação do herbicida clomazone a nanoesferas de alginato/quitosana na sorção em solos. *Quimica Nova*, 35: 102-107.
- Stanek-Tarkowska J, Czyż EA, Kaniuczak J, Poradowska A. 2017. Physicochemical properties of silt loamy soil and diversity of diatom species under winter wheat and oats. *Journal of Ecological Engineering*, 18(6): 142–151.
- Stenvanovic M, Gasica S, Pipal M, Blahova L, Brkic D, Neskovic N, Hilscherova K. 2017. Toxicity of clomazone and its formulations to zebrafish embryos (*Danio rerio*). *Aquatic Toxicology*, 188: 54–63.
- Wehr JD, Sheath RG. 2003. Freshwater habitats of algae. In: Wehr JD, Sheath RG. (Eds.) *Freshwater algae of the North America: ecology and classification*. USA: Elsevier Science.
- Wetzel RG. 1996. Benthic Algae and Nutrient Cycling in Lentic Fresh water Ecosystems. In: *Algal ecology: freshwater benthic ecosystems*. Edited by R. Jan Stevenson, Max I. Bothwell, Rex L. Lowe. Academic Press.
- Zanella R, Primel REG, Machado SLO, Gonçalves FF, Marchezan E. 2002. Monitoring of the herbicide clomazone in environmental water samples by solid-phase extraction and high-performance liquid chromatography with ultraviolet detection. *Chromatographia*, 55(9): 573–577.
- Zhao Z, Zhang Lu, Deng J, Wu J. 2015. The potential effects of phytoplankton on the occurrence of organochlorine pesticides (OCPs) and polycyclic aromatic hydrocarbons (PAHs) in water from Lake Taihu, China. *Environmental Science: Processes & Impacts*, 17(6): 1150–1156.

APRESENTAÇÃO

A tese está composta por três capítulos apresentados na forma de manuscritos a serem submetidos para a publicação em periódicos científicos.

O primeiro capítulo, “Alteration in the soil diatom assemblage after the herbicides application in the rice field”, trata do primeiro estudo da comunidade de diatomáceas de solo em área de arroz irrigado sob o impacto de herbicidas no Brasil. São apresentadas variações da riqueza, abundância relativa e da composição de espécies, antes e após a aplicação dos herbicidas. Os táxons são ilustrados em microscopia óptica. O manuscrito será submetido ao periódico *Acta Botanica Brasilica*.

O segundo capítulo, “Soil diatoms from organic rice field: a little known assemblage” trata das diatomáceas de solo em área de cultivo orgânico de arroz irrigado no Brasil. São apresentadas a riqueza, composição e abundância dos táxons, além de uma breve comparação entre os sistemas de cultivo orgânico e convencional. Os táxons são ilustrados em microscopia óptica. O manuscrito será submetido ao periódico *Acta Botanica Brasilica*.

Finalmente, o terceiro capítulo intitulado “Do phytoplankton and epiphyton communities differ between organic and conventional rice fields?” foi submetido ao periódico *Acta Botanica Brasilica* e aceito com publicação prevista para outubro de 2022. O manuscrito apresenta um estudo comparativo de riqueza, composição e abundância da comunidade fitoplanctônica e epifítica entre a lavoura de arroz irrigado orgânica e convencional.

CAPÍTULO I

Alteration in the soil diatom assemblage after the herbicides application in the rice field

Ana Paula Vestena Cassol^{1,3} 0000-0003-3662-9372,

Renato Zanella² 0000-0002-5971-1785,

Lezilda Carvalho Torgan¹ 0000-0001-6787-9441.

¹ Departamento de Botânica, Universidade Federal do Rio Grande do Sul, 9150-9900, Porto Alegre, RS, Brazil

² Departamento de Química, Universidade Federal de Santa Maria, 9710-5900, Santa Maria, RS, Brazil

³ Corresponding author: anapvcassol@gmail.com

Material aguardando publicação.

CAPÍTULO II

Soil diatoms from organic rice field: a little-known assemblage

Ana Paula Vestena Cassol^{1,2} 0000-0003-3662-9372,

Lezilda Carvalho Torgan¹ 0000-0001-6787-9441.

¹ Departamento de Botânica, Universidade Federal do Rio Grande do Sul, 9150-9900,
Porto Alegre, RS, Brazil

² Corresponding author: anapvcassol@gmail.com

Material aguardando publicação.

CAPÍTULO III

10.1590/0102-33062021abb0234

Original Article

Do phytoplankton and epiphyton communities differ between organic and conventional rice fields?

Ana Paula Vestena Cassol^{1*} <https://orcid.org/0000-0003-3662-9372>,

Renato Zanella² <https://orcid.org/0000-0002-5971-1785> and

Lezilda Carvalho Torgan¹ <https://orcid.org/0000-0001-6787-9441>

¹ Departamento de Botânica, Universidade Federal do Rio Grande do Sul, 9150-9900, Porto Alegre, RS, Brazil

² Departamento de Química, Universidade Federal de Santa Maria, 9710-5900, Santa Maria, RS, Brazil

* Corresponding author: anapvcassol@gmail.com

Received: July 20, 2021

Accepted: July 15, 2022

Abstract

This study aimed to determine if there are any differences in the attributes and composition of the phytoplankton and epiphyton communities between organic (OF) and conventional (CF) rice fields. We also strove to identify if there were any variations in these communities by comparing samples taken from two different periods (12 and 35 days) after the application of the herbicide clomazone and penoxsulam in CF. The farms are located in the Pampa Biome, Southern Brazil. Phytoplankton samples from the subsurface water and epiphyton samples from the rice stems were analyzed using the Utermöhl method. The CF and OF had distinct environmental conditions (pH, conductivity, and turbidity values), and the residual concentration of the herbicides decreased over time. There were no significant differences in epiphyton and phytoplankton density, or in phytoplankton richness, between the rice fields; only the epiphyton richness and taxonomic composition showed differences between the rice

fields. Cyanobacteria and Chlorophyceae comprised a large proportion of the epiphytic density in CF and OF, respectively. However, Bacillariophyceae and Chlorophyceae had greater densities in phytoplankton in CF and OF, respectively. The taxonomic composition of communities should be considered an effective tool to show the differences between the two cultivation systems.

Keywords: agroecosystem, artificial wetlands, community structure, herbicide, microalgae

Introduction

Wetlands are internationally recognized ecosystems that are subject to environmental protection and regulation (Millennium Ecosystem Assessment 2005). The recent classification proposed by Junk *et al.* (2013) defines three categories of Brazilian wetlands: coastal, continental, and artificial. According to this classification, artificial wetlands, such as rice farming, are the result of human activities.

It is possible to distinguish between both the conventional and organic rice cultivation systems. In Brazil, the current system in use is the conventional system that uses a large amount of pesticides, which threatens the biota and ecosystem functioning. Organic management contributes to improve the sustainability of agroecosystems by avoiding the use of chemical fertilizers and pesticides, and is based on integrative ecosystem services. According to Maltchik *et al.* (2017), encouraging sustainable practices in these areas may transfer some of the responsibilities in biodiversity conservation to production systems and, thus, contribute positively to regional biodiversity.

Rice fields are temporary environments occupied by a rich composition of fauna and flora (Bambaradeniya *et al.* 2004). The biota, especially microalgae, plays an important role in soil stabilization and fertility (Roger *et al.* 1991; Bellinger & Sigeo 2010). However, little attention has been paid to phytoplankton and epiphyton in the literature. Phytoplankton diversity was revealed by Shivakurama & Patar (2015) in a rice field in India. Additionally, the rice itself provides the substrate for periphyton colonization and growth and is used as a food source in rice–fish farming (Das *et al.* 2007; Saikia & Das 2011; 2014).

The microalgae composition in rice fields has already been registered on previous studies. Diatoms have mainly been investigated in Japan (Negoro & Higashino 1986; Ohtsuka & Fujita 2001; Fujita & Ohtsuka 2005). Furthermore, the pigmented members of Euglenophyceae were investigated by Alves da Silva & Tamanaha (2008) in Southern Brazil,

Cyanobacteria were studied by Irisarri *et al.* (2001) in Uruguayan rice fields and by Prasana & Nayak (2007) in India, and members of Chlorophyceae were investigated by Kumar & Sahu (2012) in India.

However, there is no field comparative information about algal communities in organic and conventional rice farming with herbicide application. Furthermore, the effects of herbicides on epiphytic and planktonic algae in rice fields have been described mainly in mesocosms (Sartori *et al.* 2011; Cassol *et al.* 2013; Reck *et al.* 2018; Liu *et al.* 2020). This knowledge gap motivated us to investigate the phytoplankton and epiphyton in the two rice cultivation systems. Quantifying the responses of these algal groups to agricultural systems will allow us to understand the limitations and benefits of the local biodiversity, and the results are relevant for implementing sustainable management practices with low environmental impacts (Katayama *et al.* 2019).

Our aim was to determine whether the attributes (richness, density, diversity, and equitability) and composition of phytoplankton and epiphyton communities differed between organic and conventional rice fields. The attribute variations after the application of clomazone and penoxsulam in the conventional field were also evaluated. Considering the differences in the environmental conditions of the rice fields, we expected to find distinctions among the attributes of the phytoplankton and epiphyton communities.

Materials and methods

Study area

The organic and conventional farms are located in Rio Grande do Sul, Southern Brazil at W 29°69' and S 55°85', and W 29°50' and S 55°66', respectively. The conventional farm (CF) is located 12 km from the urban center of Alegrete City (Fig. 1). A no-tillage cultivation system was used with chemical fertilization and was carried out according to the recommendations of the Irrigated Rice Brazilian Society (SOSBAI, *Sociedade Brasileira de Arroz Irrigado*, in Portuguese). The rice farm is located in the lowland of the Ibirapuitã River, which have traditionally been used for irrigated rice farming since 1986. The region is characterized by the predominant occurrence of basaltic volcanic rocks (with relief in the form of plateaus and escarpments) that are a part of the rocky and sandy riverbed (Hasenack *et al.* 2010). The Ibirapuitã River has an extension of 260 km in a sinuous course and an almost flat relief that floods in rainy seasons. These floods are supplied with organic load from livestock and agriculture, which are the predominant activities in the region.

The organic farm (OF) is located in Manoel Viana City in the Santa Maria do Ibicuí settlement, 22 km from the urban center and 45 km away from the CF (Fig. 1). Currently, 400 ha of rice are cultivated by 224 families belonging to the Association of Organic Producers of Santa Maria do Ibicuí Settlement (Ramos 2012). The rice water comes from the Ibicuí River, which extends 385 km into the lowlands and sandy riverbed in the Paleozoic lands of the Paraná sedimentary basin (Central Depression). This river has several tributaries, one of which is the Ibirapuitã River. The cultivation system used in the organic farm involved pre-germinated seedlings and does not use chemical fertilization or pesticides.




Figure 1. Location of the study sites of conventional farm (CF) and organic farm (OF) in the Ibicuí River watershed, Southern Brazil.

Sampling and data

Phytoplankton and epiphyton samples were taken after the flooding of the rice field in two periods (12 and 35 days after the spraying of herbicides). From both fields, three sample units were collected 600 m apart from each other. Three unit samples from the CF were taken on December 2, 2018 and on December 25, 2018: CF1 (29°69'09" 55°84'95"), CF2 (29°69'18" 55°85'11"), CF3 (29°69'74" 55°84'53"). Three unit samples from the OF were also taken at OF1 (29°50'26" 55°66'07"), OF2 (29°50'10" 55°66'35"), OF3 (29°50'21" 55°65'75").

Phytoplankton samples were collected from the surface of the rice water using a 100 mL glass bottle, and the epiphyton samples were collected via brushing three rice stems of 3 cm lengths, totaling a composite sample of 25 cm². This was done in both the conventional and organic fields. The samples were preserved in Lugol's alkaline solution (Sournia 1978) and housed at Alarich Schultz Herbarium (number 110934 – 110957), Museu de Ciências Naturais, Fundação Zoobotânica do Rio Grande do Sul in Porto Alegre, Brazil.

The densities of phytoplankton (ind.mL⁻¹) and epiphyton (ind.cm⁻²) were estimated according to the method of Uthermöhl (1958) in 5 mL chambers, using an inverted microscope (Zeiss Axiovert) after 24h of sedimentation. This was done via counting in transect fields, with the aim of achieving an efficiency of 80 % (Pappas & Stoermer 1996). Only individuals with plastids and those that were alive at the time of collection were quantified. Identification was performed using an inverted Zeiss microscope (640× magnification). Richness was estimated based on the number of taxa at generic and infra-specific levels. The abundant species was determined according to the method of Lobo & Leighton (1986). Shannon's diversity (H') and equitability indices (E) were used as measures of community structure.

The presence and concentration of residual herbicides (penoxsulam and clomazone) in the water were detected using the solid phase extraction method. These were then analyzed via liquid chromatography with mass spectrometry (LC-MS/MS) in the Pesticide Residue Laboratory (LARP) at the Federal University of Santa Maria, according to the method of Donato *et al.* (2012).

The water temperature (temp), hydrogen potential (pH), electrical conductivity (Cond), dissolved oxygen (DO), turbidity (Turb), and total dissolved solids (TDS) were measured in situ using a Horiba U-50 – Multiparameter probe. The data from the conventionally farmed field were obtained in the morning, but with the organically farmed field the measurements were made in the afternoon on the same sampling day.

After logarithmic transformation of the data to achieve normality, we used the two-sample T test to determine if there are differences between the periods and the cultivation systems in

both communities. Permutational analysis of variance (PERMANOVA) was performed to verify differences in the composition of species in the phytoplankton and epiphyton communities of the rice fields. Nonmetric multidimensional scaling (NMDS) was applied to the abundant species matrix to investigate the species distribution at the sites of the organically and conventionally farmed rice fields. The community matrices were transformed using Wisconsin double standardization and a Bray–Curtis dissimilarity matrix was calculated using the vegan “metaMDS” function in R. The IndVal method (Dufrêne & Legendre 1997) was used to identify indicator species in the cultivation systems. An overall analysis was performed using the R software (R Development Core Team 2021) and the Vegan package (Oksanen *et al.* 2020).

Results

Physical and chemical water conditions

The physical and chemical rice water conditions are presented in Table 1. The water temperature varied between 18.4 °C and 24.5 °C in the CF and between 22.2 °C and 31.0 °C in the OF.

The pH varied from neutral to slightly acidic in the CF, and was strongly acidic in the OF. The electrical conductivity showed a slight increase in the CF, varying between 60 $\mu\text{s cm}^{-1}$ and 100 $\mu\text{s cm}^{-1}$, whereas in the OF, the maximum electrical conductivity found in the study period was 20 $\mu\text{s cm}^{-1}$.

Turbidity varied between a minimum of 2.48 NTU and maximum of 31.20 NTU in the CF, and was greater in the OF (62 - 542 NTU). Meanwhile, total dissolved solids varied from 0.03 mg L^{-1} to 0.06 mg L^{-1} in the CF and between 0.01 mg L^{-1} and 0.02 mg L^{-1} in the OF. Dissolved oxygen showed higher values in the CF (6.5 - 14.8 mg L^{-1}) than those in the OF (4.0 - 7.8 mg L^{-1}).

Table 1. Physical and chemical variables of sample units in the conventional farm in Alegrete City, RS, and in the organic farm in Manoel Viana City, RS, 12 and 35 days after the spraying of herbicides in three sample units in periods 1 (12/02/2018) and period 2 (12/25/2018). These sample units were taken at the same day.

Farming	Period	Sample Unit	Temp (°C)	pH	Cond ($\mu\text{s cm}^{-1}$)	Turb (NTU)	DO (mg L^{-1})	TDS (g L^{-1})
Conventional	Period 1	CF 01	18.4	6.4	90	2.7	9.2	0.06
		CF 02	22	6.9	90	2.5	12.0	0.05
		CF 03	21.9	7.4	100	4.5	14.8	0.06
	Period 2	CF 01	22.8	6.1	60	31.2	7.7	0.04
		CF 02	22.9	6.2	80	24.7	6.5	0.05
		CF 03	24.5	6.4	60	20.8	7.3	0.03
Organic	Period 1	OF 01	31	5.3	10	243	7.8	0.01
		OF 02	28.4	5.8	20	174	7.3	0.01
		OF 03	28.8	5.0	20	542	4.0	0.01
	Period 2	OF 01	22.8	5.4	20	100	4.4	0.01
		OF 02	22.2	5.3	20	172	5.4	0.02
		OF 03	23.3	4.6	10	62	4.9	0.01

Herbicide concentrations in the conventionally farmed rice field

The residual concentrations of herbicides in the conventionally farmed fields and in the channel that takes water from the Ibirapuitã River to the crops are presented in Table 2. Average values of $1.29 \mu\text{g L}^{-1}$ and $0.28 \mu\text{g L}^{-1}$ were detected for clomazone and penoxsulam, respectively. The residual concentrations of the two herbicides decreased over time, and penoxsulam, which was not detected in some samples after 35 days of spraying, dissipated faster than clomazone.

Table 2. Residual concentrations of herbicides from the conventional farm after 12 days (period 1) and 35 days of spraying (period 2). Nd=non detected.

Period	Sample Unit	Clomazone ($\mu\text{g L}^{-1}$)	Penoxsulam ($\mu\text{g L}^{-1}$)
Period 1	CF 01	1.402	0.225
	CF 02	1.336	0.322
	CF 03	1.140	0.304
Period 2	CF 01	0.297	n.d.
	CF 02	0.433	0.091
	CF 03	0.184	n.d.

Phytoplankton community

The phytoplankton species richness in the CF ranged from 24 to 26 species in period 1 and from 25 to 36 species in period 2. In the OF, the richness ranged from 17 to 25 species in period 1 and from 14 to 40 species in period 2 (Fig. 2). There was no significant variation in richness between periods in either CF or OF ($p > 0.05$). There was also no significant difference in richness between the two cultivation systems ($p > 0.05$).

Specific diversity showed higher values in period 2 (2.43 – 3.05 bits. ind⁻¹) than in period 1 (1.38 - 2.23 bits. ind⁻¹) in the CF. The specific diversity seen in OF was greater in period 2 (1.81 - 2.79 bits. ind⁻¹) than in period 1 (0.45 - 2.15 bits. ind⁻¹) (Fig. 2). The equitability in the CF showed higher values in period 2 (0.72 - 0.84) than in period 1 (0.43 – 0.69) and the equitability in the OF showed higher values in period 2 (0.68 - 0.79) than in period 1 (0.15 - 0.66).

The total density of phytoplankton in CF obtained higher values in period 1 (699 - 8244 ind. mL⁻¹) than in period 2 (175 - 1094 ind. mL⁻¹). In the OF, the density obtained higher values in period 1 (4368 - 62103 ind. mL⁻¹) than in period 2 (374 - 1313 ind. mL⁻¹) (Fig. 2). There was no significant difference in density between the periods in CF, but there was a significant difference in density between the periods in OF ($p < 0.05$). Notably, there were no significant differences between the crops in these fields ($p < 0.05$).




Figure 2. Phytoplankton density (ind.mL⁻¹), richness, diversity, and equitability in conventional (CF) and organic (OF) farms in three samples from the two experimental periods.

The NMDS model showed species distribution according to the organic and conventional fields (Fig. 3). Moreover, the difference in species composition was statistically significant according to PERMANOVA analyses ($p = 0.0003$, $R^2 = 0.22$). The results of the indicator species analysis between the cultivation systems showed that *Nitzschia* cf. *vixnegligenda* Lange-Bertalot & Hofmann and *Stauroneis reichardtii* Lange-Bertalot, Cavacini, Tagliaventi & Alfinito were associated with CF ($p = 0.01$), and *Radiococcus planctonicus* J.W.G.Lund was associated with OF ($p = 0.01$). We identified 86 species belonging to seven classes in CF. The most representative class in terms of density was Bacillariophyceae, followed by Chlorophyceae and Cyanophyceae in period 1; this was followed by Euglenophyceae and Cryptophyceae in period 2. In period 1, within Bacillariophyceae there were abundant *Nitzschia palea* (Kützing) W. Smith and *Nitzschia gracilis* Hantzsch. Within Chlorophyceae, the abundance of *Dictyosphaerium* sp., *Monoraphidium caribeum* Hindák, and *Schoederia* sp. were apparent. Among the other classes,

we also noted the presence of *Aphanocapsa* sp., *Pseudanabaena* sp. (Cynaophyceae), *Trachelomonas curta* A. M. Cunha (Euglenophyceae), *Cryptomonas ovata* Ehrenberg (Cryptophyceae), *Actinotaenium perminutum* (G. S. West) Teiling, and *Zygnema* sp. (Zygnematophyceae). In period 2, within Bacillariophyceae, *Encyonema silesiacum* (Bleisch) D. G. Mann 1990, *Navicula incarum* U. Rumrich & Lange-Bertalot, *N. palea*, *N. gracilis*, *Pinnularia subcapitata* Gregory, and *S. reichardtii* were abundant. Within Euglenophyceae, the presence of *Trachelomonas verrucosa* var. *granulosa* (Playfair) Conrad & Meel was highlighted, as well as that of *Cryptomonas ovata* Ehrenberg within Cryptophyceae.




Figure 3. Nonmetric multidimensional scaling (NMDS) ordination of abundant phytoplankton species (stress value = 0.05) in the sample units (●) taken in two time periods from organic and conventional rice fields.

In the OF, we identified 79 species belonging to seven classes. Chlorophyceae followed by Zygnematophyceae were dominant in period 1, with abundance of *Dyctiosphaerium* sp., *Kirchneriella irregularis* (G. M. Smith) Korshikov, *Monoraphidium caribeum* Hindák, *M. flexuosum* Komárek, *M. griffithii* (Berkeley) Komárková-Legnerová, *Radiococcus planktonicus*,

Actinotaenium perminutum (G.S.West) Teiling, *Cosmarium* sp3, *N. palea*, and *C. ovata*. In period 2, the composition of the community was changed to having a dominance of Bacillariophyceae (*Nitzschia* spp.) in sample units 1 and 2, and Euglenophyceae (*Monomorpha* sp., *T. verrucosa* var. *granulosa*, *T. curta*, and *Cryptomonas* sp3) in sample unit 3 (Fig. 4).




Figure 4. Relative contribution (total number of individuals) of seven classes of phytoplankton in the conventional farm (a) and organic farm (b) in the three sample units from the two experimental periods (P1, P2).

Epiphytic community

The epiphyton species richness between the two systems and in the CF between the periods in the experiment was significantly different ($p < 0.05$). Higher values were observed

in period 1 (18-23 species) than in period 2 (7-17 species). In the OF, the richness varied between 30 species and 38 species in period 1 and between 10 species and 39 species in period 2; these values were not significantly different ($p < 0.05$) (Fig. 5).

Epiphyton density did not differ significantly between systems ($p > 0.05$). However, the density did vary between 20.315 ind. cm² and 36.216 ind. cm² in period 1, and between 15.355 ind. cm² and 158.058 ind. cm² in period 2 in the CF. In the OF, higher values were observed in period 2 (115.542 - 1.986782 ind. cm²) than in period 1 (50.873 – 248.288 ind. cm²). Significant differences in density between the collection periods were observed only in the OF ($p < 0.05$) (Fig. 5).

The epiphyton diversity index showed higher values in period 1 (1.24 - 2.19 bits. ind⁻¹) than in period 2 (0.71 - 1.96 bits. ind⁻¹) in the CF. In the OF, the attribute showed higher values in period 1 (1.93 - 2.63 bits. ind⁻¹) than in period 2 (0.11 - 1.57 bits. ind⁻¹). In the CF, the equitability ranged from 0.42 to 0.70 in period 1 and from 0.36 to 0.69 in period 2. In the OF, the values ranged from 0.54 to 0.72 in period 1 and ranged from 0.05 to 0.55 in period 2 (Fig. 5).




Figure 5. Epiphyton density (ind.cm²), richness, diversity and equitability in conventional (CF) and organic (OF) farms in three samples from the two experimental periods.

The NMDS model showed distribution of abundant epiphytic species corresponding to OF and CF (Fig. 6). However, according to PERMANOVA analyses, there was no significant difference in composition between the two cultivation systems ($p > 0.05$). Moreover, there were no epiphytic indicator species in either of the cultivation systems.




Figure 6. Nonmetric multidimensional scaling (NMDS) ordination of abundant epiphyton species (stress value = 0.06) in sample units (●) taken in two time periods from organic and conventional rice fields.

We identified 54 species belonging to six classes in the CF. In period 1, the most representative classes in density were Cyanophyceae and Bacillariophyceae, with an abundance of *Anagnostidinema amphibium* (C. Agardh ex Gomont) Strunecký, Bohunická, J. R. Johansen & J. Komárek ; *N. palea*; and *N. cf. vixnegligenda* . In period 2, Cyanophyceae (*A. amphibium*), Cryptophyceae (*Cryptomonas erosa* Ehrenberg , *Chroomonas cf. coerulea* (Geitler) Skuja, and Bacillariophyceae (*Nitzschia* spp.) were highlighted in sample unit 3 (Fig. 7).

We identified 91 epiphytic species in the OF. In period 1, Chlorophyceae was predominant, followed by Zygnematophyceae, with abundance of *Dictyosphaerium* sp., *K. lunaris* var. *irregularis*, *Monoraphidium circinale* Nygaard, *M. griffithii* (Berkeley) Komárková-Legnerová, *Oedogonium* sp., *R. planktonicus*, *Actinotanenium perminutum*, and *N. palea*. Cyanophyceae and Cryptophyceae dominated in period 2, with *A. amphibium*, *C. erosa*, and *C. cf. coerulea* as the main species (Fig. 7).




Figure 7. Relative contribution (total number of individuals) of different classes of epiphyton from the conventional farm (a) and organic farm (b) in three sample units from the two experimental periods (P1, P2).

Discussion

Organic versus conventional farming

Organic and conventional farming have different planting systems. Pre-germinated planting is used in the OF, where flooding occurs during soil preparation and there is the turning over of residual plants from the previous crop; whereas, in the CF, flooding occurs after the spraying of herbicides and no-tillage is used (SOSBAI 2018). These planting systems result in different environmental conditions. In the rice water in OF, higher turbidity values were recorded (> 62 NTU) because of the suspension of solids that is promoted by the practice of turning over residual plants (Furtado & Luca 2003). The lower values of pH observed (4.6 - 5.8) is because the anaerobic conditions in flooded soil as a result of decomposition process that produces organic acids (Bohnen *et al.* 2005). Lower pH values were obtained in rice fields in India that were organically fertilized (Sihi *et al.* 2020). The lower water stream flow in the

OF due to cultivation in the plots may explain the high temperature and low oxygen concentration in the water during the experimental period. Lower values of dissolved oxygen (4 - 7.8 mg L⁻¹) and pH were also observed in the OF when compared to those in the CF (Suárez-Serrano *et al.* 2010).

In the CF, the values of pH and dissolved oxygen were close to the ones registered in the Ibirapuitã River (7.1 - 9.1 and 7.02 - 9.06 mg L⁻¹, respectively) in 2018, according to the ANA (2018). The water pH ranging from slightly acidic to neutral from the CF is similar to mesocosms that underwent herbicide spraying and the control treatment (Reck *et al.* 2018; Reimche *et al.* 2015).

The values of dissolved solids and electrical conductivity were higher in CF than in OF. This result was also observed under the same conditions of rice cultivation in India due to the chemical fertilization utilized in conventional fields (Sihi *et al.* 2020). In contrast, the values of electrical conductivity in CF and OF were higher than those in the Ibirapuitã (37 - 55.8 µs cm⁻¹) and Ibicuí rivers (46.2 - 60.4 µs cm⁻¹), respectively, in 2018 (ANA 2018). This was most likely due to the higher organic matter concentration.

Effects of penoxsulam and clomazone in the environment

The herbicide penoxsulam inhibits acetolactate synthase, an enzyme responsible for building protein with a branched amino acid chain, whose inactivation results in a protein deficit that affects cell multiplication (Senseman *et al.* 2007). The results of herbicide residue analysis from rice water showed that penoxsulam degraded faster than clomazone in the environment and, after 35 days the application of the herbicide, was not detected in most of the sample units. Contact with water promotes the solubilization of chemicals and occurs in a different way for each herbicide according to its own characteristics. For example, the solubility of clomazone is higher than that of penoxsulam (Senseman *et al.* 2007). It is important to note that the aqueous photodegradation of herbicides occurs quickly in the environment; however, their products can remain for long periods and can be difficult to detect (Jabusch & Tjeerdema 2006). In addition, penoxsulam concentrations were detected during the period of rice cultivation in the superficial waters of rivers in Rio Grande do Sul (Silva *et al.* 2009), where it possibly affects other organisms.

The other herbicide evaluated was clomazone, which belongs to the isoxazolidinone chemical group and acts as an inhibitor of carotenoid biosynthesis (Senseman *et al.* 2007). Clomazone was the most frequently herbicide found in the Vacacaí and Vacacai-Mirim rivers, during rice cropping between 2000 and 2003, in the central region of Rio Grande do Sul

(Marchesan *et al.* 2007). In this study, clomazone showed a slower degradation rate than penoxsulam, which was detected in the second sampling period, 35 days after herbicide spraying. The persistence of clomazone was detected up to 130 days in rice water in the central region of Rio Grande do Sul (Zanella *et al.* 2002). Notably, the half-life of clomazone varies in the literature, ranging from 11 to 126 days (Du *et al.* 2018) and from 8 to 32 days (Noldin *et al.* 2001).

Phytoplankton and epiphyton associated with cultivation systems

Research on the algae community in organic and conventional rice fields is scarce in the literature. Comparatively, the total phytoplankton richness was similar between the two cultivation systems (79 spp. in OF and 86 spp. in CF) and was also close in value to that of natural wetlands within the territorial limits of the Pampa biome (74 taxa) (Matsubara *et al.* 2008).

In mesocosms, however, greater values of phytoplankton community richness were found in rice cultivated without the spraying of pesticides but with residual imidazolinone (151 species), such as in rice cultivated with imazapyr and imazapic herbicides (60 genera, with 44 genera observed by us), according to Cassol *et al.* (2013) and Reck *et al.* (2018), respectively. Baert *et al.* (2016) exposed the plankton diatom community at different levels of richness to different concentrations of atrazine herbicide and observed that more diverse communities were less affected by herbicide stress.

The total epiphyton richness (42 genera) was similar to the rice-fish farm in India (38 genera) (Das *et al.* 2007); this was, however, less than another rice-fish farm (97 genera) (Saikia & Das 2011). The number of epiphytic taxa (about 90 species) was similar to that found in a mesocosm under residual imidazolinone. This mesocosm was an area of cultivated rice located close to the region of this study (Cassol *et al.* 2013). The results of different studies do not seem to illustrate a unique pattern of phytoplankton and epiphyton richness between the cultivation systems.

In this study, we expected to find differences in community attributes owing to the differences in environmental conditions. However, only the difference in epiphyton richness was statistically significant. The phytoplankton community had a taxonomic homogenization after 12th and 35th days of herbicide application. The effects of herbicides on algal growth possibly occurred in the first days after spraying, followed by recovery events in the community during this time. Faster recovery events observed in other studies, associated with the phytoplankton being subjected to the water flow, can hinder the detection of punctual impacts

because the evaluation was not done 24 h after herbicide spraying. Imazapyr and imazapic herbicides, whose mechanism of action is the same as that of penoxsulam, inhibited the growth of the phytoplankton community 24 h after the spraying in mesocosms (Reck *et al.* 2018). Residual water from CF after spraying with herbicides, such as clomazone, inhibited the growth of *Chlorella vulgaris*, *Pseudokirchneriella subcapitata*, and *Desmodesmus subcapitatus* strains 2 days after spraying (Suárez-Serrano *et al.* 2010).

The decrease in epiphyton richness after herbicide spraying may be the result of the longer contact time with the chemical compound. At the same time, the species has strategies, such as mucous tubules and peduncles, for resisting environmental pressure (Rimet & Bouchez 2011). According to the sensitivity assessment, the periphyton remains exposed for longer to the pollutant that promotes community growth (Lobo *et al.* 2002).

Differences between the algal communities subjected to conventional and organic farming were observed in the phytoplankton species composition, particularly in period 1 of the experiment. In the phytoplankton of CF, Bacillariophyceae was the most representative class for the total number of individuals, with a predominance of *Nitzschia* spp. This genus is common in wetlands and tolerates high nutrient and organic matter concentrations (Sheat & Wehr 2003; Trobajo & Sullivan 2010). *Nitzschia* cf. *vixnegligenda* is an indicator species in conventional rice fields. This is a little-known taxon that has been recorded in high-electrolyte streams in Ecuador and Central Europe (Rumrich *et al.* 2000). *Nitzschia palea* was also abundant; this species is distributed in different lotic and lentic environments, and is associated with environments under stress. Trobajo *et al.* (2009) recognized this species as being tolerant to atrazine, a photosystem II inhibitor herbicide (Debenest *et al.* 2010). *Stauroneis reichardtii* has also been reported to be an indicator species in conventional rice fields. This cosmopolitan species is found in circumneutral waters with low specific conductance (Bahls 2010).

The relationships between species composition and periods after flooding were verified. Euglenophyceae and Cryptophyceae increased densities in the phytoplankton community in period 2. Cyanophyta and Euglenophyta showed a positive relationship with these factors in mesocosms, whereas Chlorophyta, Bacillariophyta, Cryptophyta, and Dinophyta showed a decrease in density (Furtado & Luca 2003). Cryptophyceae belongs to the Y functional group; it is tolerant to low luminosity and, therefore, uses the phagotrophic mode to obtain energy (Reynolds *et al.* 2002). The incidence of solar radiation in rice water decreased over the time (Cassol *et al.* 2013), which may favor the growth of heterotrophic species in period 2. In experiments, the photosystem II inhibitor herbicide may favor facultative heterotrophic species, thus preventing their deleterious effects (Debenest *et al.* 2009; Larras *et al.* 2012).

In OF, Chlorophyceae was abundant among phytoplankton during period 1 of the experiment. The association between higher turbidity recorded in this study and the presence of *R. planktonicus* has been previously described (Reynolds *et al.* 2002). The density of green algae decreased consistently during period 2. The same result was observed for higher herbicide concentrations in rice farming, showing the sensitivity of species to photosystem inhibitor chemicals (Cochard *et al.* 2014); the effect was also noted ecotoxicologically (Suárez-Serrano *et al.* 2010). *Radiococcus planktonicus*, an indicator species of organic farming, has cells that are enveloped by a mucilaginous sheath making it easy to float. It has been suggested that colonial genera such as *Aphanocapsa*, *Eudorina*, *Pandorina*, and *Sphaerocystis* are sensitive to the presence of imazapyr and imazapic herbicides which belong to the penoxsulam chemical group in rice farming mesocosms (Reck *et al.* 2018). The low abundance of green algae in CF and their predominance in OF suggest the effects that herbicide sprays have on the environment. Heterotrophic groups were also abundant in OF in period 2.

In the epiphyton, Bacillariophyceae and Cyanophyceae made notable contributions to the composition of the communities in both rice farms. However, Chlorophyceae had a higher contribution in the OF, mainly in period 1. The preference of Chlorophyceae and Cyanophyceae in colonizing the rice stem was also observed in rice–fish farming without pesticides in India, where a natural substrate with a rough surface promoted the development of even more weakly attached forms of algal flora when compared to a glass slide that was preferentially colonized by Bacillariophyceae (Saikia & Das 2011). The authors suggested that there might be a relationship between the life cycle of the substrate and periphyton diversity even though they had not been able to describe such a relationship. In bioassays with periphytic communities, the herbicide metribuzin, a photosystem II inhibitor, negatively affected Chlorophyta, whereas diatoms and Cyanobacteria showed an increase in biomass under its exposure (Gustavson *et al.* 2003).

The results of this study showed that organic and conventional farming had distinct environmental conditions and expressed differences community attributes; only epiphyton richness showed differences between conventional and organic fields. Therefore, phytoplankton and epiphyton taxonomic compositions should be considered as an effective tool to illustrate the differences between the two systems.

Acknowledgements

We would like to thank CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) for the doctoral scholarship to APVC. We are grateful to Daniel Dutra Saraiva for supporting our efforts in statistical analyses. We would also like to thank the Associação dos Produtores de Arroz Orgânico from Assentamento Santa Maria do Ibicuí for the availability of the study area.

References

- Alves-da-Silva SM, Tamanaha MS. 2008. Ocorrência de Euglenophyceae pigmentadas em rizipiscicultura na região do Vale do Itajaí, SC, Sul do Brasil. *Acta Botanica Brasilica* 22: 145-163.
- ANA- Agência Nacional de Águas. 2018. Portal HidroWeb do Sistema Nacional de Informações sobre Recursos Hídricos (SNIRH). <https://www.snirh.gov.br/hidroweb/apresentacao>. 18 Nov. 2019.
- Baert JM, De Laender F, Sabbe K, Janssen CR. 2016. Biodiversity increases functional and compositional resistance, but decreases resilience in phytoplankton communities. *Ecology* 97: 3433–3440.
- Bahls LL. 2010. Stauroneis in the Northern Rockies. In: Bahls, LL. (ed.) *Northwest Diatoms*, v.4. The Montana Diatom Collection pp.1-172.
- Bambaradeniya CNB, Edirisinghe JP, De Silva DN, Gunatilleke CVS, Ranawana KB, Wijekoon S. 2004. Biodiversity associated with an irrigated rice agro-ecosystem in Sri Lanka. *Biodiversity and Conservation* 13: 1715–1753.
- Bellinger EG, Sigeo DC. 2010. *Freshwater Algae Identification and Use as Bioindicators*. Chichester, John Wiley & Sons, Ltd.
- Bohnen H, Silva LS, Macedo VRM, Marcolin E. 2005. Ácidos orgânicos na solução de um gleissolo sob diferentes sistemas de cultivo com arroz irrigado. *Revista Brasileira de Ciência do Solo* 29: 475–480.
- Cassol APV, Oliveira MA, Figueiredo MCS, Luz DS, Sartori GMS, Marchesan E. 2013. Microalgas em Cultura de arroz: influência de diferentes manejos de adubação em áreas com residual de herbicidas (imidazolinonas). *Iheringia Série Botânica* 68: 261-271
- Cochard R, Maneepitak S, Kumar P. 2014. Aquatic faunal abundance and diversity in relation to synthetic and natural pesticide applications in rice fields of Central Thailand. *International Journal of Biodiversity Science, Ecosystem Services & Management* 10: 157–173.

- Das DN, Saikia SK, Das AK. 2007. Periphyton in rice–fish culture system: A case study from Arunachal Pradesh, India. *Renewable Agriculture Food System* 22: 316–319.
- Debenest T, Pinelli E, Coste M, *et al.* 2009. Sensitivity of freshwater periphytic diatoms to agricultural herbicides. *Aquatic Toxicology* 93: 11–17.
- Debenest T, Silvestre J, Coste M, Pinelli E. 2010. Effects of Pesticides on Freshwater Diatoms. In: Whitacre, DM (ed.). *Reviews of Environmental Contamination and Toxicology*. pp. 87-103.
- Dermiyati, Niswati A. 2014. Improving Biodiversity in Rice Paddy Fields to Promote Land Sustainability. In: Kaneko N, Yoshiura S, Kobayashi M (eds). *Sustainable Living with Environmental Risks*. Tokyo, Springer. pp. 45-55.
- Donato FF, Kemmerich M, Facco JF, *et al.* 2012. Simultaneous determination of pesticide and antibiotic residues at trace levels in water samples by SPE and LC-MS/MS. *Brazilian Journal of Analytical Chemistry* 7: 331-340.
- Du P, Wu X, Xu J, *et al.* 2018. Clomazone influence soil microbial community and soil nitrogen cycling. *Science of The Total Environment* 644: 475–485.
- Dufrêne M, Legendre P. 1997. Species Assemblages and Indicator Species: The Need for a Flexible Asymmetrical Approach. *Ecological Monographs* 67: 345-366.
- Fujita Y, Ohtsuka T. 2005. Diatoms from paddy fields in northern Laos. *Diatom* 21:71-89.
- Furtado RD, Luca SJ. 2003. Técnicas de cultivo de arroz irrigado: relação com a qualidade de água, protozoários e diversidade fitoplanctônica. *Revista Brasileira de Engenharia Agrícola e Ambiental* 7: 165–172.
- Gustavson K, Mahlenberg F, Schluter L. 2003. Effects of Exposure Duration of Herbicides on Natural Stream Periphyton Communities and Recovery. *Archives of Environmental Contamination and Toxicology* 45: 48–58.
- Hasenack H, Weber EJ, Boldrini II, Trevisan R. 2010. Mapa de sistemas ecológicos da ecorregião das savanas uruguaias em escala 1:500.000 ou superior e relatório técnico descrevendo insumos utilizados e metodologia de elaboração do mapa de sistemas ecológicos. Porto Alegre, UFRGS.
- Irisarri P, Gonnet S, Monza J. 2001. Cyanobacteria in Uruguayan rice fields: diversity, nitrogen fixing ability and tolerance to herbicides and combined nitrogen. *Journal of Biotechnology* 91:95–103.
- Jabusch TW, Tjeerdema RS. 2006. Photodegradation of Penoxsulam. *Journal of Agriculture and Food Chemistry* 54: 5958–5961.

- Junk WJ, Piedade MTF, Lourival R, et al. 2013. Brazilian wetlands: their definition, delineation, and classification for research, sustainable management and protection: Brazilian Wetlands. *Aquatic Conservation: Marine and Freshwater Ecosystem* 24:5–22.
- Katayama N, Osada Y, Mashiko M, et al. 2019. Organic farming and associated management practices benefit multiple wildlife taxa: A large-scale field study in rice paddy landscapes. *Journal of Applied Ecology* 56:1970-1981.
- Kumar A, Sahu R. 2012. Diversity of Algae (Chlorophyceae) in Paddy Fields of Lalgutwa Area, Ranchi, Jharkhand. *Journal of Applied Pharmaceutical Science* 2: 092-095.
- Larras F, Bouchez A, Rimet F, Montuelle B. 2012. Using Bioassays and Species Sensitivity Distributions to Assess Herbicide Toxicity towards Benthic Diatoms. *PLoS ONE* 7:44458. <https://doi.org/10.1371/journal.pone.0044458>.
- Liu Y, Zou G, Yuan Q, Huang W, Zhou W. 2020. Phytoplankton community characteristics in rice paddy fields under different nitrogen fertilizer applications. *Acta Physiologiae Plantarum* 42:33.
- Lobo E, Leighton G. 1986. Estructuras comunitarias de las fitocenosis planctonicas de los sistemas de desembocaduras de rios y esteros de la zona central de Chile. *Biologia Marina* 22: 1-29.
- Lobo EA, Callegaro VLM, Bender P. 2002. Utilização de algas diatomáceas epilíticas como indicadoras da qualidade da água em rios e arroios da Região Hidrográfica do Guaíba, RS, Brasil. Santa Cruz do Sul, EDUNISC.
- Maltchik L, Stenert C, Batzer DP. 2017. Can rice field management practices contribute to the conservation of species from natural wetlands? Lessons from Brazil. *Basic and Applied Ecology* 18: 50–56.
- Marchesan E, Zanella R, Avila LA, et al. 2007. Rice herbicide monitoring in two Brazilian rivers during the rice growing season. *Scientia Agricola* 64: 131–137.
- Matsubara CP, Maltchik L, Torgan LC. 2008. Diversity and Distribution of Algae in Wetlands of the Rio Grande do Sul, Brazil. *Neotropical Biology and Conservation* 7.
- Millennium Ecosystem Assessment. 2005. Ecosystems and human wellbeing: wetlands and water. <http://www.unep.org/maweb/documents/document.358.aspx.pdf>.
- Negoro K., Higashino M. 1986. Diatom vegetation of paddy fields in Japan. Report 1. Diatom vegetation of paddy fields in the vicinity of Sakurai City, Nara Prefecture. *Diatom* 2: 1-8.

- Noldin JA, Hermes LC, Fay EF, Eberhardt DS, Rossi MA. 2001. Persistência do herbicida clomazone no solo e na água quando aplicado na cultura do arroz irrigado, sistema pré-germinado. *Planta Daninha* 19:401–408.
- Ohtsuka T, Fujita Y. 2001. The diatom flora and its seasonal changes in a paddy field in Central Japan. *Nova Hedwigia* 73: 97-128.
- Oksanen J, Blanchet GF, Friendly M, et al. 2020. *Vegan: Community Ecology Package*. R package version 2:5-7. <https://CRAN.R-project.org/package=vegan>.
- Pappas JL, Stoermer EF. 1996. Quantitative method for determining a representative algal sample count. *Journal of Phycology* 32: 693-696.
- Prasanna R, Nayak S. 2007. Influence of diverse rice soil ecologies on cyanobacterial diversity and abundance. *Wetlands Ecology and Management*.15:127–134.
- R Development Core Team. 2021. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Ramos VG. 2012. *As Estratégias Sociais e Produtivas no Assentamento Santa Maria do Ibicuí, Manoel Viana – RS*. MSc Thesis, University of Santa Maria, Santa Maria, Brazil.
- Reck L, Reimche GB, Alves CR, Oliveita MA, Machado SL. 2018. Efeito Dos Herbicidas Imazapir e Imazapique na Comunidade Fitoplanctônica em Lavoura de Arroz Irrigado. *Iheringia Série Botânica* 73: 298–307.
- Reimche GB, Machado SLO, Oliveira MA, et al. 2015. Imazethapyr and imazapic, bispyribac-sodium and penoxsulam: Zooplankton and dissipation in subtropical rice paddy water. *Science of The Total Environment* 514: 68–76.
- Reynolds CS, Huszar V, Kruk C, Naselli-Flores L, Melo S. 2002. Towards a functional classification of the freshwater phytoplankton. *Journal of Plankton Research* 24: 417–428.
- Rimet F, Bouchez A. 2011. Use of diatom life-forms and ecological guilds to assess pesticide contamination in rivers: Lotic mesocosm approaches. *Ecological Indicators* 11: 489–499.
- Roger PA, Heong KL, Teng PS. 1991. Biodiversity and sustainability of wetland rice production: role and potential of microorganisms and invertebrates. In: Hawksworth D.L. (ed.) *The Biodiversity of Microorganisms and Invertebrates: Its Role in Sustainable Agriculture*. Wallingford, CAB International. pp. 117–136.
- Rumrich U, Lange-Bertalot H, Rumrich M. 2000. Diatomeen der Anden. *Iconographia diatomologica* 9: 1–649.

- Saikia S, Das D. 2011. Diversity and productivity (Chlorophyll-a and Biomass) of periphyton on natural and artificial substrates from wetland ecosystem. *Journal of Wetlands Ecology* 5: 1–9.
- Saikia SK, Das DN. 2014. Sustainable aquaculture: agro-ecological role of periphyton in ricefish farming. *Review of Aquaculture* 7: 172–186.
- Sartori GMS, Marchesan E, Luz DS, Cassol APV, et al. 2011. Manejo da adubação e seus efeitos na ocorrência de algas e na produtividade de arroz irrigado em áreas com residual de imidazolinonas. *Ciência Rural* 8:1323-1330.
- Senseman SA, Armbrust K. 2007. Weed Science Society of America. *Herbicide handbook*. Lawrence, Weed Science Society of America.
- Sheath RG, Wehr JD. 2003. Freshwater habitats of algae In: Wehr, JD. & RG.(eds.) *Freshwater Algae of North America. Ecology and classification*. Aquatic Ecological Series, Academic Press Elsevier, California, USA.
- Shivakumara LV, Pattar PV. 2015. Diversity of Phytoplanktons in Rice Fields of Davangere Taluk, Karnataka. *Journal of Marine Science Research and Development* 5: 3. doi: 10.4172/2155-9910.1000172.
- Sihi D, Dari B, Yan Z, et al. 2020. Assessment of Water Quality in Indo-Gangetic Plain of South-Eastern Asia under Organic vs. Conventional Rice Farming. *Water* 12: 960. <https://doi.org/10.3390/w12040960>.
- Silva DRO, Avila LA, Agostinetto D, et al. 2009. Monitoramento de agrotóxicos em águas superficiais de regiões orizícolas no sul do Brasil. *Ciência Rural* 39: 2383–2389.
- SOSBAI- Sociedade Sul-Brasileira de Arroz Irrigado. 2018. Arroz irrigado: recomendações técnicas da pesquisa para o Sul do Brasil / Sociedade Sul-Brasileira de Arroz Irrigado. Reunião Técnica da Cultura do Arroz Irrigado. Pelotas. Gráfica e editora Palloti.
- Sournia A. 1978. *Phytoplankton manual*. Paris, UNESCO.
- Suárez-Serrano A, Ibáñez C, Lacorte S, Barata C. 2010. Ecotoxicological effects of rice field waters on selected planktonic species: comparison between conventional and organic farming. *Ecotoxicology* 19: 1523–1535. Trobajo R, Sullivan MJ. 2010. Applied diatom studies in estuaries and shallow coastal environments. In: *The diatoms: applications for the environmental and the earth sciences*. (Smol, J.P. & Stoermer, E.F.). Cambridge, Cambridge University Press. pp. 309–323.
- Trobajo R, Clavero E, Chepurnov VA, et al. 2009. Morphological, genetic and mating diversity within the widespread bioindicator *Nitzschia palea* (Bacillariophyceae). *Phycologia* 48: 443–459.

- Utermöhl U. 1958. Zur Vervollkommnung der quantitativen Phytoplankton-Methodik. Mitteilungen der Internationalen Vereinigung für theoretische und angewandte. Limnologie 19: 1-38.
- Zanella R, Primel EG, Machado SLO, Gonçalves FF, Marchezan E. 2002. Monitoring of the herbicide clomazone in environmental water samples by solid-phase extraction and high-performance liquid chromatography with ultraviolet detection. Chromatographia 55: 573–577.

CONSIDERAÇÕES FINAIS

O presente estudo abordou uma investigação inédita para o Brasil avaliando a assembleia de diatomáceas e a comunidade de microalgas em condições de campo com ênfase no impacto de herbicidas da lavoura de arroz.

Considerando as três comunidades estudadas, as diatomáceas no solo forneceram respostas mais nítidas sobre a ação dos herbicidas através da variação da riqueza, composição e teratologias. A amostragem no sedimento abrangeu um período importante dos efeitos, detectando alterações imediatamente após a aplicação. Diferenças na riqueza de espécies entre os dois sistemas, também foram observadas no epifítton, provavelmente devido ao maior tempo de contato com os herbicidas.

Do ponto de vista prático, as microalgas participam da microbiota do solo garantindo processos importantes para a saúde do ambiente como a fixação de nitrogênio e do carbono, além do incremento dos agregados no solo. Considerando que a utilização de produtos químicos afeta a biodiversidade, estudos sobre as microalgas do solo são necessários para o que o conhecimento produzido possa ser utilizado como subsídio de práticas promotoras de diversidade, essencial para a manutenção dos referidos processos.

No atual cenário, o incentivo às práticas convencionais nos sistemas de cultivo de arroz coloca em risco a sustentabilidade da produção de alimentos. É necessário o incentivo de práticas sustentáveis para a incorporação de valores como sustentabilidade, saúde e questões sociais, além do lucro e rentabilidade nos sistemas de produção.

Finalizando, destaca-se a necessidade de avaliar a influência do sistema de plantio direto e pré-germinado na assembleia de diatomáceas em arroz irrigado. Os sistemas diferenciam-se principalmente no período de alagamento do solo, no qual condições mais úmidas são mantidas no sistema pré-germinado. Sugere-se também que assembleias de diatomáceas do solo sejam avaliadas em outros sistemas agrícolas e sob o impacto dos demais agrotóxicos utilizados atualmente.