

## SCIENTIFIC NOTE

### Record of *Podisus nigrispinus* (Dallas) (Hemiptera: Pentatomidae) Preying on *Metrogaleruca obscura* Degeer (Coleoptera: Chrysomelidae)

LA MOURA, J GRAZIA

Lab de Entomologia Sistemática, Depto de Zoologia, Univ Federal do Rio Grande do Sul, Porto Alegre, RS, Brasil

#### Keywords

*Cordia*, Heteroptera, black sage

#### Correspondence

JOCELIA GRAZIA, Lab de Entomologia Sistemática, Depto de Zoologia, Univ Federal do Rio Grande do Sul, Av. Bento Gonçalves 9500, Bloco IV, Prédio 43435, 91501-970, Porto Alegre, RS, Brasil; [jocelia@ufrgs.br](mailto:jocelia@ufrgs.br)

Edited by Madelaine Venzon – EPAMIG

Received on 25 October 2010 and accepted  
19 April 2011

#### Abstract

Asopines are predators of insects, with several species with potential as biocontrol agents of a number of pests. *Metrogaleruca obscura* (Degeer), a neotropical species of Galerucini, was introduced in Malaysia, Asia, and Mauritius, Africa, to control the spread of *Cordia curassavica* (Boraginaceae), a native plant of the neotropics. The occurrence of *Podisus nigrispinus* (Dallas) preying on *M. obscura* is recorded, and *Cordia verbenacea* (Boraginaceae) is mentioned as a host plant for *M. obscura*. A list of Chrysomelidae attacked by asopines in the neotropical region is also presented.

Many chrysomelids are important as agricultural pests or beneficial biocontrol agents of weeds (Jolivet 1997). *Metrogaleruca obscura* (Degeer) belongs to a group that includes genera of interest to agriculture, such as *Diabrotica*, *Acalymma* and *Cerotoma*. This species was also reported as *Schematiza cordiae* Barber (Wilcox 1971). It occurs in the West Indies, Venezuela, Suriname, French Guiana, Brazil (states of Amapá, Pará, Maranhão, Tocantins, Pernambuco, Bahia, Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo, Santa Catarina and Rio Grande do Sul), Bolivia, Paraguay, and Argentina (province of Salta) (Wilcox 1971, Moura 2008).

*Metrogaleruca obscura* was successfully introduced in Malaysia (Simmonds 1980), Mauritius Islands (= Mauritius) and Africa (Ung *et al* 1979, Denoth *et al* 2002, Greathead 2003) to reduce the spread of *Cordia curassavica* (Boraginaceae), a native plant of the neotropics.

Reports on predation of neotropical chrysomelids by pentatomid asopines are scarce and restricted to the subfamilies Chrysomelinae, Galerucinae, Cassidinae and Criocerinae (Table 1), which have individuals that

generally feed on host leaves. Most other subfamilies of Chrysomelidae have no records of asopines as predators, probably because their biology is poorly known. The information available is restricted to those species of economic interest of which larvae usually have subterranean and/or nocturnal habits or are protected by structures, such as a scotoshell or fecal shield (Cox 1996). They may present cicloalexia, a defensive behavior against predators in which larvae form a circle when resting to remain aggregated (Jolivet *et al* 1990).

Asopines prey mainly slow moving insects with soft body, primarily larvae of Lepidoptera (Medeiros *et al* 2003), Coleoptera and Hymenoptera. Specifically with regard to *M. obscura*, there are records of predation by *Heteroscelis servillei* Laporte (Hemiptera: Pentatomidae) in the West Indies (Callan 1948, Simmonds 1949, Herting 1973) and by *Afrius williamsi* Miller (Hemiptera: Pentatomidae) in Mauritius (Miller 1951, Tomas 1994).

This contribution aims to present a new record of predation of *M. obscura* and provide a list of chrysomelids that are preyed by asopines on the neotropics.

Specimens of *M. obscura* were collected in February

Table 1 Species of Chrysomelidae with their predators in the neotropics.

Subfamily/species	Predators	References
<b>Chrysomelinae</b>		
<i>Microtheca ochroloma</i> Stal	<i>Stiretrus decastigmus</i> (Herrich-Schaeffer)	Poncio <i>et al</i> (2010)
<i>Phaedon pertinax</i> Stal	<i>S. erythrocephalus</i> (Lepelletier & Serville)	Bondar (1953)
<i>Platyphora fasciatomaculata</i> (Stal)	<i>Stiretrus</i> sp.	Medeiros <i>et al</i> (1996)
<b>Cassidinae</b>		
<i>Acromis spinifex</i> (L.)	<i>S. decemguttatus</i> (Lep. & Serv.) <i>S. decastigmus</i> (Herrich-Schaeffer)	Bondar (1953)
<i>Botanochara sedecimpustulata</i> (Fabricius)	<i>S. decemguttatus</i> (Lep. & Serv.) <i>S. decastigmus</i> (Herrich-Schaeffer)	Bondar (1953)
<i>Zatrephina lineata</i> (Fabricius)	<i>S. decemguttatus</i> (Lep. & Serv.) <i>S. decastigmus</i> (Herrich-Schaeffer)	Bondar (1953)
<b>Galerucinae</b>		
<i>Cacoscelis marginata</i> (F.)	<i>Stiretrus</i> sp.	Lordello (1952)
<i>Coelomera lanio</i> (Dalman)	<i>Oplomus catena</i> (Drury)	Silveira <i>et al</i> (2002)
<i>Metrogaleruca obscura</i> (Degeer)	<i>Heteroscelis servillei</i> Laporte <i>Podisus nigrispinus</i> (Dallas)	Callan (1948), Simmonds (1949), Herting (1943) NEW RECORD
<i>Diabrotica speciosa</i> (Germar)	<i>Oplomus cruentus</i> (Burmeister)	Lima (1968)
<i>Pedilia</i> sp.	<i>P. aenescens</i> (Stal)	Duckett (1989)
<b>Criocerinae</b>		
<i>Lema apicalis</i> Lacordaire	<i>Podisus</i> sp.	Medeiros <i>et al</i> (1996)
<i>Lema reticulosa</i> Clark	<i>Podisus</i> sp.	Medeiros <i>et al</i> (1996)

2008 at the Centro Pluridisciplinar de Pesquisas Químicas, Biológicas e Agrícolas (CPQBA), Universidade Estadual de Campinas, Campinas, state of São Paulo, Brazil, defoliating leaves of *Cordia verbenacea* (Boraginaceae). This plant, originally described from the state of Rio de Janeiro, Brazil, is distributed in Brazil (from Rio de Janeiro to Rio Grande do Sul states), Bolivia, Paraguay, and Argentina. It is popularly known as black sage and widely used as a medicinal plant with anti-inflammatory action.

Two asopine species were observed preying on individuals of *M. obscura*: *H. servillei* and *Podisus nigrispinus* (Dallas), with the first record of the last species preying on this chrysomelid beetle.

The potential medical importance of *C. verbenacea* makes necessary to study predators of *M. obscura* to reduce or eliminate damage caused by herbivory. *Metrogaleruca obscura* has been registered only in *Cordia* species as host plants.

### Acknowledgments

To Centroflora (Botucatu, SP), for sending the material for identification. To Dr. A. Leyva for reviewing the English

version of the manuscript. LAM and JG acknowledge the grant support of “Conselho Nacional de Desenvolvimento Científico e Tecnológico” – CNPq.

### References

- Bondar G (1953) Pragas novas nas plantas do Brasil, II. Boletim Campo 9: 20-24.
- Callan EM (1948) The Pentatomidae, Cydnidae and Scutelleridae of Trinidad, B.W. I. Proc R Entomol Soc Lond Ser B Taxon 17: 115-124.
- Cox ML (1996) Insect predators of Chrysomelidae, p.23-91 In Jolivet P, Cox ML (eds) Chrysomelidae biology. Vol. 2. Ecological studies. Amsterdam, Academic Publishing, 465p.
- Denoth M, Frid L, Myers JH (2002) Multiple agents in biological control: improving the odds? Biol Control 24: 20-30.
- Duckett CN (1989) Natural history of *Pedilia* sp. A and its interactions with other herbivores of *Passiflora pittieri*. Entomography 6: 381-389.
- Greathead DJ (2003) Historical overview of biological control in Africa, p.1-26 In Neuenschwander P, Borgemeister C, Longewaldt J (eds) Biological control in IPM systems in Africa. Wellington, CAB International, 456p.

- Jolivet P (1997) Biologie des coléoptères Chrysomélides. Paris, Boubée, 279p.
- Jolivet P, Vasconcellos Neto J, Weinstein P (1990) Cycloalexy: a new concept in the larval defense of insects. *Insecta Mundi* 4: 133-142.
- Lima A da C (1968) Quarto catálogo dos insetos que vivem nas plantas do Brasil: seus parasitos e predadores. parte 2, n. 1, p.417-239. Rio de Janeiro, Ministério da Agricultura.
- Lordello LGE (1952) Insetos que vivem sobre o maracujazeiro. II. Contribuição ao conhecimento de *Cacoscelis famelica* (F., 1787) (Col. Chrysomelidae). *Dusenica* 3: 387-393.
- Medeiros L, Ferro DN, Mafra Neto A (1996) Association of chrysomelid beetles with solanaceous plants in the south of Brazil, p.339-363 In Jolivet PHA, Cox ML (eds) Chrysomelidae biology, vol. 2: Ecological studies. Amsterdam, SPB Academic, 465p.
- Medeiros RS, Ramalho FS, Zanuncio JC, Serrão JE (2003) Effect of temperature on life table parameters of *Podisus nigrispinus* (Het., Pentatomidae) fed with *Alabama argillacea* (Lep., Noctuidae) larvae. *J Appl Entomol* 127: 209-213.
- Miller NCE (1951) A new species of *Afrius* (Hem., Pentatomidae) predacious on *Schematiza cordiae* Barb., in Mauritius. *Bull Entomol Res* 42: 183-184.
- Moura L de A (2008) Análise cladística dos gêneros da Seção Schematizites (Coleoptera, Chrysomelidae, Galerucinae, Galerucini). Tese de doutorado, Porto Alegre, Universidade Federal do Rio Grande do Sul, 105p.
- Poncio S, Dequech STB, Sturza VS, Lissner RAD, Perlin LF, Rosalino PK, Ribeiro L do P (2010) Primeiro relato de *Stiretrus decastigmus* no Brasil predando *Microtheca ochroloma*. *Ci Ru* 40: 1203-1205.
- Silveira RD, Anjos N dos, Zanuncio JC (2002) Natural enemies of *Coelomera lanio* (Coleoptera: Chrysomelidae) in the region of Viçosa, Minas Gerais, Brazil. *Rev Biol Trop* 50: 117-120.
- Simmonds FG (1980) Biological control of *Cordia curassavica* (Boraginaceae) in Malaysia. *Entomophaga* 25: 363-364.
- Simmonds FJ (1949) Insects attacking *Cordia macrostachya* (Jacq.) Roem. and Schult. in the West Indies. II. *Schematiza cordiae* Barb. (Coleoptera, Galerucidae). *Can Entomol* 81: 275-282.
- Thomas DB (1994) Taxonomic synopsis of the Old World asopine genera (Heteroptera: Pentatomidae). *Insecta Mundi* 8: 145-212.
- Ung SH, Yunus A, Chin WH (1979) Biological control of *Cordia curassavica* in Malaysia by *Schematiza cordiae* Barb. *Malay Agric J* 52: 154-165.
- Wilcox JA (1971) Coleopterorum catalogus. Chrysomelidae: Galerucinae, 2 ed., s'-Gravenhage, W. Junk, v.78, n. 1, 220p.
-